

wwPDB X-ray Structure Validation Summary Report (i)

Jul 10, 2024 – 01:27 am BST

PDB ID : 9FE8

Title : Crystal Structure of reduced NuoEF variant P228R(NuoF) from Aquifex ae-

olicus

Authors: Wohlwend, D.; Friedrich, T.; Goeppert-Asadollahpour, S.

Deposited on : 2024-05-17

Resolution : 2.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.37.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Percentile relative to X-ray structures of similar resolution

Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(\mathring{A}))$
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	160	8% 85%	12% •
1	С	160	93%	6% •
2	В	434	90%	6% •
2	D	434	89%	7% •



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 9829 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called NADH-quinone oxidoreductase subunit E.

\mathbf{Mol}	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	10001	С	- '	0	S	0	0	0
			1268	821	204	234	9	_	-	
1	C	159	Total	С	N	O	S	0	1	0
1		109	1296	837	211	239	9	U		

• Molecule 2 is a protein called NADH-quinone oxidoreductase subunit F.

\mathbf{Mol}	Chain	Residues		Atoms					AltConf	Trace
2	B	418	Total	С	N	О	S	0	1	0
2	Ъ	410	3298	2118	551	616	13	U	1	
2	D	418	Total	С	N	Ο	S	0	9	0
2	ש	410	3323	2136	555	619	13	0)	

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	228	ARG	PRO	engineered mutation	UNP O66841
В	427	ALA	-	expression tag	UNP O66841
В	428	GLY	-	expression tag	UNP O66841
В	429	HIS	-	expression tag	UNP O66841
В	430	HIS	- expression tag		UNP O66841
В	431	HIS	-	expression tag	UNP O66841
В	432	HIS	-	expression tag	UNP O66841
В	433	HIS	_	expression tag	UNP O66841
В	434	HIS	-	expression tag	UNP O66841
D	228	ARG	PRO	engineered mutation	UNP O66841
D	427	ALA	_	expression tag	UNP O66841
D	428	GLY	-	expression tag	UNP O66841
D	429	HIS	-	expression tag	UNP O66841
D	430	HIS	-	expression tag	UNP O66841
D	431	HIS	-	expression tag	UNP O66841
D	432	HIS	_	expression tag	UNP O66841


Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	433	HIS	-	expression tag	UNP O66841
D	434	HIS	-	expression tag	UNP O66841

• Molecule 3 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Fe S 4 2 2	0	0
3	С	1	Total Fe S 4 2 2	0	0

 \bullet Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	В	1	Total O S 5 4 1	0	0
4	С	1	Total O S 5 4 1	0	0
4	С	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0

• Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total Na 3 3	0	0
5	В	3	Total Na 3 3	0	0
5	С	1	Total Na 1 1	0	0
5	D	1	Total Na 1 1	0	0


• Molecule 6 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe_4S_4) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	1	Total Fe S 8 4 4	0	0
6	D	1	Total Fe S 8 4 4	0	0

• Molecule 7 is 1-DEOXY-1-(7,8-DIMETHYL-2,4-DIOXO-3,4-DIHYDRO-2H-BENZO[G]P TERIDIN-1-ID-10(5H)-YL)-5-O-PHOSPHONATO-D-RIBITOL (three-letter code: FNR) (formula: $C_{17}H_{23}N_4O_9P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	D	1	Total	С	N	О	Р	0	0
'	Б	1	31	17	4	9	1	U	0


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	D	1	Total	С	N	О	Р	0	0
'	D	1	31	17	4	9	1	U	0


• Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf							
8	В	1	Total C O	0	0							
	Б	1	6 3 3	· ·	Ů							
Q	В	1	Total C O	0	0							
0	Ъ	Б	Ъ	Ъ	Б	Б	D	Б	1	6 3 3	U	U
Q	В	1	Total C O	0	0							
	Ъ	D 1	6 3 3	U								
0	D	1	Total C O	0	0							
0	D	1	6 3 3	U	U							
R	D	1	Total C O	0	0							
	ע	1	6 3 3	U								

• Molecule 9 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (three-letter code: MPO) (formula: $C_7H_{15}NO_4S$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
0	D	1	Total	С	N	О	S	0	0
	ש	1	13	7	1	4	1	0	

• Molecule 10 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	52	Total O 53 53	0	1
10	В	198	Total O 198 198	0	0
10	С	55	Total O 55 55	0	0
10	D	171	Total O 171 171	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NADH-quinone oxidoreductase subunit E





4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 21 21 21	Depositor	
Cell constants	63.08Å 116.22Å 190.26Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	49.64 - 2.34	Depositor	
recontion (11)	49.59 - 2.34	EDS	
% Data completeness	99.4 (49.64-2.34)	Depositor	
(in resolution range)	99.4 (49.59-2.34)	EDS	
R_{merge}	(Not available)	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	2.20 (at 2.34Å)	Xtriage	
Refinement program	REFMAC 5.8.0425	Depositor	
D D.	0.184 , 0.206	Depositor	
R, R_{free}	0.190 , 0.209	DCC	
R_{free} test set	2968 reflections (4.98%)	wwPDB-VP	
Wilson B-factor (Å ²)	38.2	Xtriage	
Anisotropy	0.290	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 29.7	EDS	
L-test for twinning ²	$ < L > = 0.51, < L^2> = 0.34$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
F_o, F_c correlation	0.96	EDS	
Total number of atoms	9829	wwPDB-VP	
Average B, all atoms (Å ²)	45.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 43.85 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6611e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MPO, GOL, FNR, SO4, FES, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.35	0/1297	0.68	0/1752	
1	С	0.37	0/1325	0.72	0/1790	
2	В	0.39	0/3378	0.71	0/4571	
2	D	0.38	0/3404	0.68	1/4603 (0.0%)	
All	All	0.38	0/9404	0.70	$1/12716 \ (0.0\%)$	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	В	0	1
2	D	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

N	Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
	2	D	18	MET	CG-SD-CE	5.04	108.27	100.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	В	401	ARG	Sidechain
2	D	87	ARG	Sidechain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1268	0	1269	17	0
1	С	1296	0	1292	5	0
2	В	3298	0	3259	18	0
2	D	3323	0	3286	17	0
3	A	4	0	0	0	0
3	С	4	0	0	0	0
4	A	10	0	0	0	0
4	В	5	0	0	0	0
4	С	10	0	0	0	0
4	D	5	0	0	0	0
5	A	3	0	0	0	0
5	В	3	0	0	0	0
5	С	1	0	0	0	0
5	D	1	0	0	0	0
6	В	8	0	0	0	0
6	D	8	0	0	1	0
7	В	31	0	22	0	0
7	D	31	0	22	0	0
8	В	18	0	24	2	0
8	D	12	0	16	0	0
9	D	13	0	15	0	0
10	A	53	0	0	1	0
10	В	198	0	0	1	0
10	С	55	0	0	0	0
10	D	171	0	0	1	0
All	All	9829	0	9205	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1 Atom-2		$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)	
2:B:384:ASN:HD22	2:B:404:ARG:HH21	1.06	1.00	
1:A:38:GLN:HE22	1:A:74:ASP:H	1.31	0.77	

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	Clash overlap (Å)
2:B:68:GLY:HA3	8:B:504:GOL:H31	1.75	0.68
2:B:384:ASN:ND2	2:B:404:ARG:HH21	1.87	0.67
2:B:233:MET:HE1	2:B:241:ILE:HD11	1.78	0.66

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	154/160 (96%)	149 (97%)	5 (3%)	0	100	100
1	С	158/160 (99%)	154 (98%)	4 (2%)	0	100	100
2	В	417/434 (96%)	405 (97%)	11 (3%)	1 (0%)	47	55
2	D	419/434 (96%)	406 (97%)	12 (3%)	1 (0%)	47	55
All	All	1148/1188 (97%)	1114 (97%)	32 (3%)	2 (0%)	47	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	95	GLU
2	В	95	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Pe	erce	ntiles
1	A	142/146~(97%)	142 (100%)	0	10	00	100
1	С	144/146 (99%)	142 (99%)	2 (1%)		67	78
2	В	342/357~(96%)	340 (99%)	2 (1%)	8	86	92
2	D	344/357 (96%)	341 (99%)	3 (1%)	,	78	87
All	All	972/1006 (97%)	965 (99%)	7 (1%)	9	84	90

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	116	PRO
2	D	2	ARG
2	D	393	CYS
2	D	9	ARG
1	С	43	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	220	ASN
2	D	286	ASN
2	D	362	ASN
2	D	344	HIS
2	В	220	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 26 ligands modelled in this entry, 8 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Во	ond leng	ths	В	ond ang	eles
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	С	203	-	4,4,4	0.32	0	6,6,6	0.28	0
4	SO4	A	202	5	4,4,4	0.35	0	6,6,6	0.12	0
7	FNR	В	502	-	32,33,33	0.37	0	40,50,50	0.61	0
7	FNR	D	502	-	32,33,33	0.37	0	40,50,50	0.57	0
8	GOL	В	503	-	5,5,5	0.16	0	5,5,5	0.28	0
4	SO4	A	203	-	4,4,4	0.33	0	6,6,6	0.11	0
3	FES	С	201	1	0,4,4	-	-	-		
6	SF4	В	501	2	0,12,12	-	-	-		
8	GOL	В	504	-	5,5,5	0.18	0	5,5,5	0.57	0
4	SO4	В	506	-	4,4,4	0.33	0	6,6,6	0.08	0
8	GOL	В	505	-	5,5,5	0.13	0	5,5,5	0.35	0
9	MPO	D	503	-	13,13,13	0.73	1 (7%)	17,17,17	0.76	0
6	SF4	D	501	2	0,12,12	-	-	-		
8	GOL	D	504	-	5,5,5	0.10	0	5,5,5	0.35	0
4	SO4	D	506	-	4,4,4	0.29	0	6,6,6	0.16	0
4	SO4	С	202	-	4,4,4	0.32	0	6,6,6	0.08	0
8	GOL	D	505	-	5,5,5	0.10	0	5,5,5	0.29	0
3	FES	A	201	1	0,4,4	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	FNR	В	502	-	-	4/18/18/18	0/3/3/3
7	FNR	D	502	-	-	3/18/18/18	0/3/3/3
3	FES	С	201	1	-	-	0/1/1/1
8	GOL	В	505	-	-	0/4/4/4	-
8	GOL	В	504	-	-	2/4/4/4	-
9	MPO	D	503	-	-	0/7/15/15	0/1/1/1

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SF4	В	501	2	-	-	0/6/5/5
6	SF4	D	501	2	-	-	0/6/5/5
8	GOL	D	504	_	-	0/4/4/4	-
8	GOL	В	503	-	-	2/4/4/4	-
8	GOL	D	505	-	-	0/4/4/4	-
3	FES	A	201	1	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
9	D	503	MPO	O3-S1	2.57	1.56	1.47

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	В	504	GOL	C1-C2-C3-O3
8	В	504	GOL	O2-C2-C3-O3
7	D	502	FNR	C4'-C5'-O5'-P
7	В	502	FNR	C4'-C5'-O5'-P
8	В	503	GOL	O1-C1-C2-C3

There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	В	503	GOL	1	0
8	В	504	GOL	1	0
6	D	501	SF4	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.




The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	156/160 (97%)	0.32	12 (7%) 13 20	33, 56, 79, 112	0
1	С	159/160 (99%)	0.05	6 (3%) 40 51	30, 46, 75, 115	0
2	В	418/434 (96%)	-0.26	1 (0%) 95 97	29, 40, 62, 90	0
2	D	418/434 (96%)	-0.35	2 (0%) 91 95	31, 41, 59, 95	0
All	All	1151/1188 (96%)	-0.17	21 (1%) 68 76	29, 42, 68, 115	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	6	PHE	5.0
1	С	2	PHE	4.7
1	С	6	PHE	4.2
1	A	5	GLU	3.9
1	A	10	GLU	3.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
8	GOL	В	504	6/6	0.79	0.23	72,81,85,86	0
8	GOL	В	505	6/6	0.83	0.15	64,75,79,80	0
4	SO4	В	506	5/5	0.88	0.28	94,97,102,106	0
5	NA	В	509	1/1	0.89	0.35	59,59,59,59	0
5	NA	A	204	1/1	0.89	0.23	57,57,57	0
5	NA	A	206	1/1	0.89	0.13	47,47,47,47	0
8	GOL	В	503	6/6	0.90	0.17	46,50,53,54	0
5	NA	В	508	1/1	0.91	0.42	58,58,58,58	0
8	GOL	D	504	6/6	0.91	0.17	55,62,68,76	0
8	GOL	D	505	6/6	0.91	0.18	68,69,72,74	0
5	NA	A	205	1/1	0.92	0.22	59,59,59,59	0
9	MPO	D	503	13/13	0.92	0.23	58,78,91,94	0
4	SO4	A	203	5/5	0.94	0.16	74,76,79,80	0
4	SO4	A	202	5/5	0.94	0.16	78,87,97,99	0
5	NA	С	204	1/1	0.94	0.53	53,53,53,53	0
5	NA	D	507	1/1	0.94	0.25	43,43,43,43	0
4	SO4	С	202	5/5	0.94	0.22	75,77,88,90	0
4	SO4	С	203	5/5	0.95	0.19	59,67,71,74	0
5	NA	В	507	1/1	0.95	0.11	45,45,45,45	0
4	SO4	D	506	5/5	0.95	0.11	57,62,71,81	0
7	FNR	D	502	31/31	0.97	0.12	30,33,34,36	0
7	FNR	В	502	31/31	0.98	0.10	28,32,34,37	0
6	SF4	В	501	8/8	0.99	0.10	33,34,35,36	0
6	SF4	D	501	8/8	0.99	0.12	31,33,34,34	0
3	FES	A	201	4/4	1.00	0.10	33,33,33,35	0
3	FES	С	201	4/4	1.00	0.12	32,32,32,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.












Electron density around SF4 B 501:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around SF4 D 501:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around FES A 201:


 $2mF_o$ -DF_c (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

