



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 11, 2023 – 08:35 am GMT

PDB ID : 5FJN
Title : Structure of L-Amino acid deaminase from *Proteus myxofaciens* in complex with anthranilate
Authors : Motta, P.; Molla, G.; Pollegioni, L.; Nardini, M.
Deposited on : 2015-10-11
Resolution : 1.75 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	456	
1	B	456	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 7620 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-AMINO ACID DEAMINASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	447	Total 3464	2202	592	659	11	0	4	0
1	B	447	Total 3480	2214	592	663	11	0	7	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 53	27	9	15	2	0	0
2	B	1	Total 53	27	9	15	2	0	0

- Molecule 3 is 2-AMINOBENZOIC ACID (three-letter code: BE2) (formula: $C_7H_7NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			10	7	1	2		
3	B	1	Total	C	N	O	0	0
			10	7	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	263	Total	O	0	4
			267	267		
4	B	279	Total	O	0	4
			283	283		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-AMINO ACID DEAMINASE



- Molecule 1: L-AMINO ACID DEAMINASE



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.81Å 93.32Å 75.07Å 90.00° 103.22° 90.00°	Depositor
Resolution (Å)	41.19 – 1.75 39.33 – 1.75	Depositor EDS
% Data completeness (in resolution range)	97.9 (41.19-1.75) 97.9 (39.33-1.75)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.156 , 0.197 0.166 , 0.204	Depositor DCC
R_{free} test set	4818 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.6	Xtrriage
Anisotropy	0.060	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.022 for l,-k,h	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7620	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, BE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3550	0.52	0/4811
1	B	0.30	0/3575	0.54	0/4845
All	All	0.30	0/7125	0.53	0/9656

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3464	0	3473	8	0
1	B	3480	0	3494	6	0
2	A	53	0	31	4	0
2	B	53	0	31	3	0
3	A	10	0	3	0	0
3	B	10	0	3	0	0
4	A	267	0	0	0	0
4	B	283	0	0	1	0
All	All	7620	0	7035	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:ASN:O	1:A:184:VAL:HG13	1.90	0.71
1:B:412:VAL:HG13	2:B:500:FAD:C8M	2.31	0.60
1:A:412:VAL:HG13	2:A:500:FAD:C8M	2.33	0.59
1:A:128[A]:ILE:HG13	1:A:214:TYR:CG	2.40	0.57
1:A:98:TYR:H	2:A:500:FAD:C6	2.20	0.53
1:B:412:VAL:HG13	2:B:500:FAD:HM83	1.90	0.53
1:A:59:VAL:HG11	1:A:241:VAL:HG21	1.92	0.51
1:B:98:TYR:H	2:B:500:FAD:C6	2.26	0.49
1:A:412:VAL:HG13	2:A:500:FAD:HM81	1.96	0.47
1:B:172:ARG:HD2	4:B:2148:HOH:O	2.17	0.44
1:B:318:PHE:HB2	1:B:344:LEU:HD12	2.00	0.43
1:A:412:VAL:HG13	2:A:500:FAD:HM83	2.00	0.43
1:A:348:ILE:HD11	1:A:352:LEU:HD22	2.03	0.40
1:B:244:GLU:HG3	1:B:245:LYS:HG3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	449/456 (98%)	434 (97%)	15 (3%)	0	100	100
1	B	452/456 (99%)	436 (96%)	16 (4%)	0	100	100
All	All	901/912 (99%)	870 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/377 (99%)	372 (100%)	1 (0%)	92	89
1	B	376/377 (100%)	376 (100%)	0	100	100
All	All	749/754 (99%)	748 (100%)	1 (0%)	93	91

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	302	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	500	-	53,58,58	1.32	7 (13%)	68,89,89	1.34	10 (14%)
3	BE2	A	1476	-	10,10,10	1.87	1 (10%)	13,13,13	0.67	0
3	BE2	B	1476	-	10,10,10	1.88	1 (10%)	13,13,13	0.62	0
2	FAD	B	500	-	53,58,58	1.33	6 (11%)	68,89,89	1.36	8 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	500	-	-	1/30/50/50	0/6/6/6
3	BE2	A	1476	-	-	0/4/4/4	0/1/1/1
3	BE2	B	1476	-	-	0/4/4/4	0/1/1/1
2	FAD	B	500	-	-	1/30/50/50	0/6/6/6

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1476	BE2	C1-CA	5.53	1.49	1.41
3	B	1476	BE2	C1-CA	5.53	1.49	1.41
2	A	500	FAD	C9A-C5X	5.12	1.49	1.41
2	B	500	FAD	C9A-C5X	4.94	1.49	1.41
2	B	500	FAD	C4X-N5	3.32	1.37	1.30
2	A	500	FAD	C8-C7	3.14	1.48	1.40
2	B	500	FAD	C1'-C2'	-3.03	1.48	1.52
2	B	500	FAD	C8-C7	3.00	1.48	1.40
2	A	500	FAD	C4X-N5	2.82	1.36	1.30
2	A	500	FAD	C5A-C4A	2.34	1.47	1.40
2	A	500	FAD	C1'-C2'	-2.31	1.49	1.52
2	B	500	FAD	C5A-C4A	2.22	1.46	1.40
2	A	500	FAD	C4-N3	-2.12	1.34	1.38
2	A	500	FAD	C2A-N3A	2.06	1.35	1.32
2	B	500	FAD	C4-N3	-2.03	1.35	1.38

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	FAD	N3A-C2A-N1A	-3.71	122.89	128.68
2	B	500	FAD	N3A-C2A-N1A	-3.62	123.02	128.68
2	A	500	FAD	O4-C4-C4X	-3.25	117.98	126.60
2	B	500	FAD	C4-C4X-N5	3.23	122.83	118.23
2	B	500	FAD	C4'-C3'-C2'	-3.02	107.08	113.36
2	B	500	FAD	O4-C4-C4X	-2.91	118.88	126.60
2	A	500	FAD	C4-C4X-N5	2.56	121.88	118.23
2	A	500	FAD	C4A-C5A-N7A	-2.48	106.81	109.40
2	B	500	FAD	O2-C2-N1	-2.47	117.73	121.83
2	A	500	FAD	C4X-C4-N3	2.41	119.32	113.19
2	B	500	FAD	C4X-C4-N3	2.39	119.26	113.19
2	A	500	FAD	O2-C2-N1	-2.33	117.96	121.83
2	A	500	FAD	C4X-C10-N1	-2.25	119.51	124.73
2	A	500	FAD	C4'-C3'-C2'	-2.23	108.73	113.36
2	A	500	FAD	C10-N1-C2	2.08	121.05	116.90
2	B	500	FAD	C4X-C10-N1	-2.06	119.95	124.73
2	A	500	FAD	C4-N3-C2	-2.06	121.84	125.64
2	B	500	FAD	C4A-C5A-N7A	-2.04	107.27	109.40

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	FAD	O4B-C4B-C5B-O5B
2	B	500	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.


2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	FAD	4	0
2	B	500	FAD	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	447/456 (98%)	0.36	40 (8%) 9 12	22, 35, 61, 113	0
1	B	447/456 (98%)	0.44	37 (8%) 11 14	20, 31, 55, 113	0
All	All	894/912 (98%)	0.40	77 (8%) 10 13	20, 33, 60, 113	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	337	LEU	20.4
1	B	337	LEU	14.7
1	B	28	MET	14.6
1	B	342	LEU	11.2
1	B	338	GLY	10.9
1	A	341	GLU	9.5
1	A	342	LEU	9.3
1	A	338	GLY	8.6
1	B	474	LYS	8.6
1	B	341	GLU	8.5
1	A	474	LYS	7.8
1	B	340	GLY	7.4
1	A	235	GLY	7.1
1	A	29	VAL	6.6
1	B	29	VAL	6.4
1	B	31	ARG	5.8
1	B	32	ASP	5.6
1	A	28	MET	5.6
1	A	339	GLY	5.5
1	A	340	GLY	5.2
1	B	336	LEU	4.9
1	A	336	LEU	4.8
1	B	353	PHE	4.2
1	A	353	PHE	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	176	GLY	3.8
1	A	318	PHE	3.7
1	B	97	ALA	3.5
1	B	363	LYS	3.5
1	A	189	PRO	3.5
1	A	398	VAL	3.5
1	B	318	PHE	3.5
1	B	365	ASP	3.4
1	A	146	GLU	3.3
1	A	144	ALA	3.2
1	B	202	VAL	3.1
1	A	150	ASP	3.0
1	B	30	ARG	3.0
1	A	344	LEU	3.0
1	A	473	LYS	2.9
1	B	339	GLY	2.9
1	B	473	LYS	2.9
1	A	177	GLU	2.9
1	A	403	GLN	2.7
1	B	65[A]	ILE	2.7
1	B	161	GLU	2.7
1	B	367	LYS	2.7
1	B	366	GLU	2.6
1	B	335	HIS	2.6
1	A	394	THR	2.6
1	B	334	MET	2.6
1	A	175	LYS	2.5
1	A	166	ASP	2.5
1	A	97	ALA	2.5
1	A	397	PRO	2.5
1	B	412	VAL	2.4
1	B	102	ILE	2.4
1	A	334	MET	2.4
1	B	101	ILE	2.4
1	A	102	ILE	2.3
1	B	438	VAL	2.3
1	A	153	GLN	2.3
1	B	383	GLU	2.3
1	B	279	LEU	2.2
1	B	166	ASP	2.2
1	A	335	HIS	2.2
1	A	350	GLU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	401	LYS	2.2
1	A	191	THR	2.2
1	B	357	LYS	2.1
1	B	182[A]	ARG	2.1
1	A	159	ALA	2.1
1	A	163	ALA	2.1
1	A	31	ARG	2.1
1	A	187	GLN	2.0
1	B	157	LYS	2.0
1	A	365	ASP	2.0
1	B	439	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	BE2	A	1476	10/10	0.90	0.12	41,45,49,55	0
3	BE2	B	1476	10/10	0.92	0.13	31,38,41,41	0
2	FAD	A	500	53/53	0.97	0.10	21,23,27,31	0
2	FAD	B	500	53/53	0.97	0.15	19,21,24,26	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 500:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.