



# wwPDB EM Validation Summary Report ⓘ

Dec 30, 2024 – 09:19 PM EST

PDB ID : 8FJK  
EMDB ID : EMD-29243  
Title : Golden Shiner Reovirus Core Polar Vertex  
Authors : Stevens, A.S.; Zhou, Z.H.  
Deposited on : 2022-12-19  
Resolution : 3.30 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev113  
MolProbity : 4.02b-467  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





























Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415
RNA backbone	6643	2191

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1273	
2	B	718	
3	C	1138	
3	D	1138	
3	E	1138	
3	F	1138	
3	G	1138	
3	H	1138	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	I	1138	 93% 6%
3	J	1138	 99%
3	K	1138	 96%
3	L	1138	 99%
4	M	94	 98%
4	N	94	 32% 68%
4	k	94	 32% 68%
4	l	94	 32% 68%
4	m	94	 32% 68%
5	V	411	 99%
5	W	411	 100%
5	X	411	 99%
5	a	411	 99%
5	b	411	 100%
5	d	411	 100%
5	e	411	 100%
5	g	411	 99%
5	h	411	 99%
5	n	411	 100%
6	Y	1297	 99%
6	Z	1297	 99%
6	c	1297	 99%
6	f	1297	 99%
6	i	1297	 99%
7	a1	52	 98%
			 100%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
7	b1	52	100%
8	a2	60	100%
8	b2	60	100%
9	a3	40	100%
9	b3	40	100%
10	a4	18	100%
10	b4	18	100%
11	a5	38	100%
11	b5	38	100%
12	a6	30	100%
12	b6	30	100%

## 2 Entry composition [i](#)

There are 13 unique types of molecules in this entry. The entry contains 193908 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase VP2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1273	9978	6375	1736	1822	45	0	0

- Molecule 2 is a protein called Microtubule-associated protein VP5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	718	5593	3588	971	1016	18	0	0

- Molecule 3 is a protein called Major inner capsid protein VP3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	1075	8300	5296	1421	1534	49	0	0
3	D	1138	8761	5578	1502	1628	53	0	0
3	E	1106	8534	5440	1463	1579	52	0	0
3	F	1138	8761	5578	1502	1628	53	0	0
3	G	1098	8477	5403	1454	1568	52	0	0
3	H	1130	8699	5540	1491	1615	53	0	0
3	I	1065	8226	5244	1412	1519	51	0	0
3	J	1130	8699	5540	1491	1615	53	0	0
3	K	1098	8477	5403	1454	1568	52	0	0
3	L	1138	8761	5578	1502	1628	53	0	0

- Molecule 4 is a protein called Major inner capsid protein VP3.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	M	94	Total	C	N	O	S	0	0
			640	377	113	149	1		
4	N	30	Total	C	N	O	S	0	0
			210	128	36	45	1		
4	k	30	Total	C	N	O	S	0	0
			211	128	36	46	1		
4	l	30	Total	C	N	O	S	0	0
			211	128	36	46	1		
4	m	30	Total	C	N	O	S	0	0
			211	128	36	46	1		

- Molecule 5 is a protein called Clamp protein VP6.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	V	411	Total	C	N	O	S	0	0
			3138	2008	544	570	16		
5	W	411	Total	C	N	O	S	0	0
			3138	2008	544	570	16		
5	X	411	Total	C	N	O	S	0	0
			3138	2008	544	570	16		
5	a	411	Total	C	N	O	S	0	0
			3138	2008	544	570	16		
5	b	411	Total	C	N	O	S	0	0
			3138	2008	544	570	16		
5	d	411	Total	C	N	O	S	0	0
			3138	2008	544	570	16		
5	e	411	Total	C	N	O	S	0	0
			3138	2008	544	570	16		
5	g	411	Total	C	N	O	S	0	0
			3138	2008	544	570	16		
5	h	411	Total	C	N	O	S	0	0
			3138	2008	544	570	16		
5	n	411	Total	C	N	O	S	0	0
			3138	2008	544	570	16		

- Molecule 6 is a protein called Outer capsid protein VP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	Y	1297	Total	C	N	O	S	0	0
			9963	6382	1694	1860	27		
6	Z	1297	Total	C	N	O	S	0	0
			9963	6382	1694	1860	27		
6	c	1297	Total	C	N	O	S	0	0
			9963	6382	1694	1860	27		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
6	f	1297	Total	C	N	O	S	0	0
			9963	6382	1694	1860	27		
6	i	1297	Total	C	N	O	S	0	0
			9963	6382	1694	1860	27		

- Molecule 7 is a RNA chain called RNA (52-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
7	a1	52	Total	C	N	O	P	0	0
			1089	494	182	362	51		
7	b1	52	Total	C	N	O	P	0	0
			1089	494	182	362	51		

- Molecule 8 is a RNA chain called RNA (60-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
8	a2	60	Total	C	N	O	P	0	0
			1257	570	210	418	59		
8	b2	60	Total	C	N	O	P	0	0
			1257	570	210	418	59		

- Molecule 9 is a RNA chain called RNA (40-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
9	a3	40	Total	C	N	O	P	0	0
			837	380	140	278	39		
9	b3	40	Total	C	N	O	P	0	0
			837	380	140	278	39		

- Molecule 10 is a RNA chain called RNA (18-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
10	a4	18	Total	C	N	O	P	0	0
			375	171	63	124	17		
10	b4	18	Total	C	N	O	P	0	0
			375	171	63	124	17		

- Molecule 11 is a RNA chain called RNA (38-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
11	a5	38	Total	C	N	O	P	0	0
			795	361	133	264	37		
11	b5	38	Total	C	N	O	P	0	0
			795	361	133	264	37		

- Molecule 12 is a RNA chain called RNA (30-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
12	a6	30	Total	C	N	O	P	0	0
			627	285	105	208	29		
12	b6	30	Total	C	N	O	P	0	0
			627	285	105	208	29		

- Molecule 13 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
13	E	1	Total	Zn	0
			1	1	
13	G	1	Total	Zn	0
			1	1	
13	I	1	Total	Zn	0
			1	1	
13	K	1	Total	Zn	0
			1	1	




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA-directed RNA polymerase VP2



- Molecule 2: Microtubule-associated protein VP5




- Molecule 3: Major inner capsid protein VP3




- Molecule 3: Major inner capsid protein VP3







• Molecule 3: Major inner capsid protein VP3




• Molecule 3: Major inner capsid protein VP3





• Molecule 3: Major inner capsid protein VP3




• Molecule 3: Major inner capsid protein VP3




• Molecule 3: Major inner capsid protein VP3







• Molecule 3: Major inner capsid protein VP3




• Molecule 3: Major inner capsid protein VP3





• Molecule 3: Major inner capsid protein VP3



• Molecule 4: Major inner capsid protein VP3




• Molecule 4: Major inner capsid protein VP3



• Molecule 4: Major inner capsid protein VP3




THR	ALA	SER	PRO	ALA	ASP	THR	ASN	VAL	VAL	PRO	PRO	LYS	ASP	ALA	PRO	THR	THR	ASN	SER	PRO	PRO	SER	SER	THR	THR	SER	PRO	ASN	GLN	GLN	ALA	ALA	ALA	ASP	ASN	GLN	GLN	GLN	ALA	ALA	ILE	VAL	SER	SER	GLN	GLY	ASN	ASN	ALA	VAL	GLY	ASP	SER	ALA	PRO	SER	THR	SER	VAL
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----



• Molecule 4: Major inner capsid protein VP3




THR	ALA	SER	PRO	ALA	ASP	THR	ASN	VAL	VAL	PRO	PRO	LYS	ASP	ALA	PRO	THR	THR	ASN	SER	PRO	PRO	SER	SER	THR	THR	SER	PRO	ASN	GLN	GLN	ALA	ALA	ALA	ASP	ASN	GLN	GLN	GLN	ALA	ALA	ILE	VAL	SER	SER	GLN	GLY	ASN	ASN	ALA	VAL	GLY	ASP	SER	ALA	PRO	SER	THR	SER	VAL
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----



• Molecule 4: Major inner capsid protein VP3




THR	ALA	SER	PRO	ALA	ASP	THR	ASN	VAL	VAL	PRO	PRO	LYS	ASP	ALA	PRO	THR	THR	ASN	SER	PRO	PRO	SER	SER	THR	THR	SER	PRO	ASN	GLN	GLN	ALA	ALA	ALA	ASP	ASN	GLN	GLN	GLN	ALA	ALA	ILE	VAL	SER	SER	GLN	GLY	ASN	ASN	ALA	VAL	GLY	ASP	SER	ALA	PRO	SER	THR	SER	VAL
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----




• Molecule 5: Clamp protein VP6



• Molecule 5: Clamp protein VP6



• Molecule 5: Clamp protein VP6



• Molecule 5: Clamp protein VP6

Chain a:  99%



- Molecule 5: Clamp protein VP6

Chain b:  100%



- Molecule 5: Clamp protein VP6

Chain d:  100%



- Molecule 5: Clamp protein VP6

Chain e:  100%



- Molecule 5: Clamp protein VP6

Chain g:  99%



- Molecule 5: Clamp protein VP6

Chain h:  99%




- Molecule 5: Clamp protein VP6


Chain n:  100%




- Molecule 6: Outer capsid protein VP1




- Molecule 6: Outer capsid protein VP1




- Molecule 6: Outer capsid protein VP1



- Molecule 6: Outer capsid protein VP1




- Molecule 6: Outer capsid protein VP1



- Molecule 7: RNA (52-MER)







● Molecule 7: RNA (52-MER)



● Molecule 8: RNA (60-MER)




● Molecule 8: RNA (60-MER)




● Molecule 9: RNA (40-MER)




● Molecule 9: RNA (40-MER)




● Molecule 10: RNA (18-MER)




● Molecule 10: RNA (18-MER)



● Molecule 11: RNA (38-MER)



● Molecule 11: RNA (38-MER)



● Molecule 12: RNA (30-MER)



● Molecule 12: RNA (30-MER)





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	119182	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	45	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	81000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.039	Depositor
Minimum map value	-0.018	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.00665	Depositor
Map size (Å)	435.2, 435.2, 435.2	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.36, 1.36, 1.36	Depositor

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/10261	0.50	0/14016
2	B	0.31	0/5734	0.52	0/7849
3	C	0.33	0/8516	0.49	0/11674
3	D	0.33	0/8985	0.49	0/12316
3	E	0.33	0/8755	0.49	0/12001
3	F	0.33	0/8985	0.49	0/12316
3	G	0.33	0/8696	0.49	0/11917
3	H	0.33	0/8922	0.49	0/12229
3	I	0.33	0/8439	0.49	0/11562
3	J	0.33	0/8922	0.49	0/12229
3	K	0.33	0/8696	0.49	0/11917
3	L	0.33	0/8985	0.49	0/12316
4	M	0.28	0/650	0.46	0/898
4	N	0.30	0/212	0.46	0/291
4	k	0.28	0/213	0.47	0/291
4	l	0.27	0/213	0.46	0/291
4	m	0.26	0/213	0.47	0/291
5	V	0.32	0/3234	0.47	0/4444
5	W	0.33	0/3234	0.48	0/4444
5	X	0.32	0/3234	0.47	0/4444
5	a	0.33	0/3234	0.48	0/4444
5	b	0.32	0/3234	0.47	0/4444
5	d	0.33	0/3234	0.48	0/4444
5	e	0.32	0/3234	0.47	0/4444
5	g	0.32	0/3234	0.48	0/4444
5	h	0.32	0/3234	0.47	0/4444
5	n	0.33	0/3234	0.48	0/4444
6	Y	0.31	0/10233	0.51	0/14057
6	Z	0.31	0/10233	0.51	0/14057
6	c	0.31	0/10233	0.51	0/14057
6	f	0.31	0/10233	0.51	0/14057
6	i	0.31	0/10233	0.51	0/14057

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
7	a1	0.16	0/1218	0.73	0/1890
7	b1	0.17	0/1218	0.77	0/1890
8	a2	0.17	0/1406	0.77	0/2182
8	b2	0.16	0/1406	0.73	0/2182
9	a3	0.16	0/936	0.73	0/1452
9	b3	0.17	0/936	0.77	0/1452
10	a4	0.18	0/419	0.76	0/649
10	b4	0.16	0/419	0.72	0/649
11	a5	0.16	0/889	0.73	0/1379
11	b5	0.17	0/889	0.76	0/1379
12	a6	0.16	0/701	0.72	0/1087
12	b6	0.17	0/701	0.76	0/1087
All	All	0.31	0/200040	0.51	0/276407

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1271/1273 (100%)	1212 (95%)	59 (5%)	0	100	100
2	B	716/718 (100%)	688 (96%)	28 (4%)	0	100	100
3	C	1073/1138 (94%)	1026 (96%)	47 (4%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	D	1136/1138 (100%)	1097 (97%)	39 (3%)	0	100	100
3	E	1104/1138 (97%)	1064 (96%)	40 (4%)	0	100	100
3	F	1136/1138 (100%)	1094 (96%)	42 (4%)	0	100	100
3	G	1096/1138 (96%)	1056 (96%)	40 (4%)	0	100	100
3	H	1128/1138 (99%)	1070 (95%)	58 (5%)	0	100	100
3	I	1061/1138 (93%)	1022 (96%)	39 (4%)	0	100	100
3	J	1128/1138 (99%)	1075 (95%)	53 (5%)	0	100	100
3	K	1096/1138 (96%)	1047 (96%)	49 (4%)	0	100	100
3	L	1136/1138 (100%)	1083 (95%)	53 (5%)	0	100	100
4	M	92/94 (98%)	91 (99%)	1 (1%)	0	100	100
4	N	28/94 (30%)	28 (100%)	0	0	100	100
4	k	28/94 (30%)	26 (93%)	2 (7%)	0	100	100
4	l	28/94 (30%)	27 (96%)	1 (4%)	0	100	100
4	m	28/94 (30%)	27 (96%)	1 (4%)	0	100	100
5	V	409/411 (100%)	401 (98%)	8 (2%)	0	100	100
5	W	409/411 (100%)	400 (98%)	9 (2%)	0	100	100
5	X	409/411 (100%)	403 (98%)	6 (2%)	0	100	100
5	a	409/411 (100%)	400 (98%)	9 (2%)	0	100	100
5	b	409/411 (100%)	402 (98%)	7 (2%)	0	100	100
5	d	409/411 (100%)	400 (98%)	9 (2%)	0	100	100
5	e	409/411 (100%)	403 (98%)	6 (2%)	0	100	100
5	g	409/411 (100%)	397 (97%)	12 (3%)	0	100	100
5	h	409/411 (100%)	401 (98%)	8 (2%)	0	100	100
5	n	409/411 (100%)	397 (97%)	12 (3%)	0	100	100
6	Y	1295/1297 (100%)	1254 (97%)	41 (3%)	0	100	100
6	Z	1295/1297 (100%)	1255 (97%)	40 (3%)	0	100	100
6	c	1295/1297 (100%)	1255 (97%)	40 (3%)	0	100	100
6	f	1295/1297 (100%)	1253 (97%)	42 (3%)	0	100	100
6	i	1295/1297 (100%)	1255 (97%)	39 (3%)	1 (0%)	48	76
All	All	23850/24436 (98%)	23009 (96%)	840 (4%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
6	i	1291	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1092/1092 (100%)	1085 (99%)	7 (1%)	84	90
2	B	619/619 (100%)	615 (99%)	4 (1%)	84	90
3	C	917/972 (94%)	913 (100%)	4 (0%)	89	93
3	D	972/972 (100%)	967 (100%)	5 (0%)	86	91
3	E	946/972 (97%)	938 (99%)	8 (1%)	79	87
3	F	972/972 (100%)	961 (99%)	11 (1%)	70	82
3	G	938/972 (96%)	933 (100%)	5 (0%)	86	91
3	H	964/972 (99%)	954 (99%)	10 (1%)	73	84
3	I	910/972 (94%)	905 (100%)	5 (0%)	86	91
3	J	964/972 (99%)	958 (99%)	6 (1%)	84	90
3	K	938/972 (96%)	930 (99%)	8 (1%)	75	85
3	L	972/972 (100%)	963 (99%)	9 (1%)	75	85
4	M	73/73 (100%)	71 (97%)	2 (3%)	40	65
4	N	24/73 (33%)	24 (100%)	0	100	100
4	k	24/73 (33%)	24 (100%)	0	100	100
4	l	24/73 (33%)	24 (100%)	0	100	100
4	m	24/73 (33%)	24 (100%)	0	100	100
5	V	325/325 (100%)	322 (99%)	3 (1%)	75	85
5	W	325/325 (100%)	323 (99%)	2 (1%)	84	90
5	X	325/325 (100%)	322 (99%)	3 (1%)	75	85
5	a	325/325 (100%)	322 (99%)	3 (1%)	75	85
5	b	325/325 (100%)	324 (100%)	1 (0%)	91	94
5	d	325/325 (100%)	323 (99%)	2 (1%)	84	90

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	e	325/325 (100%)	323 (99%)	2 (1%)	84	90
5	g	325/325 (100%)	322 (99%)	3 (1%)	75	85
5	h	325/325 (100%)	322 (99%)	3 (1%)	75	85
5	n	325/325 (100%)	323 (99%)	2 (1%)	84	90
6	Y	1089/1089 (100%)	1077 (99%)	12 (1%)	70	82
6	Z	1089/1089 (100%)	1081 (99%)	8 (1%)	81	88
6	c	1089/1089 (100%)	1082 (99%)	7 (1%)	84	90
6	f	1089/1089 (100%)	1081 (99%)	8 (1%)	81	88
6	i	1089/1089 (100%)	1079 (99%)	10 (1%)	75	85
All	All	20068/20491 (98%)	19915 (99%)	153 (1%)	77	87

5 of 153 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	b	255	ARG
6	i	243	GLN
6	c	929	MET
6	f	711	ASP
6	i	1296	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
6	Y	630	GLN
5	b	158	HIS
5	a	338	HIS
6	f	630	GLN
3	J	353	GLN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
10	a4	17/18 (94%)	0	0
10	b4	17/18 (94%)	0	0
11	a5	37/38 (97%)	0	0
11	b5	37/38 (97%)	0	0
12	a6	29/30 (96%)	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
12	b6	29/30 (96%)	0	0
7	a1	51/52 (98%)	0	0
7	b1	51/52 (98%)	0	0
8	a2	59/60 (98%)	0	0
8	b2	59/60 (98%)	0	0
9	a3	39/40 (97%)	0	0
9	b3	39/40 (97%)	0	0
All	All	464/476 (97%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.




## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-29243. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


### 6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 178




Y Index: 125




Z Index: 168


### 6.3.2 Raw map



X Index: 178



Y Index: 125




Z Index: 168


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.00665. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 2845  $\text{nm}^3$ ; this corresponds to an approximate mass of 2570 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.303 Å<sup>-1</sup>



## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.303 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.30	-	-
Author-provided FSC curve	3.26	3.71	3.30
Unmasked-calculated*	3.48	4.14	3.58

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-29243 and PDB model 8FJK. Per-residue inclusion information can be found in section 3 on page 9.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.00665 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.00665).







































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 95% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary





















The table lists the average atom inclusion at the recommended contour level (0.00665) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8270	 0.4670
A	 0.8100	 0.4790
B	 0.7420	 0.4450
C	 0.8750	 0.5130
D	 0.8660	 0.5050
E	 0.8730	 0.5110
F	 0.8680	 0.5070
G	 0.8750	 0.5140
H	 0.8710	 0.5060
I	 0.8750	 0.5150
J	 0.8680	 0.5030
K	 0.8770	 0.5110
L	 0.8750	 0.5060
M	 0.7880	 0.4980
N	 0.7600	 0.4950
V	 0.9110	 0.4990
W	 0.9250	 0.5110
X	 0.9140	 0.4970
Y	 0.8700	 0.4560
Z	 0.8690	 0.4590
a	 0.9230	 0.5120
a1	 0.0540	 0.0240
a2	 0.0350	 0.0340
a3	 0.0000	 0.0060
a4	 0.0030	 0.0280
a5	 0.0110	 0.0020
a6	 0.0000	 0.0230
b	 0.9110	 0.4940
b1	 0.0140	 0.0320
b2	 0.0190	 0.0290
b3	 0.0010	 -0.0000
b4	 0.0000	 0.0310
b5	 0.0200	 0.0340
b6	 0.0000	 0.0220
c	 0.8680	 0.4580



*Continued on next page...*

*Continued from previous page...*

Chain	Atom inclusion	Q-score
d	 0.9280	 0.5130
e	 0.9090	 0.5020
f	 0.8740	 0.4590
g	 0.9270	 0.5160
h	 0.9190	 0.5020
i	 0.8720	 0.4560
k	 0.8040	 0.5010
l	 0.7890	 0.4830
m	 0.7660	 0.4600
n	 0.9250	 0.5200