

Full wwPDB X-ray Structure Validation Report (i)

Oct 25, 2022 – 07:00 PM EDT

PDB ID : 7FK1

Title : PanDDA analysis group deposition – Aar2/RNaseH in complex with fragment

P04A06 from the F2X-Universal Library

Authors: Barthel, T.; Wollenhaupt, J.; Lima, G.M.A.; Wahl, M.C.; Weiss, M.S.

Deposited on : 2022-08-26

Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.31.2

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	258	78%	10% • • 8%
2	В	308	9%	7% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



\mathbf{Mol}	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	W3H	A	2101[A]	-	-	-	X
3	W3H	A	2101[B]	-	-	-	X



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mo	l Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	237	Total 1997	C 1281	N 332	O 372	S 12	0	11	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

• Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	В	300	Total 2580	C 1654	N 421	O 485	S 20	0	9	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
В	-3	GLY	=	expression tag	UNP P32357
В	-2	ALA	-	expression tag	UNP P32357
В	-1	MET	-	expression tag	UNP P32357
В	0	ALA	-	expression tag	UNP P32357
В	166	SER	LEU	$\operatorname{conflict}$	UNP P32357
В	167	SER	LYS	$\operatorname{conflict}$	UNP P32357
В	?	-	LEU	deletion	UNP P32357
В	?	-	GLN	deletion	UNP P32357
В	?	-	LYS	deletion	UNP P32357
В	?	-	ALA	deletion	UNP P32357
В	?	-	GLY	deletion	UNP P32357
В	?	-	SER	deletion	UNP P32357
В	?	-	LYS	deletion	UNP P32357



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	?	-	MET	deletion	UNP P32357
В	?	-	GLU	deletion	UNP P32357
В	?	-	ALA	deletion	UNP P32357
В	?	-	LYS	deletion	UNP P32357
В	?	-	ASN	deletion	UNP P32357
В	?	-	GLU	deletion	UNP P32357
В	170	SER	ASP	conflict	UNP P32357

• Molecule 3 is (3R)-3-(4-chlorobenzene-1-sulfonyl) butanoic acid (three-letter code: W3H) (formula: $C_{10}H_{11}ClO_4S$).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
3	Λ	1	Total	С	Cl	О	S	0	1
3	3 A	1	32	20	2	8	2	U	1
9	D	1	Total	С	Cl	О	S	0	0
3	D	1	16	10	1	4	1	U	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	40	Total O 40 40	0	0
4	В	29	Total O 29 29	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pre-mRNA-splicing factor 8





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	89.61Å 82.69Å 94.59Å	Donositor
a, b, c, α , β , γ	90.00° 108.02° 90.00°	Depositor
Resolution (Å)	45.26 - 1.76	Depositor
Resolution (A)	45.26 - 1.76	EDS
% Data completeness	99.0 (45.26-1.76)	Depositor
(in resolution range)	99.0 (45.26-1.76)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.95 (at 1.76Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
D D.	0.280 , 0.315	Depositor
R, R_{free}	0.262 , 0.296	DCC
R_{free} test set	2101 reflections (3.27%)	wwPDB-VP
Wilson B-factor (Å ²)	42.1	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 48.5	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4694	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.53% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: W3H

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		nd lengths	Bond angles		
MIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.74	1/2044 (0.0%)	0.94	$12/2770 \ (0.4\%)$	
2	В	0.62	0/2651	0.72	$2/3581 \ (0.1\%)$	
All	All	0.67	$1/4695 \ (0.0\%)$	0.82	$14/6351 \ (0.2\%)$	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	В	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
1	A	1884	PRO	N-CA	11.99	1.67	1.47

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	1884	PRO	CB-CA-C	8.52	133.31	112.00
1	A	1884	PRO	N-CA-CB	-8.01	93.69	103.30
1	A	1840	TYR	CB-CG-CD2	7.91	125.75	121.00
1	A	1842	GLU	N-CA-CB	6.24	121.84	110.60
1	A	1842	GLU	CB-CA-C	-6.05	98.31	110.40
1	A	1846	ASN	CB-CA-C	-5.99	98.42	110.40
1	A	1849	LYS	CB-CA-C	-5.88	98.64	110.40
2	В	230[A]	ASN	CA-C-O	5.72	132.12	120.10



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
2	В	230[B]	ASN	CA-C-O	5.72	132.12	120.10
1	A	1992	TYR	CB-CG-CD1	5.68	124.41	121.00
1	A	1992	TYR	CB-CG-CD2	-5.49	117.71	121.00
1	A	1838	SER	C-N-CA	5.27	134.89	121.70
1	A	1836	ASN	C-N-CA	5.04	134.30	121.70
1	A	1841	ALA	C-N-CA	5.01	134.23	121.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1959	THR	Mainchain
2	В	54[A]	MET	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1997	0	2028	25	4
2	В	2580	0	2450	12	4
3	A	32	0	0	1	0
3	В	16	0	0	1	0
4	A	40	0	0	2	0
4	В	29	0	0	2	0
All	All	4694	0	4478	38	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:A:1884:PRO:N	1:A:1884:PRO:CA	1.67	1.36
1:A:1846:ASN:ND2	1:A:1846:ASN:O	2.06	0.89
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ2	1.35	0.73




 $Continued\ from\ previous\ page...$

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	$-$ overlap (\AA)
1:A:1835:MET:HA	1:A:1835:MET:HE3	1.75	0.69
1:A:1834:ALA:HB2	1:A:1958:PRO:HG2	1.76	0.68
2:B:53:SER:O	4:B:501:HOH:O	2.13	0.67
1:A:1833:GLY:O	1:A:1957:ARG:HB3	1.95	0.66
1:A:1840:TYR:CE2	3:A:2101[A]:W3H:CL	2.92	0.60
1:A:1884:PRO:N	1:A:1884:PRO:C	2.53	0.59
1:A:1842:GLU:O	1:A:1842:GLU:HG3	2.05	0.56
2:B:12:GLU:HG3	2:B:25:LYS:HA	1.86	0.55
1:A:1938:LYS:NZ	4:A:2201:HOH:O	2.34	0.54
3:B:401:W3H:C	3:B:401:W3H:C5	2.85	0.54
1:A:2058:LEU:C	1:A:2058:LEU:HD23	2.29	0.53
1:A:1848:ILE:H	1:A:1931[A]:LYS:NZ	2.05	0.53
2:B:275:LEU:CD2	2:B:283:LEU:HD13	2.40	0.52
1:A:1847:ASP:HB3	1:A:1931[A]:LYS:HB2	1.90	0.52
2:B:16:GLY:HA3	2:B:45:HIS:CE1	2.45	0.52
2:B:287:ARG:O	2:B:291:ILE:HD13	2.09	0.52
1:A:2069:VAL:C	4:A:2224:HOH:O	2.48	0.51
1:A:1834:ALA:CB	1:A:1958:PRO:HG2	2.39	0.51
2:B:277:GLU:CD	2:B:277:GLU:H	2.14	0.51
1:A:2027:LEU:O	1:A:2028:SER:C	2.49	0.50
1:A:1835:MET:SD	1:A:1957:ARG:NH2	2.84	0.50
1:A:1847:ASP:HB3	1:A:1931[B]:LYS:HG3	1.94	0.48
2:B:275:LEU:HD22	2:B:283:LEU:HD13	1.95	0.48
1:A:1998:ARG:NH1	1:A:2045:ASP:OD2	2.41	0.47
2:B:110:GLU:HG2	2:B:110:GLU:O	2.15	0.47
1:A:1848:ILE:N	1:A:1931[A]:LYS:HZ2	2.09	0.45
2:B:258:LYS:HD2	2:B:258:LYS:H	1.82	0.45
1:A:1835:MET:SD	1:A:1842:GLU:HG2	2.57	0.45
1:A:1933:ILE:HG21	1:A:1944:LEU:HD21	2.00	0.44
2:B:28:GLN:CG	2:B:29:PRO:HD2	2.49	0.43
2:B:1:MET:N	4:B:504:HOH:O	2.52	0.42
2:B:1:MET:HB3	2:B:35:ASP:HA	2.02	0.42
1:A:2066:LYS:HD3	1:A:2067:TYR:CZ	2.55	0.42
1:A:2005:PHE:O	1:A:2009:THR:HG23	2.21	0.41
1:A:2066:LYS:HD3	1:A:2067:TYR:CE1	2.56	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:1841:ALA:O	2:B:115:TYR:CE1[4_555]	1.53	0.67





Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:1846:ASN:CB	2:B:112:ASP:OD2[4_555]	1.75	0.45
1:A:2069:VAL:O	2:B:52:SER:OG[2_655]	1.98	0.22
1:A:1846:ASN:CG	2:B:112:ASP:OD2[4_555]	2.07	0.13

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	247/258 (96%)	236 (96%)	9 (4%)	2 (1%)	19 6
2	В	306/308 (99%)	293 (96%)	13 (4%)	0	100 100
All	All	553/566 (98%)	529 (96%)	22 (4%)	2 (0%)	34 17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1846	ASN
1	A	2028	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	227/233 (97%)	212 (93%)	15 (7%)	16 3
2	В	287/284 (101%)	283 (99%)	4 (1%)	67 52
All	All	514/517~(99%)	495 (96%)	19 (4%)	38 12



All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1835	MET
1	A	1837	SER
1	A	1839	ASN
1	A	1840	TYR
1	A	1846	ASN
1	A	1849	LYS
1	A	1884	PRO
1	A	1885	LYS
1	A	1931[A]	LYS
1	A	1931[B]	LYS
1	A	1962	ARG
1	A	1979[A]	MET
1	A	1979[B]	MET
1	A	1979[C]	MET
1	A	2058	LEU
2	В	77	LEU
2	В	258	LYS
2	В	295	SER
2	В	317	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	В	300	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Chain Res Link Bond lengths		Bond angles					
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	W3H	A	2101[A]	-	14,16,16	0.69	0	21,23,23	1.91	4 (19%)
3	W3H	A	2101[B]	-	14,16,16	0.51	0	21,23,23	0.60	0
3	W3H	В	401	-	14,16,16	6.96	4 (28%)	21,23,23	2.65	11 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	W3H	A	2101[A]	-	-	4/16/16/16	0/1/1/1
3	W3H	A	2101[B]	-	-	1/16/16/16	0/1/1/1
3	W3H	В	401	-	-	0/16/16/16	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
3	В	401	W3H	O2-S	-20.17	1.25	1.44
3	В	401	W3H	O3-S	-11.99	1.33	1.44
3	В	401	W3H	C1-S	-10.50	1.58	1.79
3	В	401	W3H	O-C3	-3.05	1.20	1.30

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
3	A	2101[A]	W3H	C-C1-S	-6.33	91.97	109.90
3	В	401	W3H	O2-S-O3	-5.99	113.22	118.71
3	В	401	W3H	C5-C4-S	5.45	124.14	119.44
3	A	2101[A]	W3H	C4-S-C1	4.22	113.19	105.14
3	В	401	W3H	C4-S-C1	-3.67	98.15	105.14



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
3	В	401	W3H	C5-C4-C9	-3.62	115.39	120.44
3	В	401	W3H	O3-S-C1	3.32	113.13	108.16
3	В	401	W3H	C6-C5-C4	3.19	122.75	119.45
3	В	401	W3H	C8-C9-C4	2.74	122.28	119.45
3	В	401	W3H	O2-S-C1	2.56	111.99	108.16
3	В	401	W3H	O2-S-C4	2.46	110.57	108.18
3	В	401	W3H	O1-C3-C2	-2.19	115.79	122.80
3	A	2101[A]	W3H	O2-S-C1	-2.14	104.94	108.16
3	В	401	W3H	O-C3-O1	2.12	128.58	123.30
3	A	2101[A]	W3H	C1-C2-C3	-2.05	108.03	113.10

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2101[A]	W3H	C5-C4-S-O3
3	A	2101[A]	W3H	C5-C4-S-C1
3	A	2101[A]	W3H	C9-C4-S-O3
3	A	2101[B]	W3H	C-C1-C2-C3
3	A	2101[A]	W3H	C9-C4-S-C1


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2101[A]	W3H	1	0
3	В	401	W3H	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	$237/258 \ (91\%)$	1.19	26 (10%) 5 7	33, 48, 89, 193	0
2	В	300/308 (97%)	0.86	29 (9%) 7 10	33, 50, 97, 136	0
All	All	537/566 (94%)	1.00	55 (10%) 6 9	33, 49, 96, 193	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1841	ALA	27.1
1	A	1839	ASN	14.0
2	В	1	MET	12.2
1	A	1843	LEU	11.2
1	A	1838	SER	10.8
1	A	1835	MET	10.6
1	A	1836	ASN	9.5
1	A	1837	SER	9.3
2	В	170	SER	8.9
2	В	172	PRO	8.7
1	A	1833	GLY	8.3
1	A	1844	PHE	8.3
1	A	1840	TYR	8.2
1	A	1842	GLU	7.5
1	A	1845	ASN	7.3
2	В	173	ALA	6.7
2	В	54[A]	MET	6.6
1	A	1846	ASN	5.6
1	A	2027	LEU	5.2
2	В	108	ILE	5.2
1	A	1834	ALA	5.0
2	В	171	ASP	4.8
2	В	109	ASP	4.5
2	В	174	HIS	4.3



Continued from previous page...

Mol	Chain	Res	Type	RSRZ	
2	В	150	ASN	4.1	
2	В	110	GLU	4.1	
2	В	317	LEU	3.6	
1	A	2046	GLU	3.5	
2	В	107	LYS	3.3	
2	В	293	LEU	3.3	
1	A	1953	ASN	3.1	
1	A	2032	ILE	2.8	
1	A	2028	SER	2.8	
2	В	286	GLU	2.8	
1	A	1969	MET	2.7	
2	В	39	GLY	2.6	
2	В	38	ILE	2.6	
1	A	2033	THR	2.6	
2	В	84	ARG	2.6	
1	A	2069	VAL	2.6	
2	В	316	LEU	2.5	
2	В	259	HIS	2.5	
2	В	62	CYS	2.4	
2	В	275	LEU	2.3	
2	В	181	ILE	2.3	
1	A	1866	PHE	2.3	
1	A	1957	ARG	2.3	
2	В	92	ILE	2.3	
2	В	148	GLN	2.2	
1	A	1861	THR	2.2	
2	В	43	VAL	2.2	
2	В	247	ILE	2.1	
1	A	1863	HIS	2.1	
2	В	94	HIS	2.1	
2	В	101	MET	2.1	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	W3H	A	2101[A]	16/16	0.59	0.47	20,20,20,20	16
3	W3H	A	2101[B]	16/16	0.59	0.47	30,30,30,30	16
3	W3H	В	401	16/16	0.81	0.44	20,20,20,20	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around W3H A 2101 (B):

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around W3H B 401:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

