

# Full wwPDB X-ray Structure Validation Report (i)

#### Sep 28, 2024 – 11:42 PM EDT

PDB ID : 1FOF

Title : CRYSTAL STRUCTURE OF THE CLASS D BETA-LACTAMASE OXA-10 Authors : Paetzel, M.; Danel, F.; de Castro, L.; Mosimann, S.C.; Page, M.G.P.; Stry-

nadka, N.C.J.

Deposited on : 2000-08-28

Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp

with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain	
1	A	246	87%	13%
1	В	246	80%	19% •



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4355 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called BETA LACTAMASE OXA-10.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	246	Total 1940	C 1240	11	O 368	S 6	0	0	0
1	В	246	Total 1936	C 1238	- 1	O 366	S 6	0	0	0

• Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Co 1 1	0	0
2	В	1	Total Co 1 1	0	0

• Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0

### $\bullet$ Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	270	Total O 270 270	0	0
4	В	197	Total O 197 197	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: BETA LACTAMASE OXA-10





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	48.40Å 96.20Å 125.70Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.79 - 2.00	Depositor
% Data completeness	93.3 (19.79-2.00)	Depositor
(in resolution range)	30.9 (13.13 2.00)	Depositor
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
$R, R_{free}$	0.177 , 0.208	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4355	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP



## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles	
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.34	0/1981	0.58	0/2675
1	В	0.31	0/1977	0.55	0/2670
All	All	0.32	0/3958	0.56	0/5345

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1940	0	1933	17	0
1	В	1936	0	1929	34	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
3	A	5	0	0	0	0
3	В	5	0	0	0	0
4	A	270	0	0	1	0
4	В	197	0	0	0	0
All	All	4355	0	3862	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.



All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A. 1	A.1 0	Interatomic	Clash
Atom-1	Atom-2	${ m distance} \ ({ m \AA})$	overlap (Å)
1:A:24:GLU:HG3	1:A:56:LEU:HG	1.54	0.90
1:B:138:LYS:HE3	1:B:179:SER:CB	2.05	0.87
1:B:70:LYS:HE2	1:B:115:SER:HA	1.60	0.80
1:B:176:ASN:HB3	1:B:182:LYS:HD3	1.68	0.75
4:A:727:HOH:O	1:B:230:THR:HG21	1.87	0.73
1:B:138:LYS:HE3	1:B:179:SER:HB3	1.69	0.72
1:B:247:LEU:O	1:B:250:ARG:HG2	1.96	0.66
1:A:255:THR:O	1:A:259:GLU:HG3	1.98	0.64
1:B:117:VAL:N	1:B:118:PRO:HD2	2.15	0.61
1:B:255:THR:O	1:B:259:GLU:HG3	2.01	0.61
1:A:24:GLU:CG	1:A:56:LEU:HG	2.29	0.59
1:B:25:ASN:C	1:B:25:ASN:HD22	2.06	0.58
1:A:44:CYS:SG	1:A:167:VAL:HG11	2.44	0.57
1:A:110:GLY:O	1:A:114:VAL:HG22	2.03	0.57
1:B:153:PHE:HA	1:B:157:GLY:HA3	1.86	0.57
1:B:243:ASN:ND2	1:B:245:SER:OG	2.38	0.57
1:B:91:LYS:HA	1:B:105:ASP:OD1	2.05	0.56
1:B:247:LEU:O	1:B:250:ARG:CG	2.57	0.52
1:B:164:VAL:O	1:B:168:GLU:HG3	2.10	0.51
1:A:109:ARG:O	1:A:113:GLN:HG2	2.09	0.51
1:B:102:TRP:CE2	1:B:116:ALA:HA	2.45	0.51
1:A:98:ALA:HB3	1:A:102:TRP:CZ3	2.45	0.50
1:B:93:ASP:OD2	1:B:95:LYS:HD2	2.11	0.49
1:A:61:LYS:HD3	1:A:63:TYR:OH	2.13	0.49
1:B:220:ALA:O	1:B:238:ASN:HA	2.13	0.49
1:A:164:VAL:O	1:A:168:GLU:HG3	2.13	0.49
1:B:230:THR:CG2	1:B:230:THR:O	2.62	0.48
1:B:98:ALA:HB3	1:B:102:TRP:CZ3	2.50	0.47
1:B:138:LYS:HE3	1:B:179:SER:HB2	1.94	0.47
1:A:121:GLN:O	1:A:125:ARG:HG3	2.16	0.46
1:B:92:TRP:CZ2	1:B:103:GLU:HB3	2.52	0.45
1:B:230:THR:O	1:B:230:THR:HG22	2.17	0.45
1:B:117:VAL:O	1:B:121:GLN:HG3	2.17	0.45
1:B:24:GLU:HB2	1:B:56:LEU:HG	1.98	0.44
1:A:247:LEU:O	1:A:250:ARG:HG2	2.18	0.44
1:A:220:ALA:O	1:A:238:ASN:HA	2.19	0.43
1:B:129:GLU:HG3	1:B:146:ILE:HD11	2.00	0.43
1:A:64:LEU:HD21	1:A:155:LEU:O	2.19	0.43
1:B:113:GLN:C	1:B:115:SER:H	2.21	0.43
1:A:101:GLN:O	1:A:104:ARG:NH1	2.52	0.42

Continued on next page...



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:PHE:CE2	1:A:250:ARG:HA	2.54	0.42
1:B:151:ASP:OD2	1:B:152:LYS:HD2	2.18	0.42
1:B:93:ASP:C	1:B:95:LYS:H	2.21	0.42
1:A:171:GLU:HA	1:A:234:PHE:CE2	2.54	0.42
1:B:25:ASN:C	1:B:25:ASN:ND2	2.71	0.42
1:B:103:GLU:O	1:B:104:ARG:HB3	2.20	0.42
1:B:164:VAL:O	1:B:167:VAL:HG22	2.20	0.42
1:B:26:THR:HA	1:B:29:ASN:ND2	2.34	0.42
1:B:113:GLN:C	1:B:115:SER:N	2.73	0.42
1:B:171:GLU:HA	1:B:234:PHE:CE2	2.55	0.42
1:A:101:GLN:HG3	1:A:102:TRP:CD1	2.57	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers   Percentil		$\mathbf{ntiles}$
1	A	$244/246 \ (99\%)$	236 (97%)	8 (3%)	0	100	100
1	В	$244/246 \ (99\%)$	234 (96%)	10 (4%)	0	100	100
All	All	488/492 (99%)	470 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	210/210 (100%)	208 (99%)	2 (1%)	73 78		
1	В	209/210 (100%)	204 (98%)	5 (2%)	44 47		
All	All	419/420 (100%)	412 (98%)	7 (2%)	56 61		

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	ASN
1	A	165	ASN
1	В	25	ASN
1	В	73	ASN
1	В	165	ASN
1	В	166	GLN
1	В	250	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	A	73	ASN
1	A	87	HIS
1	A	101	GLN
1	A	158	GLN
1	В	25	ASN
1	В	38	ASN
1	В	73	ASN
1	В	144	GLN
1	В	166	GLN
1	В	243	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol T	Т	Chain	Res	Link	Bond lengths			Bond angles		
	Type				Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	SO4	В	489	-	4,4,4	0.31	0	6,6,6	0.10	0
3	SO4	A	488	-	4,4,4	0.34	0	6,6,6	0.12	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

