

# Full wwPDB X-ray Structure Validation Report (i)

Nov 27, 2024 – 12:20 pm GMT

PDB ID : 9FRV

Title: Arginase 2 in complex with inhibitor

Authors : Petersen, j. Deposited on : 2024-06-19

Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.40

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	164625	3436 (2.08-2.04)
Clashscore	180529	3661 (2.08-2.04)
Ramachandran outliers	177936	3649 (2.08-2.04)
Sidechain outliers	177891	3649 (2.08-2.04)
RSRZ outliers	164620	3436 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	336	7% 84%	10% • 5%
1	В	336	27% 87%	9% •
1	С	336	7%	10% • 5%



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7423 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Arginase-2, mitochondrial.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	318	Total	С	N	О	S	0	0	0
1	A	310	2430	1536	422	463	9	0	U	
1	D	324	Total	С	N	О	S	0	0	0
1	Ъ	324	2477	1567	428	473	9	0	0	
1	C	318	Total	С	N	О	S	0	0	0
1		310	2430	1536	422	463	9	U	U	

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6	MET	-	initiating methionine	UNP P78540
A	7	HIS	-	expression tag	UNP P78540
A	8	HIS	-	expression tag	UNP P78540
A	9	HIS	-	expression tag	UNP P78540
A	10	HIS	-	expression tag	UNP P78540
A	11	HIS	-	expression tag	UNP P78540
A	12	HIS	-	expression tag	UNP P78540
A	13	GLY	-	expression tag	UNP P78540
A	14	GLY	-	expression tag	UNP P78540
A	15	GLY	-	expression tag	UNP P78540
A	16	GLU	-	expression tag	UNP P78540
A	17	ASN	-	expression tag	UNP P78540
A	18	LEU	-	expression tag	UNP P78540
A	19	TYR	-	expression tag	UNP P78540
A	20	PHE	-	expression tag	UNP P78540
A	21	GLN	-	expression tag	UNP P78540
В	6	MET	-	initiating methionine	UNP P78540
В	7	HIS	-	expression tag	UNP P78540
В	8	HIS	-	expression tag	UNP P78540
В	9	HIS	-	expression tag	UNP P78540
В	10	HIS	-	expression tag	UNP P78540
В	11	HIS	-	expression tag	UNP P78540
В	12	HIS	_	expression tag	UNP P78540



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
В	13	GLY	-	expression tag	UNP P78540
В	14	GLY	-	expression tag	UNP P78540
В	15	GLY	-	expression tag	UNP P78540
В	16	GLU	-	expression tag	UNP P78540
В	17	ASN	-	expression tag	UNP P78540
В	18	LEU	-	expression tag	UNP P78540
В	19	TYR	-	expression tag	UNP P78540
В	20	PHE	-	expression tag	UNP P78540
В	21	GLN	-	expression tag	UNP P78540
С	6	MET	-	initiating methionine	UNP P78540
С	7	HIS	-	expression tag	UNP P78540
С	8	HIS	-	expression tag	UNP P78540
С	9	HIS	-	expression tag	UNP P78540
С	10	HIS	-	expression tag	UNP P78540
С	11	HIS	-	expression tag	UNP P78540
С	12	HIS	-	expression tag	UNP P78540
С	13	GLY	-	expression tag	UNP P78540
С	14	GLY	-	expression tag	UNP P78540
С	15	GLY	-	expression tag	UNP P78540
С	16	GLU	-	expression tag	UNP P78540
С	17	ASN	-	expression tag	UNP P78540
С	18	LEU	-	expression tag	UNP P78540
С	19	TYR	-	expression tag	UNP P78540
С	20	PHE	-	expression tag	UNP P78540
С	21	GLN	-	expression tag	UNP P78540

• Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mn 2 2	0	0
2	В	2	Total Mn 2 2	0	0
2	С	2	Total Mn 2 2	0	0

• Molecule 3 is [(4 {S},5 {S})-4-(aminomethyl)-5-azanyl-6-oxidanyl-6-oxidanylidene-hexyl ]-\$l^{3}-oxidanyl-bis(oxidanyl)boron (three-letter code: A1IFX) (formula:  $C_7H_{18}BN_2O_5$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	С	1	Total	В	С	N	О	0	0
3		1	15	1	7	2	5	0	

#### • Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	25	Total O 25 25	0	0
4	В	13	Total O 13 13	0	0
4	С	27	Total O 27 27	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Arginase-2, mitochondrial











## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	46.75Å 134.59Å 145.08Å	Donositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.50 - 2.06	Depositor
Resolution (A)	44.50 - 2.06	EDS
% Data completeness	95.1 (44.50-2.06)	Depositor
(in resolution range)	95.0 (44.50-2.06)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.96 (at 2.07Å)	Xtriage
Refinement program	BUSTER 2.11.7 PACIOREK	Depositor
D D.	0.279 , 0.315	Depositor
$R, R_{free}$	0.294 , $0.337$	DCC
$R_{free}$ test set	2097 reflections (3.83%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.7	Xtriage
Anisotropy	0.443	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.31, 32.0	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.53, < L^2>=0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7423	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, A1IFX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	$\mathbf{angles}$
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.42	0/2484	0.64	0/3382
1	В	0.43	0/2531	0.61	0/3444
1	С	0.44	0/2484	0.66	0/3382
All	All	0.43	0/7499	0.64	0/10208

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2430	0	2410	19	0
1	В	2477	0	2451	10	0
1	С	2430	0	2410	16	0
2	A	2	0	0	0	0
2	В	2	0	0	0	0
2	С	2	0	0	0	0
3	С	15	0	0	0	0
4	A	25	0	0	0	0
4	В	13	0	0	0	0
4	С	27	0	0	0	0
All	All	7423	0	7271	45	0



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
4 4 400 CLNIN	4 4 400 DDO HDO	distance (Å)	overlap (Å)
1:A:162:GLN:N	1:A:163:PRO:HD2	2.15	0.61
1:B:57:ARG:HH21	1:B:325:GLN:HE22	1.47	0.60
1:A:151:PRO:HD2	1:A:176:PRO:HD2	1.84	0.59
1:A:162:GLN:H	1:A:163:PRO:HD2	1.66	0.59
1:A:262:ALA:HB2	1:A:302:ALA:HB2	1.88	0.55
1:C:262:ALA:HB1	1:C:298:ASN:O	2.06	0.55
1:A:252:ILE:HD13	1:A:294:LEU:HD11	1.89	0.54
1:C:74:THR:HB	1:C:94:LEU:HD21	1.90	0.54
1:A:255:PHE:CE1	1:A:276:GLY:HA3	2.43	0.54
1:C:147:ASP:HB3	1:C:163:PRO:HD2	1.91	0.52
1:A:57:ARG:HD2	1:A:325:GLN:HE22	1.76	0.50
1:B:145:HIS:CE1	1:B:265:THR:HB	2.46	0.50
1:C:151:PRO:HD2	1:C:176:PRO:HD2	1.93	0.50
1:C:81:LEU:H	1:C:81:LEU:HD23	1.77	0.49
1:B:130:HIS:CE1	1:B:290:SER:HB3	2.48	0.49
1:B:138:CYS:HB2	1:B:238:LEU:HD22	1.95	0.48
1:C:57:ARG:HD2	1:C:325:GLN:HE22	1.78	0.48
1:C:124:ILE:HA	1:C:167:LEU:HD11	1.95	0.48
1:C:243:GLN:HB3	1:C:288:LEU:HD21	1.97	0.47
1:A:248:LEU:HD22	1:A:283:ILE:HG21	1.96	0.46
1:A:57:ARG:HH21	1:A:325:GLN:HE22	1.62	0.46
1:A:169:ARG:HG2	1:A:186:PRO:HB2	1.98	0.45
1:C:202:ASP:HB3	1:C:204:PRO:HD2	2.00	0.43
1:A:33:PHE:CZ	1:A:35:GLN:HB2	2.53	0.43
1:B:195:TYR:O	1:B:196:ILE:HG13	2.19	0.43
1:C:57:ARG:HH21	1:C:325:GLN:HE22	1.65	0.43
1:C:162:GLN:N	1:C:163:PRO:CD	2.81	0.43
1:C:138:CYS:HB2	1:C:238:LEU:HD22	2.01	0.43
1:A:284:HIS:HB2	1:A:322:SER:O	2.19	0.43
1:A:255:PHE:CZ	1:A:276:GLY:HA3	2.54	0.42
1:B:195:TYR:HB3	1:B:198:LEU:HD11	2.01	0.42
1:C:158:ASN:HB3	1:C:160:HIS:CD2	2.54	0.42
1:B:143:ASP:HB3	1:B:145:HIS:O	2.19	0.42
1:C:314:LEU:O	1:C:318:VAL:HG23	2.19	0.42
1:A:162:GLN:H	1:A:163:PRO:CD	2.29	0.42
1:A:138:CYS:HB2	1:A:238:LEU:HD22	2.02	0.42
1:A:262:ALA:HB1	1:A:298:ASN:O	2.19	0.42
		0	



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap (Å)} \end{array}$
1:C:262:ALA:HB2	1:C:302:ALA:HB2	2.02	0.41
1:A:326:THR:HG23	1:A:329:GLY:H	1.85	0.41
1:B:169:ARG:HG2	1:B:186:PRO:HB2	2.03	0.41
1:B:198:LEU:HD13	1:B:216:TYR:HD2	1.86	0.41
1:B:314:LEU:O	1:B:318:VAL:HG23	2.20	0.41
1:C:228:GLN:HA	1:C:278:TYR:CE2	2.55	0.41
1:A:162:GLN:N	1:A:163:PRO:CD	2.82	0.40
1:A:199:ARG:HD3	1:A:219:MET:HG3	2.03	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	316/336~(94%)	302 (96%)	13 (4%)	1 (0%)	37	31
1	В	321/336 (96%)	300 (94%)	18 (6%)	3 (1%)	14	7
1	С	316/336 (94%)	299 (95%)	16 (5%)	1 (0%)	37	31
All	All	953/1008 (94%)	901 (94%)	47 (5%)	5 (0%)	25	17

#### All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	GLN
1	В	86	ILE
1	В	162	GLN
1	С	162	GLN
1	В	239	ILE



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	$268/283 \; (95\%)$	258 (96%)	10 (4%)	29 23
1	В	273/283 (96%)	261 (96%)	12 (4%)	24 18
1	С	268/283 (95%)	257 (96%)	11 (4%)	26 20
All	All	809/849 (95%)	776 (96%)	33 (4%)	26 20

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	ASP
1	A	115	THR
1	A	120	HIS
1	A	137	LEU
1	A	160	HIS
1	A	177	GLN
1	A	182	SER
1	A	265	THR
1	A A	294	LEU
1	A	327	ARG
1	В	37	GLN
1	В	105	ARG
1	В	113	CYS
1	В	115	THR
1	В	120	HIS
1	В	137	LEU
1	В	160	HIS
1	В	265	THR
1	В	294	LEU
1	В	311	THR
1	В	327	ARG
1	В	335	ASP
1	C C C	37	GLN
1	С	51	GLU
1	С	105	ARG
1	С	120	HIS



Continued from previous page...

Mol	Chain	Res	Type
1	С	160	HIS
1	С	168	LEU
1	С	177	GLN
1	С	213	ASP
1	С	294	LEU
1	С	300	GLN
1	С	321	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	177	GLN
1	В	325	GLN
1	С	228	GLN
1	С	284	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 6 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Type Chain	Pog	Link	Bond lengths		Bond angles			
IVIOI			rtes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	A1IFX	С	403	2	10,14,14	0.81	0	13,19,19	1.11	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	$\operatorname{Res}$	Link	Chirals	Torsions	Rings
3	A1IFX	С	403	2	-	3/14/17/17	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	Atoms	${f Z}$	$Observed(^o)$	$Ideal(^{o})$
3	С	403	A1IFX	O5-C1-O6	-2.84	117.65	124.09

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	С	403	A1IFX	C2-C3-C4-C10
3	С	403	A1IFX	C1-C2-C3-C4
3	С	403	A1IFX	C8-C3-C4-C10

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	A	318/336~(94%)	0.93	23 (7%) 23 25	39, 52, 90, 141	0
1	В	324/336~(96%)	1.49	91 (28%) 2 1	42, 73, 136, 156	0
1	С	318/336 (94%)	0.97	23 (7%) 23 25	34, 52, 89, 125	0
All	All	960/1008 (95%)	1.13	137 (14%) 7 7	34, 57, 118, 156	0

All (137) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	В	71	LEU	4.9	
1	С	33	PHE	4.8	
1	A	337	LEU	4.5	
1	В	92	VAL	4.5	
1	В	86	ILE	4.2	
1	В	106	ALA	4.1	
1	A	338	PRO	4.1	
1	A	26	VAL	4.0	
1	A	339	THR	3.9	
1	В	107	VAL	3.9	
1	В	191	ALA	3.9	
1	В	302	ALA	3.8	
1	В	329	GLY	3.7	
1	В	99	LEU	3.6	
1	В	76	VAL	3.6	
1	В	111	TYR	3.5	
1	В	33	PHE	3.5	
1	В	308	ALA	3.4	
1	В	157	GLY	3.4	
1	В	36	GLY	3.4	
1	В	68	PHE	3.2	
1	В	171	LEU	3.2	
1	В	113	CYS	3.2	



Continued from previous page.

Mol 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Chain B B	238	Type LEU	3.2	
1 1 1				3.2	
1 1		188	ILE	3.2	
1	В	207	PHE	3.1	
	В	152	LEU	3.1	
1	В	168	LEU	3.1	
1	С	86	ILE	3.1	
1	В	87	VAL	3.1	
1	A	332	ILE	3.0	
1	В	333	VAL	3.0	
1	В	301	LEU	3.0	
1	В	184	ILE	3.0	
1	A	94	LEU	3.0	
1	В	193	ILE	2.9	
1	В	264	GLY	2.9	
1	В	42	VAL	2.9	
1	В	98	GLU	2.9	
1	В	297	VAL	2.9	
1	В	31	ALA	2.8	
1	В	181	PHE	2.8	
1	A	111	TYR	2.8	
1	A	237	LEU	2.8	
1	В	151	PRO	2.8	
1	В	94	LEU	2.8	
1	В	209	LEU	2.8	
1	В	289	LEU	2.8	
1	В	74	THR	2.8	
1	A	340	PRO	2.8	
1	В	166	PHE	2.7	
1	В	29	ILE	2.7	
1	В	49	ILE	2.7	
1	В	176	PRO	2.7	
1	В	82	TYR	2.7	
1	В	195	TYR	2.7	
1	A	25	SER	2.7	
1	В	85	LEU	2.7	
1	В	81	LEU	2.6	
1	В	103	VAL	2.6	
1	В	237	LEU	2.6	
1	В	93	GLY	2.6	
1	A	188	ILE	2.6	
1	В	122	LEU	2.6	
1	A	255	PHE d on nex	2.5	



Continued from previous page...

Continued from previous page							
Mol	Chain	Res	Type	RSRZ			
1	В	135	PRO	2.5			
1	В	178	LEU	2.5			
1	В	314	LEU	2.5			
1	В	102	VAL	2.5			
1	В	194	VAL	2.5			
1	В	299	PRO	2.5			
1	В	134	CYS	2.5			
1	A	262	ALA	2.5			
1	В	150	THR	2.5			
1	С	279	ILE	2.5			
1	В	140	VAL	2.5			
1	В	153	THR	2.5			
1	В	89	PRO	2.4			
1	В	149	ASN	2.4			
1	В	28	VAL	2.4			
1	A	144	ALA	2.4			
1	В	340	PRO	2.4			
1	В	146	ALA	2.4			
1	A	108	SER	2.4			
1	В	137	LEU	2.3			
1	С	102	VAL	2.3			
1	С	140	VAL	2.3			
1	С	290	SER	2.3			
1	В	174	LYS	2.3			
1	С	81	LEU	2.3			
1	В	159	LEU	2.3			
1	В	141	TRP	2.3			
1	В	130	HIS	2.3			
1	В	54	LEU	2.3			
1	В	311	THR	2.3			
1	В	164	VAL	2.2			
1	В	167	LEU	2.2			
1	В	179	PRO	2.2			
1	В	84	ASN	2.2			
1	С	188	ILE	2.2			
1	A	54	LEU	2.2			
1	A	301	LEU	2.2			
1	С	137	LEU	2.2			
1	В	212	TYR	2.2			
1	С	320	ALA	2.2			
1	С	297	VAL	2.2			
1	С	85	LEU	2.2			
		1					



Continued from previous page...

Mol	Chain	Res	Type	RSRZ	
1	A	334	TYR	2.2	
1	С	82	TYR	2.2	
1	В	100	ALA	2.2	
1	В	260	ALA	2.2	
1	В	117	GLY	2.2	
1	С	51	GLU	2.1	
1	В	187	CYS	2.1	
1	С	336	GLN	2.1	
1	С	26	VAL	2.1	
1	С	87	VAL	2.1	
1	С	124	ILE	2.1	
1	В	190	SER	2.1	
1	В	263	THR	2.1	
1	С	276	GLY	2.1	
1	В	139	VAL	2.1	
1	A	80	ASP	2.1	
1	A	166	PHE	2.1	
1	С	116	LEU	2.1	
1	С	222	ILE	2.1	
1	В	95	ALA	2.1	
1	В	183	TRP	2.1	
1	С	42	VAL	2.1	
1	В	108	SER	2.1	
1	С	227	ILE	2.1	
1	A	82	TYR	2.0	
1	В	148	ILE	2.0	
1	В	323	PHE	2.0	
1	A	204	PRO	2.0	
1	В	52	ALA	2.0	
1	A	42	VAL	2.0	

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	A1IFX	С	403	15/15	0.83	0.11	47,53,55,56	0
2	MN	В	501	1/1	0.92	0.06	61,61,61,61	0
2	MN	A	501	1/1	0.92	0.06	50,50,50,50	0
2	MN	В	500	1/1	0.93	0.05	65,65,65,65	0
2	MN	С	402	1/1	0.96	0.04	49,49,49,49	0
2	MN	A	500	1/1	0.96	0.05	53,53,53,53	0
2	MN	С	401	1/1	0.97	0.04	52,52,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers (i)

There are no such residues in this entry.

