



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 09:54 am BST

PDB ID : 2FTW
Title : Crystal structure of dihydropyrimidinase from dictyostelium discoideum
Authors : Lohkamp, B.; Dobritzsch, D.
Deposited on : 2006-01-25
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called dihydropyrimidine amidohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	484	3818	2422	668	704	24	9	14	0


There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	158	KCX	LYS	MODIFIED RESIDUE	UNP Q86LT2
A	504	PRO	-	EXPRESSION TAG	UNP Q86LT2
A	505	GLY	-	EXPRESSION TAG	UNP Q86LT2
A	506	ASP	-	EXPRESSION TAG	UNP Q86LT2
A	507	ASP	-	EXPRESSION TAG	UNP Q86LT2
A	508	ASP	-	EXPRESSION TAG	UNP Q86LT2
A	509	ASP	-	EXPRESSION TAG	UNP Q86LT2
A	510	LYS	-	EXPRESSION TAG	UNP Q86LT2
A	511	HIS	-	EXPRESSION TAG	UNP Q86LT2
A	512	HIS	-	EXPRESSION TAG	UNP Q86LT2
A	513	HIS	-	EXPRESSION TAG	UNP Q86LT2
A	514	HIS	-	EXPRESSION TAG	UNP Q86LT2
A	515	HIS	-	EXPRESSION TAG	UNP Q86LT2
A	516	HIS	-	EXPRESSION TAG	UNP Q86LT2
A	517	HIS	-	EXPRESSION TAG	UNP Q86LT2
A	518	HIS	-	EXPRESSION TAG	UNP Q86LT2
A	519	SER	-	EXPRESSION TAG	UNP Q86LT2
A	520	GLY	-	EXPRESSION TAG	UNP Q86LT2
A	521	ASP	-	EXPRESSION TAG	UNP Q86LT2

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 3 4	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	280	Total O 280 280	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	84.60Å 89.60Å 134.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.05 24.99 – 2.05	Depositor EDS
% Data completeness (in resolution range)	93.9 (25.00-2.05) 93.9 (24.99-2.05)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.04Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.204 , 0.252 0.212 , 0.258	Depositor DCC
R_{free} test set	1557 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtrriage
Anisotropy	0.163	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4107	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MLI, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	1/3929 (0.0%)	0.62	0/5321

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	159	CYS	CB-SG	-5.72	1.72	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3818	0	3797	25	0
2	A	2	0	0	0	0
3	A	7	0	2	1	0
4	A	280	0	0	2	0
All	All	4107	0	3799	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:251[B]:ILE:HD11	1:A:306:THR:HA	1.57	0.86
1:A:328:THR:HB	1:A:422:VAL:HG11	1.62	0.81
1:A:64:ASP:OD2	1:A:324[A]:THR:HG22	2.00	0.60
1:A:242:PRO:HD3	1:A:474:ILE:CD1	2.35	0.56
1:A:9[A]:THR:HG22	1:A:32:ASN:H	1.71	0.55
1:A:288:TRP:CZ2	1:A:338:LYS:HA	2.43	0.54
1:A:324[A]:THR:HG23	1:A:326:ASN:H	1.73	0.53
1:A:455:VAL:HB	1:A:458:SER:OG	2.11	0.51
1:A:84[A]:ILE:HG21	1:A:421:ALA:HB2	1.94	0.50
1:A:370:TRP:CD2	3:A:524:MLI:H12	2.47	0.49
1:A:471:PHE:HA	1:A:474:ILE:HD12	1.94	0.48
1:A:207:GLU:HG3	4:A:722:HOH:O	2.15	0.46
1:A:289:ARG:NH1	4:A:721:HOH:O	2.49	0.46
1:A:375:ARG:HG2	1:A:380:GLU:HB2	1.99	0.44
1:A:182[A]:LYS:NZ	1:A:239:VAL:O	2.43	0.44
1:A:49[B]:LYS:HE2	1:A:445:ILE:HD12	1.99	0.44
1:A:64:ASP:OD2	1:A:324[A]:THR:CG2	2.64	0.42
1:A:10:ILE:HD11	1:A:51:VAL:CG2	2.50	0.42
1:A:406:ASP:O	1:A:434:GLY:HA2	2.19	0.41
1:A:80:ASP:HA	1:A:84[B]:ILE:CD1	2.50	0.41
1:A:279:ASP:OD1	1:A:279:ASP:C	2.58	0.41
1:A:249:GLN:HG3	1:A:298:PRO:HG2	2.02	0.41
1:A:251[B]:ILE:HD12	1:A:309:VAL:HB	2.02	0.41
1:A:117:LYS:NZ	1:A:121:ASP:OD2	2.53	0.40
1:A:255:ASP:OD1	1:A:313:TYR:OH	2.29	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	495/521 (95%)	473 (96%)	21 (4%)	1 (0%)	47 39

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	350	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	413/444 (93%)	405 (98%)	8 (2%)	57 53

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	SER
1	A	40	LYS
1	A	77	VAL
1	A	116	TRP
1	A	148	LEU
1	A	289	ARG
1	A	346	ASN
1	A	418	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	290	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	A	158	1,2	7,11,12	1.12	1 (14%)	4,12,14	1.26	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	158	1,2	-	4/7/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	158	KCX	CE-NZ	2.23	1.50	1.45

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	KCX	CE-NZ-CX	-2.26	119.12	122.95

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	158	KCX	CA-CB-CG-CD
1	A	158	KCX	N-CA-CB-CG
1	A	158	KCX	CE-CD-CG-CB
1	A	158	KCX	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MLI	A	524	-	0,6,6	0.00	-	0,7,7	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLI	A	524	-	-	0/0/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	524	MLI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	483/521 (92%)	0.23	10 (2%) 63 67	18, 25, 34, 42	11 (2%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	7	THR	4.7
1	A	489	ASP	4.5
1	A	75	GLY	3.4
1	A	9[A]	THR	3.1
1	A	74	MET	2.8
1	A	48	ILE	2.6
1	A	295	VAL	2.5
1	A	46	GLU	2.3
1	A	331	ALA	2.0
1	A	166	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	A	158	12/13	0.92	0.17	22,23,26,26	3

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	A	523	1/1	0.65	0.18	36,36,36,36	1
3	MLI	A	524	7/7	0.78	0.20	30,34,38,40	0
2	ZN	A	522	1/1	0.97	0.07	41,41,41,41	1

6.5 Other polymers [i](#)

There are no such residues in this entry.