



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2023 – 06:45 PM EDT

PDB ID : 2FU5
Title : structure of Rab8 in complex with MSS4
Authors : Itzen, A.; Pylypenko, O.; Goody, R.S.; Rak, A.
Deposited on : 2006-01-26
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	117	 3% 83% 8% • 9%
1	B	117	 3% 83% 7% • 9%
2	C	183	 19% 62% 16% • 20%
2	D	183	 17% 74% 11% • 14%

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4473 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide exchange factor MSS4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	107	841	530	149	154	8	0	0	0
1	B	107	847	533	152	154	8	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	GLN	-	cloning artifact	UNP P47224
A	8	GLY	-	cloning artifact	UNP P47224
A	9	HIS	-	cloning artifact	UNP P47224
A	10	MET	-	cloning artifact	UNP P47224
B	7	GLN	-	cloning artifact	UNP P47224
B	8	GLY	-	cloning artifact	UNP P47224
B	9	HIS	-	cloning artifact	UNP P47224
B	10	MET	-	cloning artifact	UNP P47224

- Molecule 2 is a protein called Ras-related protein Rab-8A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	146	1190	762	202	221	5	0	0	0
2	D	157	1284	820	220	239	5	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	98	Total	O	0	0
			98	98		
5	B	114	Total	O	0	0
			114	114		
5	C	40	Total	O	0	0
			40	40		
5	D	49	Total	O	0	0
			49	49		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Guanine nucleotide exchange factor MSS4




- Molecule 1: Guanine nucleotide exchange factor MSS4



- Molecule 2: Ras-related protein Rab-8A



- Molecule 2: Ras-related protein Rab-8A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	40.92Å 49.85Å 83.48Å 102.88° 97.46° 90.12°	Depositor
Resolution (Å)	19.17 – 2.00 19.17 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (19.17-2.00) 94.7 (19.17-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 2.01Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.197 , 0.252 0.198 , 0.250	Depositor DCC
R_{free} test set	2036 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	24.5	Xtrriage
Anisotropy	0.394	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 60.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4473	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BME, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/857	0.79	1/1152 (0.1%)
1	B	0.79	0/863	0.72	0/1159
2	C	0.50	0/1205	0.63	0/1610
2	D	0.50	0/1301	0.59	0/1740
All	All	0.65	0/4226	0.67	1/5661 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	18	ARG	NE-CZ-NH1	5.35	122.97	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	841	0	804	5	0
1	B	847	0	815	6	0
2	C	1190	0	1203	22	0
2	D	1284	0	1283	12	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	4	0	5	0	0
4	B	4	0	5	0	0
5	A	98	0	0	1	0
5	B	114	0	0	2	0
5	C	40	0	0	0	0
5	D	49	0	0	1	0
All	All	4473	0	4115	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:150:THR:HB	2:D:157:ASN:HA	1.82	0.61
2:C:7:TYR:HE1	2:C:55:LYS:HG2	1.64	0.61
2:D:100:ARG:NH2	5:D:215:HOH:O	2.34	0.60
1:A:110:LYS:HG3	5:A:4359:HOH:O	2.02	0.60
2:D:140:ALA:HB1	2:D:145:ILE:O	2.00	0.59
1:B:99:ILE:HD12	5:B:4349:HOH:O	2.03	0.59
1:A:97:CYS:SG	1:A:99:ILE:HD12	2.44	0.58
2:C:157:ASN:HB3	2:C:160:ASN:ND2	2.22	0.55
2:D:87:VAL:HG12	2:D:119:LEU:HB3	1.88	0.55
2:C:13:LEU:HD22	2:C:162:PHE:HZ	1.73	0.53
2:D:99:ILE:O	2:D:103:ILE:HG13	2.09	0.52
1:A:90:LYS:HD2	1:A:107:LEU:HD21	1.92	0.52
2:C:17:SER:HB2	2:C:122:LYS:NZ	2.25	0.51
2:C:13:LEU:CD2	2:C:162:PHE:HZ	2.24	0.51
2:C:92:ASN:HD21	2:C:94:LYS:HB3	1.76	0.50
2:C:49:THR:OG1	2:C:58:LYS:HE3	2.11	0.49
2:C:7:TYR:CD1	2:C:57:ILE:CD1	2.96	0.49
1:B:45:LEU:HD13	1:B:65:LEU:HD12	1.94	0.48
1:B:49:ARG:HG3	5:B:4417:HOH:O	2.14	0.48
2:C:13:LEU:CD2	2:C:162:PHE:CZ	2.97	0.48
2:C:17:SER:HB2	2:C:122:LYS:HZ1	1.78	0.48
2:C:106:ILE:HG21	2:C:116:LYS:HD3	1.95	0.48
1:A:12:SER:HA	1:A:18:ARG:NH2	2.30	0.46
2:D:17:SER:HB3	2:D:89:ASP:HB2	1.98	0.46
1:A:93:VAL:HG12	1:A:100:GLY:HA3	1.99	0.45
2:D:69:ARG:HG3	2:D:71:ARG:O	2.16	0.45
2:C:103:ILE:H	2:C:103:ILE:HG13	1.58	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:36:THR:O	2:D:40:THR:HG23	2.18	0.43
2:C:13:LEU:HD22	2:C:162:PHE:CZ	2.52	0.43
2:C:103:ILE:HD11	2:C:139:LEU:HD21	2.01	0.43
2:D:157:ASN:ND2	2:D:160:ASN:ND2	2.66	0.43
1:B:16:ARG:HA	1:B:36:ALA:O	2.19	0.43
2:C:103:ILE:HA	2:C:106:ILE:HD12	2.00	0.42
2:C:103:ILE:CD1	2:C:139:LEU:HD11	2.49	0.42
2:D:157:ASN:ND2	2:D:160:ASN:HD22	2.17	0.42
2:C:140:ALA:HB2	2:C:147:PHE:HB2	2.01	0.42
2:C:13:LEU:HD21	2:C:162:PHE:CZ	2.55	0.41
2:D:87:VAL:O	2:D:99:ILE:HD11	2.20	0.41
1:B:79:ASN:HA	2:C:47:ILE:O	2.20	0.41
2:C:94:LYS:HA	2:C:94:LYS:HD2	1.92	0.41
2:C:122:LYS:HB2	2:C:122:LYS:HZ2	1.86	0.41
1:B:72:GLU:HA	1:B:111:ASN:O	2.20	0.40
2:D:15:GLY:O	2:D:87:VAL:HG23	2.21	0.40
2:C:36:THR:HG22	2:C:37:PHE:H	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/117 (88%)	101 (98%)	2 (2%)	0	100	100
1	B	103/117 (88%)	102 (99%)	1 (1%)	0	100	100
2	C	136/183 (74%)	127 (93%)	6 (4%)	3 (2%)	6	2
2	D	147/183 (80%)	141 (96%)	6 (4%)	0	100	100
All	All	489/600 (82%)	471 (96%)	15 (3%)	3 (1%)	25	19

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	76	ALA
2	C	176	ASN
2	C	128	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	90/101 (89%)	89 (99%)	1 (1%)	73	78
1	B	91/101 (90%)	88 (97%)	3 (3%)	38	37
2	C	127/155 (82%)	118 (93%)	9 (7%)	14	10
2	D	135/155 (87%)	128 (95%)	7 (5%)	23	19
All	All	443/512 (86%)	423 (96%)	20 (4%)	27	24

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ARG
1	B	7	GLN
1	B	49	ARG
1	B	72	GLU
2	C	36	THR
2	C	64	THR
2	C	77	TYR
2	C	78	TYR
2	C	79	ARG
2	C	103	ILE
2	C	119	LEU
2	C	121	ASN
2	C	127	ASP
2	D	34	ASN
2	D	35	SER
2	D	69	ARG
2	D	98	ASN
2	D	111	SER
2	D	113	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	157	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	7	GLN
2	C	92	ASN
2	C	121	ASN
2	C	160	ASN
2	D	34	ASN
2	D	105	ASN
2	D	109	HIS
2	D	130	GLN
2	D	155	ASN
2	D	157	ASN
2	D	160	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BME	A	4322	1	3,3,3	0.18	0	1,2,2	1.09	0
4	BME	B	4321	1	3,3,3	0.40	0	1,2,2	1.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BME	A	4322	1	-	0/1/1/1	-
4	BME	B	4321	1	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	107/117 (91%)	-0.07	4 (3%) 41 41	19, 27, 42, 61	0
1	B	107/117 (91%)	-0.12	3 (2%) 53 51	19, 27, 42, 59	0
2	C	146/183 (79%)	1.28	34 (23%) 0 0	28, 52, 81, 86	0
2	D	157/183 (85%)	1.16	32 (20%) 1 0	24, 55, 75, 119	0
All	All	517/600 (86%)	0.67	73 (14%) 2 2	19, 42, 75, 119	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	125	VAL	10.4
2	C	74	THR	8.3
2	C	177	TRP	7.6
2	C	19	VAL	7.5
2	D	177	TRP	7.5
2	C	151	SER	6.9
2	D	155	ASN	6.5
2	C	156	ILE	6.4
2	C	128	LYS	6.3
2	C	109	HIS	5.9
2	C	77	TYR	5.9
2	D	66	GLY	5.7
2	D	112	ALA	5.7
2	C	127	ASP	5.5
2	D	178	LYS	5.2
2	D	176	ASN	5.0
2	D	32	ALA	4.8
2	C	152	ALA	4.7
2	D	124	ASP	4.5
2	C	110	ALA	4.4
2	D	34	ASN	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	65	ALA	4.2
2	C	18	GLY	4.2
2	D	17	SER	4.2
2	D	33	PHE	4.1
2	D	154	ALA	3.9
2	C	125	VAL	3.9
2	D	127	ASP	3.8
2	D	156	ILE	3.8
2	C	126	ASN	3.7
2	D	35	SER	3.7
2	C	16	ASP	3.6
2	C	113	ASP	3.5
2	D	110	ALA	3.5
2	C	17	SER	3.4
2	C	111	SER	3.3
2	C	79	ARG	3.3
2	C	75	THR	3.2
2	C	78	TYR	3.2
2	C	102	TRP	3.2
2	D	71	ARG	3.2
2	D	15	GLY	3.2
1	B	50	LYS	3.1
2	D	74	THR	3.1
2	C	76	ALA	3.1
2	C	112	ALA	3.1
2	C	107	GLU	3.0
1	A	7	GLN	2.9
1	B	49	ARG	2.9
2	C	106	ILE	2.9
2	D	128	LYS	2.9
1	B	61	ASP	2.8
2	D	119	LEU	2.8
2	D	77	TYR	2.8
2	C	175	LYS	2.7
2	C	88	TYR	2.6
2	D	109	HIS	2.6
2	C	176	ASN	2.6
1	A	50	LYS	2.5
2	D	68	GLU	2.5
2	D	94	LYS	2.5
2	C	159	GLU	2.4
2	C	3	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	79	ARG	2.4
2	D	113	ASP	2.4
2	D	86	LEU	2.4
1	A	61	ASP	2.4
1	A	8	GLY	2.3
2	C	157	ASN	2.3
2	D	75	THR	2.2
2	C	120	GLY	2.2
2	D	118	ILE	2.1
2	C	108	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BME	A	4322	4/4	0.96	0.11	38,39,41,45	0
4	BME	B	4321	4/4	0.97	0.09	38,39,39,41	0
3	ZN	A	502	1/1	1.00	0.08	24,24,24,24	0
3	ZN	B	501	1/1	1.00	0.08	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.