



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 30, 2023 – 01:42 PM EST

PDB ID : 8FUY
Title : Glucose-6-phosphate 1-dehydrogenase (G6PDH) from *Crithidia fasciculata* (citrate bound)
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID); Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2023-01-18
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.32.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	568	 4% 76% 9% 15%
1	B	568	 4% 77% 9% 14%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CIT	A	602	-	-	-	X

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 7647 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucose-6-phosphate 1-dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	485	Total 3795	C 2416	N 652	O 712	S 15	0	0	0
1	B	486	Total 3788	C 2408	N 651	O 714	S 15	0	0	0

- Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 13	C 6	O 7	0	0
2	A	1	Total 13	C 6	O 7	0	0
2	B	1	Total 13	C 6	O 7	0	0
2	B	1	Total 13	C 6	O 7	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	B	1	Total Cl 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total O 4 4	0	0
4	B	6	Total O 6 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glucose-6-phosphate 1-dehydrogenase



- Molecule 1: Glucose-6-phosphate 1-dehydrogenase



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	121.51Å 124.91Å 161.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.55 – 2.45 87.10 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.8 (43.55-2.45) 99.9 (87.10-2.45)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 2.45Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.203 , 0.227 0.201 , 0.223	Depositor DCC
R_{free} test set	2188 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	58.1	Xtrriage
Anisotropy	0.600	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 59.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.021 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7647	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3876	0.52	0/5244
1	B	0.29	0/3869	0.53	0/5238
All	All	0.29	0/7745	0.53	0/10482

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3795	0	3732	39	0
1	B	3788	0	3707	40	0
2	A	26	0	10	1	0
2	B	26	0	10	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	0	0	0
4	B	6	0	0	0	0
All	All	7647	0	7459	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:117:ASP:HB3	1:B:120:LYS:HB2	1.61	0.81
1:A:100:LEU:HD12	1:B:455:LEU:CD1	2.14	0.78
1:A:100:LEU:CD1	1:B:455:LEU:CD1	2.65	0.74
1:A:100:LEU:CD1	1:B:455:LEU:HD12	2.22	0.70
1:B:384:VAL:HG22	1:B:402:LYS:HG3	1.73	0.69
1:A:70:ARG:HH12	1:A:180:ALA:HB3	1.58	0.68
1:A:100:LEU:CD1	1:B:455:LEU:HD13	2.26	0.66
1:B:119:GLU:O	1:B:123:HIS:ND1	2.33	0.61
1:A:446:MET:HE2	1:A:467:LEU:HD12	1.82	0.61
1:A:100:LEU:HD12	1:B:455:LEU:HD13	1.84	0.59
1:A:119:GLU:O	1:A:123:HIS:ND1	2.36	0.59
1:A:191:PRO:HG3	1:A:220:PRO:HD2	1.85	0.58
1:A:100:LEU:HD11	1:B:455:LEU:HD12	1.86	0.58
1:B:359:SER:HB3	1:B:364:ILE:HB	1.86	0.58
1:B:172:ASP:HA	1:B:179:LYS:HD2	1.86	0.57
1:A:180:ALA:HA	1:A:208:ARG:HG3	1.86	0.56
1:A:80:SER:HA	1:A:121:TRP:HH2	1.72	0.55
1:A:100:LEU:HD12	1:B:455:LEU:HD12	1.83	0.55
1:B:117:ASP:OD1	1:B:120:LYS:NZ	2.37	0.54
1:B:481:ASP:HB3	1:B:484:GLU:HG3	1.90	0.53
1:A:433:GLU:HB2	1:A:447:LYS:HD3	1.90	0.53
1:B:77:LEU:HA	1:B:110:TYR:HB3	1.90	0.53
1:A:484:GLU:OE2	1:B:455:LEU:HD11	2.09	0.51
1:B:402:LYS:HD2	2:B:602:CIT:O3	2.10	0.51
1:B:284:THR:HG23	1:B:402:LYS:HD3	1.91	0.51
1:B:359:SER:O	1:B:361:ASP:N	2.43	0.51
1:B:500:ARG:HD2	1:B:502:ASP:OD1	2.10	0.51
1:A:63:VAL:HG13	1:A:67:GLN:HB2	1.92	0.50
1:B:122:LYS:HE3	1:B:147:VAL:HB	1.93	0.50
1:A:60:VAL:O	1:A:62:LYS:HE3	2.11	0.49
1:A:137:SER:HA	1:A:140:VAL:HG23	1.95	0.49
1:A:123:HIS:NE2	1:A:141:GLU:OE2	2.35	0.48
1:A:348:LYS:HD2	1:A:517:GLN:NE2	2.29	0.47
1:B:148:THR:HG22	1:B:149:TYR:N	2.30	0.47
1:B:180:ALA:HA	1:B:208:ARG:HG3	1.97	0.46
1:A:384:VAL:HG22	1:A:402:LYS:HG3	1.96	0.46
1:B:285:PHE:O	1:B:403:ALA:HA	2.14	0.46
1:A:174:PHE:O	1:A:179:LYS:NZ	2.46	0.46
1:B:123:HIS:NE2	1:B:141:GLU:OE2	2.50	0.45
1:B:124:GLU:HG3	1:B:125:THR:HG23	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:LYS:HB3	1:A:144:LEU:HD22	1.98	0.45
1:A:286:LYS:HB2	1:A:411:TYR:HB3	1.97	0.45
1:B:250:TYR:HA	1:B:253:LYS:HG3	1.99	0.45
1:B:172:ASP:O	1:B:179:LYS:NZ	2.41	0.44
1:B:280:CYS:HB2	1:B:547:PHE:CE1	2.52	0.44
1:B:415:ARG:HE	2:B:602:CIT:C1	2.29	0.44
1:A:77:LEU:HA	1:A:110:TYR:HB3	2.00	0.44
1:B:402:LYS:NZ	2:B:602:CIT:O3	2.37	0.44
1:A:421:GLU:OE2	1:B:323:LYS:NZ	2.38	0.44
1:B:126:LEU:HD23	1:B:126:LEU:HA	1.82	0.43
1:A:417:GLN:HG2	1:A:433:GLU:HG2	2.00	0.43
1:B:56:CYS:HB3	1:B:57:GLU:H	1.65	0.43
1:A:111:ALA:O	1:A:151:THR:HA	2.19	0.42
1:A:127:THR:HG22	1:A:130:PHE:CE2	2.53	0.42
1:A:465:THR:HG21	1:B:469:LEU:HD11	2.01	0.42
1:B:310:ASN:OD1	1:B:310:ASN:N	2.53	0.42
1:A:208:ARG:H	1:A:208:ARG:HG2	1.58	0.42
1:A:148:THR:HG22	1:A:149:TYR:N	2.35	0.42
1:A:254:GLU:OE2	1:B:453:PRO:HA	2.21	0.41
1:A:451:LYS:HE2	1:A:451:LYS:HB2	1.81	0.41
1:A:273:TRP:O	1:A:393:ARG:NH1	2.51	0.41
1:B:317:ALA:HA	1:B:339:LEU:CD2	2.51	0.41
1:A:112:ARG:O	1:A:151:THR:HG23	2.20	0.41
1:A:172:ASP:HA	1:A:179:LYS:HD2	2.01	0.41
1:B:208:ARG:H	1:B:208:ARG:HG2	1.66	0.41
1:A:254:GLU:OE2	1:B:451:LYS:NZ	2.51	0.41
2:A:601:CIT:O3	2:A:601:CIT:O7	2.23	0.40
1:A:70:ARG:NH1	1:A:180:ALA:HB3	2.30	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	481/568 (85%)	468 (97%)	13 (3%)	0	100	100
1	B	482/568 (85%)	461 (96%)	21 (4%)	0	100	100
All	All	963/1136 (85%)	929 (96%)	34 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	400/493 (81%)	396 (99%)	4 (1%)	76	84
1	B	398/493 (81%)	392 (98%)	6 (2%)	65	76
All	All	798/986 (81%)	788 (99%)	10 (1%)	69	79

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	CYS
1	A	208	ARG
1	A	415	ARG
1	A	463	HIS
1	B	115	GLN
1	B	139	HIS
1	B	208	ARG
1	B	351	CYS
1	B	363	SER
1	B	463	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CIT	B	601	-	12,12,12	1.12	0	17,17,17	1.32	2 (11%)
2	CIT	B	602	-	12,12,12	1.10	0	17,17,17	1.30	2 (11%)
2	CIT	A	602	-	12,12,12	1.12	0	17,17,17	1.30	2 (11%)
2	CIT	A	601	-	12,12,12	1.01	0	17,17,17	1.66	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	B	601	-	-	3/16/16/16	-
2	CIT	B	602	-	-	13/16/16/16	-
2	CIT	A	602	-	-	9/16/16/16	-
2	CIT	A	601	-	-	6/16/16/16	-

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	CIT	O6-C6-C3	4.20	120.35	113.05
2	B	602	CIT	O5-C6-C3	-3.57	117.19	122.25
2	B	601	CIT	O5-C6-C3	-3.57	117.20	122.25
2	A	602	CIT	O5-C6-C3	-3.53	117.25	122.25
2	B	601	CIT	O6-C6-C3	2.48	117.36	113.05
2	A	602	CIT	O6-C6-C3	2.47	117.33	113.05
2	A	601	CIT	C3-C4-C5	-2.46	107.86	113.81
2	B	602	CIT	O6-C6-C3	2.42	117.25	113.05
2	A	601	CIT	O4-C5-C4	2.08	121.02	114.35

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	CIT	O7-C3-C4-C5
2	A	602	CIT	C2-C3-C4-C5
2	A	602	CIT	O7-C3-C4-C5
2	A	602	CIT	C6-C3-C4-C5
2	A	602	CIT	O7-C3-C6-O5
2	A	602	CIT	O7-C3-C6-O6
2	A	602	CIT	C4-C3-C6-O5
2	A	602	CIT	C4-C3-C6-O6
2	B	602	CIT	O7-C3-C6-O5
2	B	602	CIT	O7-C3-C6-O6
2	B	602	CIT	C4-C3-C6-O5
2	B	602	CIT	C4-C3-C6-O6
2	A	601	CIT	C2-C3-C4-C5
2	A	601	CIT	C1-C2-C3-C6
2	A	601	CIT	C6-C3-C4-C5
2	A	601	CIT	C1-C2-C3-O7
2	B	602	CIT	O7-C3-C4-C5
2	B	602	CIT	C6-C3-C4-C5
2	B	601	CIT	C1-C2-C3-C6
2	B	602	CIT	C2-C3-C4-C5
2	B	602	CIT	C1-C2-C3-C6
2	B	601	CIT	C1-C2-C3-O7
2	B	602	CIT	C1-C2-C3-O7
2	B	601	CIT	C1-C2-C3-C4
2	A	602	CIT	C1-C2-C3-C4
2	A	602	CIT	C1-C2-C3-C6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	602	CIT	C1-C2-C3-C4
2	A	601	CIT	C1-C2-C3-C4
2	B	602	CIT	C3-C4-C5-O3
2	B	602	CIT	O2-C1-C2-C3
2	B	602	CIT	C3-C4-C5-O4

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	602	CIT	3	0
2	A	601	CIT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	485/568 (85%)	0.41	25 (5%) 27 24	42, 81, 132, 177	0
1	B	486/568 (85%)	0.46	23 (4%) 31 29	43, 80, 135, 191	0
All	All	971/1136 (85%)	0.44	48 (4%) 29 27	42, 80, 134, 191	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	127	THR	5.8
1	B	550	GLN	5.7
1	B	118	VAL	4.5
1	A	163	LEU	3.8
1	A	129	TYR	3.7
1	A	56	CYS	3.6
1	B	180	ALA	3.6
1	B	56	CYS	3.6
1	B	124	GLU	3.6
1	B	375	LYS	3.4
1	A	124	GLU	3.2
1	B	126	LEU	3.2
1	B	457	GLU	3.2
1	A	547	PHE	3.2
1	A	208	ARG	3.1
1	B	470	THR	3.1
1	A	160	PHE	3.1
1	B	371	GLU	3.1
1	A	121	TRP	3.0
1	A	149	TYR	2.9
1	A	550	GLN	2.9
1	B	128	LYS	2.8
1	B	139	HIS	2.8
1	A	526	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	226	GLU	2.7
1	B	174	PHE	2.7
1	B	117	ASP	2.7
1	A	382	PHE	2.6
1	A	131	SER	2.6
1	B	178	GLU	2.6
1	A	235	ILE	2.6
1	A	459	LEU	2.5
1	B	177	GLU	2.5
1	A	371	GLU	2.4
1	B	123	HIS	2.4
1	A	204	GLY	2.4
1	A	383	ALA	2.3
1	A	549	PHE	2.2
1	B	480	PRO	2.2
1	B	116	ASP	2.2
1	B	143	PHE	2.2
1	A	128	LYS	2.2
1	B	137	SER	2.1
1	A	356	TYR	2.1
1	A	158	GLU	2.1
1	A	69	ARG	2.1
1	A	201	ILE	2.0
1	A	229	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CL	B	603	1/1	0.55	0.17	100,100,100,100	0
2	CIT	A	601	13/13	0.69	0.24	94,104,111,116	0
2	CIT	A	602	13/13	0.72	0.40	82,95,104,108	0
2	CIT	B	601	13/13	0.74	0.26	88,103,110,111	0
2	CIT	B	602	13/13	0.85	0.21	78,83,97,98	0
3	CL	A	603	1/1	0.93	0.10	102,102,102,102	0

6.5 Other polymers [i](#)

There are no such residues in this entry.