



wwPDB EM Validation Summary Report ⓘ

Jan 2, 2025 – 03:19 pm GMT

PDB ID : 9FUX
EMDB ID : EMD-50781
Title : Cryo-EM structure of the Nipah virus polymerase (L) bound to the tetrameric phosphoprotein (P)
Authors : Balikci, E.; Gunl, F.; Carrique, L.; Keown, J.R.; Fodor, E.; Grimes, J.M.
Deposited on : 2024-06-26
Resolution : 2.49 Å(reported)
Based on initial model : .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2243	
2	C	708	
2	D	708	
2	E	708	
2	F	708	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14556 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase L.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1286	10357	6599	1769	1922	67	0	0

- Molecule 2 is a protein called Phosphoprotein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	C	116	927	580	160	181	6	1	0
2	D	108	861	545	143	166	7	0	0
2	E	198	1591	999	265	319	8	0	0
2	F	103	818	516	135	160	7	0	0


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
3	A	2	Total	Zn	0
			2	2	

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RNA-directed RNA polymerase L



V1362	V1370	V1371	M1372	S1373	L1377	L1381	E1382	L1391	V1392	Q1393	Q1394	M1396	L1397	L1398	G1399	Y1420	H1421	L1422	H1423	C1429	D1459	P1462	Y1463
LEU	HIS	ASP	VAL	LEU	ALA	LEU	THR	VAL	LEU	GLU	ILE	THR	THR	LYS	ALA	ASP	ASP	LEU	LYS	GLN	HIS	ASP	SER
SER	ILE	LYS	TRP	ALA	GLU	HIS	ARG	PRO	ARG	GLY	ASN	HIS	MET	VAL	ASP	LEU	LEU	LEU	LEU	VAL	HIS	LYS	SER
GLN	PRO	TYR	LEU	HIS	GLN	ASP	PHE	LYS	SER	GLN	ASN	LEU	VAL	THR	LEU	MET	ASN	LEU	LEU	ASN	ASN	PHE	SER
VAL	ILE	ASP	LEU	TYR	ALA	HIS	HIS	PRO	PRO	TRP	ILE	ASP	ASN	PRO	GLN	GLY	ARG	VAL	VAL	ILE	LYS	SER	ASP
LEU	ARG	ARG	ILE	ILE	LYS	GLN	LEU	ARG	VAL	THR	THR	THR	THR	THR	THR	ASN	THR	ASP	ASP	ASN	PRO	PRO	LYS
ALA	SER	LYS	GLY	PHE	ASN	ASN	ARG	ARG	VAL	ARG	ASN	HIS	TYR	ARG	ARG	SER	ALA	ALA	ALA	SER	ASN	ALA	LEU
LEU	TYR	GLN	THR	THR	LEU	GLY	SER	ILE	PHE	TYR	ASN	GLY	ASP	GLY	ASP	GLY	PHE	PRO	ILE	ALA	ASN	GLY	ARG
PRO	GLU	THR	TRP	ILE	ASN	ASP	THR	TYR	PRO	TYR	ILE	ARG	THR	ALA	VAL	GLY	GLY	VAL	VAL	GLU	GLY	VAL	ASP
GLY	LEU	VAL	SER	ILE	ALA	LEU	THR	PRO	ILE	ILE	ILE	ARG	PHE	MET	GLY	PHE	GLY	VAL	VAL	ASP	PRO	VAL	LEU
LYS	VAL	LEU	HIS	ASN	LEU	HIS	ASP	ASP	GLN	GLY	GLN	THR	VAL	ILE	SER	HIS	ASP	TYR	TYR	ASN	TYR	ASN	LEU
GLN	ALA	LEU	PRO	ASN	PRO	GLU	ILE	VAL	GLY	SER	THR	THR	ASP	ILE	LEU	GLN	ASP	ASP	ASP	ASN	MET	ALA	ARG
THR	ARG	ILE	GLY	THR	CYS	VAL	THR	LYS	VAL	THR	THR	ASP	ILE	LYS	PHE	THR	THR	THR	THR	THR	LYS	ARG	ARG
GLU	ARG	GLU	ASN	GLY	PHE	LYS	VAL	VAL	ASP	GLU	GLY	THR	GLU	VAL	VAL	ILE	ARG	ARG	ARG	ARG	ARG	ARG	ILE

● Molecule 2: Phosphoprotein



ASP	LYS	LEU	LEU	VAL	ASP	PHE	GLY	ASP	GLN	THR	GLN	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
GLY	MET	SER	LYS	ASP	GLY	VAL	VAL	ASP	SER	THR	SER	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
GLU	CYS	THR	GLY	TYR	GLY	THR	PRO	ARG	ASP	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
SER	THR	THR	GLY	ASN	PRO	THR	ALA	PHE	THR	ASP	PRO	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
ASP	GLU	ALA	ASP	GLN	LEU	PHE	GLY	ASP	GLY	VAL	ILE	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	490675	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1400	Depositor
Maximum defocus (nm)	2400	Depositor
Magnification	Not provided	
Image detector	TFS FALCON 4i (4k x 4k)	Depositor
Maximum map value	66.063	Depositor
Minimum map value	-54.750	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.908	Depositor
Recommended contour level	3	Depositor
Map size (\AA)	373.9136, 373.9136, 373.9136	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.7303, 0.7303, 0.7303	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.34	0/10571	0.53	3/14294 (0.0%)
2	C	0.26	0/941	0.45	0/1266
2	D	0.25	0/875	0.44	0/1180
2	E	0.26	0/1611	0.48	0/2176
2	F	0.25	0/831	0.48	0/1124
All	All	0.32	0/14829	0.51	3/20040 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	210	LEU	CA-CB-CG	5.87	128.81	115.30
1	A	242	LEU	CA-CB-CG	-5.85	101.85	115.30
1	A	1138	LEU	CA-CB-CG	5.57	128.11	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10357	0	10379	155	0
2	C	927	0	931	34	0
2	D	861	0	868	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	1591	0	1582	37	0
2	F	818	0	816	21	0
3	A	2	0	0	0	0
All	All	14556	0	14576	224	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 224 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:536:ILE:HD11	2:D:537:GLU:HG2	1.51	0.93
1:A:291:GLU:HG2	1:A:292:PRO:HD3	1.53	0.90
1:A:474:LYS:HD3	1:A:561:ARG:HH21	1.43	0.84
1:A:916:SER:OG	1:A:929:ASN:OD1	1.99	0.79
1:A:1147:SER:HA	1:A:1151:LYS:HD2	1.64	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1276/2243 (57%)	1219 (96%)	55 (4%)	2 (0%)	44	64
2	C	115/708 (16%)	108 (94%)	7 (6%)	0	100	100
2	D	106/708 (15%)	106 (100%)	0	0	100	100
2	E	192/708 (27%)	188 (98%)	4 (2%)	0	100	100
2	F	101/708 (14%)	101 (100%)	0	0	100	100
All	All	1790/5075 (35%)	1722 (96%)	66 (4%)	2 (0%)	50	69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1382	GLU
1	A	1145	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1156/2045 (56%)	1155 (100%)	1 (0%)	92	97
2	C	107/624 (17%)	107 (100%)	0	100	100
2	D	100/624 (16%)	100 (100%)	0	100	100
2	E	185/624 (30%)	185 (100%)	0	100	100
2	F	96/624 (15%)	96 (100%)	0	100	100
All	All	1644/4541 (36%)	1643 (100%)	1 (0%)	92	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	222	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-50781. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 245



Y Index: 237



Z Index: 222


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 146 nm³; this corresponds to an approximate mass of 132 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i



*Reported resolution corresponds to spatial frequency of 0.402 \AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-50781 and PDB model 9FUX. Per-residue inclusion information can be found in section 3 on page 4.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 3.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3).













9.4 Atom inclusion [i](#)



At the recommended contour level, 95% of all backbone atoms, 92% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9170	 0.5700
A	 0.9620	 0.6150
C	 0.7940	 0.4400
D	 0.7770	 0.4350
E	 0.8260	 0.4960
F	 0.8190	 0.4330

