



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2023 – 11:43 PM EDT

PDB ID : 8FV8
Title : E coli. CTP synthase in complex with dF-dCTP + ADP
Authors : Holyoak, T.; McLeod, M.J.; Tran, N.
Deposited on : 2023-01-18
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	545	
1	BBB	545	

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9216 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CTP synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	535	Total 4267	C 2699	N 742	O 801	S 25	0	14	0
1	BBB	535	Total 4231	C 2676	N 737	O 794	S 24	0	11	0

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			P
3	AAA	1	Total	C	F	N	O	P	0	0
			30	9	2	3	13	3		
3	AAA	1	Total	C	F	N	O	P	0	0
			30	9	2	3	13	3		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
4	AAA	1	Total	Na	0	0
			1	1		
4	BBB	1	Total	Na	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	AAA	266	Total	O	0	0
			266	266		
5	BBB	282	Total	O	0	0
			282	282		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CTP synthase



- Molecule 1: CTP synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	158.01Å 109.43Å 129.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.50 – 2.05 89.96 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.8 (83.50-2.05) 99.8 (89.96-2.05)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 2.05Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.197 , 0.216 0.204 , 0.221	Depositor DCC
R_{free} test set	6950 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	38.6	Xtrriage
Anisotropy	0.224	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9216	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTF, NA, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.64	0/4356	0.73	0/5899
1	BBB	0.64	0/4325	0.72	0/5857
All	All	0.64	0/8681	0.72	0/11756

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	4267	0	4289	16	0
1	BBB	4231	0	4266	17	0
2	AAA	54	0	24	2	0
2	BBB	54	0	24	4	0
3	AAA	60	0	20	0	0
4	AAA	1	0	0	0	0
4	BBB	1	0	0	0	0
5	AAA	266	0	0	2	0
5	BBB	282	0	0	1	0
All	All	9216	0	8623	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:BBB:802[A]:ADP:O2B	5:BBB:901:HOH:O	2.04	0.74
1:AAA:150[A]:SER:OG	5:AAA:701:HOH:O	2.08	0.70
1:BBB:154:LEU:HD13	1:BBB:201[B]:ILE:CD1	2.27	0.64
1:AAA:382:MET:HA	1:AAA:511:ALA:HB1	1.85	0.59
1:BBB:382:MET:HA	1:BBB:511:ALA:HB1	1.86	0.58
1:AAA:18:LYS:N	2:AAA:601[B]:ADP:O1B	2.37	0.56
1:BBB:154:LEU:HD13	1:BBB:201[B]:ILE:HD11	1.88	0.55
1:AAA:154:LEU:HD13	1:AAA:201[B]:ILE:CD1	2.40	0.52
1:AAA:154:LEU:HD13	1:AAA:201[B]:ILE:HD12	1.90	0.52
1:BBB:45:ILE:O	1:BBB:93:GLY:HA3	2.12	0.49
1:BBB:71:LEU:C	1:BBB:71:LEU:HD12	2.34	0.49
2:AAA:601[A]:ADP:O4'	1:BBB:182:ALA:HB1	2.12	0.48
1:BBB:177:VAL:HG12	1:BBB:186:VAL:HB	1.97	0.47
1:BBB:355:TYR:CD1	1:BBB:404:PHE:HB3	2.50	0.47
1:BBB:48:ASP:HB2	1:BBB:49:PRO:HD2	1.97	0.46
1:BBB:42:ASP:O	1:BBB:91:THR:HA	2.17	0.44
1:AAA:355:TYR:CD1	1:AAA:404:PHE:HB3	2.53	0.44
1:AAA:401:SER:HA	1:AAA:413:VAL:O	2.19	0.43
1:AAA:459:ASN:O	1:AAA:459:ASN:CG	2.56	0.43
1:AAA:182:ALA:HB1	2:BBB:802[B]:ADP:C4'	2.49	0.42
1:BBB:154:LEU:HD13	1:BBB:201[B]:ILE:HD12	2.01	0.42
1:AAA:294[B]:MET:HE2	1:AAA:294[B]:MET:HB2	1.88	0.42
1:AAA:177:VAL:HG12	1:AAA:186:VAL:HB	2.02	0.42
1:BBB:189:LYS:HB3	1:BBB:190:PRO:HD3	2.02	0.42
1:BBB:17:GLY:HA3	2:BBB:802[A]:ADP:C8	2.54	0.42
1:AAA:182:ALA:HB1	2:BBB:802[B]:ADP:H4'	2.02	0.41
1:AAA:189:LYS:HB3	1:AAA:190:PRO:HD3	2.03	0.41
1:BBB:116:ILE:HA	1:BBB:117:PRO:HA	1.87	0.41
1:BBB:1:MET:HE1	1:BBB:266:LEU:HD21	2.02	0.41
1:BBB:8:VAL:HA	1:BBB:172:MET:O	2.20	0.41
1:AAA:38:ILE:HA	1:AAA:138:LEU:O	2.21	0.40
1:BBB:172:MET:SD	1:BBB:209[B]:ILE:HD11	2.61	0.40
1:AAA:75:HIS:HE1	5:AAA:768:HOH:O	2.04	0.40
1:AAA:180[B]:MET:HG3	1:AAA:187:LYS:CG	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	545/545 (100%)	531 (97%)	14 (3%)	0	100	100
1	BBB	542/545 (99%)	530 (98%)	12 (2%)	0	100	100
All	All	1087/1090 (100%)	1061 (98%)	26 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	467/461 (101%)	467 (100%)	0	100	100
1	BBB	464/461 (101%)	460 (99%)	4 (1%)	78	79
All	All	931/922 (101%)	927 (100%)	4 (0%)	91	91

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	BBB	56	GLN
1	BBB	105	ARG
1	BBB	455	ARG
1	BBB	508	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADP	BBB	802[B]	-	24,29,29	0.65	0	29,45,45	0.83	1 (3%)
2	ADP	BBB	802[A]	-	24,29,29	0.66	0	29,45,45	0.76	1 (3%)
2	ADP	AAA	601[B]	-	24,29,29	0.64	0	29,45,45	0.73	1 (3%)
3	GTF	AAA	602	4	25,31,31	0.53	0	35,50,50	0.99	3 (8%)
3	GTF	AAA	603	4	25,31,31	0.62	0	35,50,50	1.07	2 (5%)
2	ADP	AAA	601[A]	-	24,29,29	0.68	0	29,45,45	0.78	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	BBB	802[B]	-	-	6/12/32/32	0/3/3/3
2	ADP	BBB	802[A]	-	-	0/12/32/32	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	AAA	601[B]	-	-	3/12/32/32	0/3/3/3
3	GTF	AAA	602	4	-	2/22/42/42	0/2/2/2
3	GTF	AAA	603	4	-	5/22/42/42	0/2/2/2
2	ADP	AAA	601[A]	-	-	4/12/32/32	0/3/3/3

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	AAA	603	GTF	PB-O3A-PA	3.33	144.25	132.83
3	AAA	602	GTF	PB-O3A-PA	3.04	143.25	132.83
3	AAA	603	GTF	C4'-O4'-C1'	-2.70	104.60	109.45
3	AAA	602	GTF	C4'-O4'-C1'	-2.38	105.17	109.45
2	AAA	601[A]	ADP	C5-C6-N6	2.33	123.89	120.35
2	BBB	802[B]	ADP	C5-C6-N6	2.27	123.80	120.35
2	BBB	802[A]	ADP	C5-C6-N6	2.24	123.76	120.35
2	AAA	601[B]	ADP	C5-C6-N6	2.22	123.72	120.35
3	AAA	602	GTF	F2-C2'-F1	2.09	107.59	105.20

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	601[A]	ADP	C5'-O5'-PA-O2A
2	AAA	601[A]	ADP	C5'-O5'-PA-O3A
2	BBB	802[B]	ADP	C5'-O5'-PA-O1A
2	BBB	802[B]	ADP	C5'-O5'-PA-O2A
2	BBB	802[B]	ADP	C5'-O5'-PA-O3A
2	AAA	601[B]	ADP	O4'-C4'-C5'-O5'
2	AAA	601[B]	ADP	C3'-C4'-C5'-O5'
2	BBB	802[B]	ADP	O4'-C4'-C5'-O5'
2	BBB	802[B]	ADP	C3'-C4'-C5'-O5'
3	AAA	603	GTF	PB-O3B-PG-O3G
3	AAA	603	GTF	PA-O3A-PB-O1B
3	AAA	602	GTF	PB-O3B-PG-O1G
2	AAA	601[A]	ADP	PB-O3A-PA-O2A
3	AAA	602	GTF	PB-O3A-PA-O2A
2	AAA	601[A]	ADP	C5'-O5'-PA-O1A
2	BBB	802[B]	ADP	PB-O3A-PA-O2A
3	AAA	603	GTF	PB-O3A-PA-O1A

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
3	AAA	603	GTF	PA-O3A-PB-O2B
2	AAA	601[B]	ADP	C5'-O5'-PA-O3A
3	AAA	603	GTF	PB-O3A-PA-O2A


There are no ring outliers.


4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	BBB	802[B]	ADP	2	0
2	BBB	802[A]	ADP	2	0
2	AAA	601[B]	ADP	1	0
2	AAA	601[A]	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	535/545 (98%)	0.72	44 (8%) 11 12	25, 50, 84, 102	0
1	BBB	535/545 (98%)	0.59	28 (5%) 27 29	26, 48, 79, 93	0
All	All	1070/1090 (98%)	0.66	72 (6%) 17 19	25, 49, 81, 102	0

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	481	ILE	7.3
1	AAA	419	TRP	5.1
1	AAA	395	ASN	4.8
1	BBB	335	THR	4.6
1	AAA	505	ASN	4.3
1	BBB	44	TYR	4.1
1	BBB	419	TRP	4.1
1	AAA	389	TYR	3.6
1	AAA	428	VAL	3.6
1	AAA	494	ASP	3.6
1	AAA	462	THR	3.5
1	AAA	44[A]	TYR	3.5
1	AAA	487	ARG	3.3
1	AAA	485	GLY	3.3
1	AAA	339	GLU	3.3
1	BBB	409	LYS	3.3
1	BBB	395	ASN	3.2
1	BBB	57	HIS	3.2
1	AAA	410	TYR	3.2
1	BBB	437	GLY	3.2
1	AAA	438	THR	3.2
1	BBB	428	VAL	3.2
1	AAA	495	ASP	3.1
1	AAA	456	GLN	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	BBB	487	ARG	3.1
1	BBB	495	ASP	3.1
1	AAA	393	VAL	3.0
1	BBB	416	ILE	3.0
1	AAA	405	VAL	3.0
1	BBB	410	TYR	3.0
1	BBB	101	ARG	2.9
1	BBB	494	ASP	2.9
1	AAA	486	LEU	2.9
1	AAA	498	VAL	2.9
1	AAA	367	PHE	2.9
1	AAA	397	GLU	2.8
1	AAA	491	ARG	2.8
1	AAA	504	PRO	2.8
1	BBB	423	ASN	2.8
1	BBB	504	PRO	2.8
1	BBB	407	ASP	2.8
1	AAA	416	ILE	2.7
1	AAA	437	GLY	2.7
1	AAA	394	ALA	2.7
1	BBB	397	GLU	2.6
1	AAA	101	ARG	2.6
1	BBB	481	ILE	2.6
1	BBB	393	VAL	2.4
1	BBB	366	ARG	2.4
1	AAA	496	GLN	2.4
1	AAA	372	ASN	2.4
1	BBB	1	MET	2.4
1	AAA	343	GLY	2.4
1	AAA	342	LYS	2.3
1	AAA	355	TYR	2.3
1	AAA	56	GLN	2.3
1	AAA	1	MET	2.2
1	AAA	245	TYR	2.2
1	BBB	342	LYS	2.2
1	BBB	398	ASN	2.2
1	AAA	422	GLU	2.2
1	AAA	451	ASP	2.2
1	AAA	425	ASN	2.2
1	AAA	484	ALA	2.2
1	BBB	438	THR	2.2
1	AAA	409	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	BBB	245	TYR	2.1
1	BBB	505	ASN	2.1
1	AAA	421	ASP	2.1
1	AAA	449	VAL	2.1
1	AAA	340	ILE	2.0
1	BBB	425	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ADP	BBB	802[A]	27/27	0.82	0.21	50,60,61,62	27
2	ADP	BBB	802[B]	27/27	0.82	0.21	49,55,58,59	27
2	ADP	AAA	601[A]	27/27	0.88	0.19	35,40,42,43	27
2	ADP	AAA	601[B]	27/27	0.88	0.19	48,51,57,57	27
4	NA	AAA	604	1/1	0.91	0.07	43,43,43,43	0
4	NA	BBB	801	1/1	0.95	0.07	37,37,37,37	0
3	GTF	AAA	602	30/30	0.98	0.16	25,27,30,32	0
3	GTF	AAA	603	30/30	0.98	0.15	27,29,31,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ADP BBB 802 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ADP BBB 802 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP AAA 601 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP AAA 601 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTF AAA 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GTF AAA 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.