



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 11, 2023 – 12:43 PM EST

PDB ID : 8G4P
Title : Crystal structure of the peanut allergen Ara h 2 bound by two neutralizing antibodies 13T1 and 13T5
Authors : Pedersen, L.C.; Mueller, G.A.; Min, J.
Deposited on : 2023-02-10
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	231	 7% 93% 5%
2	D	216	 4% 96%
3	A	232	 2% 92%
4	B	214	 1% 96%
5	E	135	 4% 69% 30%

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 14173 atoms, of which 6706 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 13t1 Fab heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	C	225	3091	1017	1482	261	324	7	0	0	0

- Molecule 2 is a protein called 13T1 Fab light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	D	212	3037	989	1464	264	316	4	0	0	0

- Molecule 3 is a protein called 13T5 Fab heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	A	222	3246	1055	1577	282	326	6	0	3	0

- Molecule 4 is a protein called 13T5 Fab light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
4	B	214	3188	1014	1561	273	335	5	0	0	0

- Molecule 5 is a protein called Ara h 2 allergen.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
5	E	94	1248	409	574	123	131	11	0	1	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	26	GLY	-	expression tag	UNP A0A445BYI5
E	27	SER	-	expression tag	UNP A0A445BYI5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	28	ALA	-	expression tag	UNP A0A445BYI5
E	29	ALA	-	expression tag	UNP A0A445BYI5
E	30	ALA	-	expression tag	UNP A0A445BYI5

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
6	C	1	10	2	6	2	0	0
6	A	1	10	2	6	2	0	0
6	A	1	10	2	6	2	0	0
6	B	1	10	2	6	2	0	0
6	B	1	10	2	6	2	0	0
6	B	1	10	2	6	2	0	0

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	1	Total	Cl	0	0
			1	1		
7	D	2	Total	Cl	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	3	Total	Cl	0	0
			3	3		
7	B	1	Total	Cl	0	0
			1	1		

- Molecule 8 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	C	1	Total	O	S	0	0
			5	4	1		
8	D	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		
8	E	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
9	A	1	26	8	12	1	5	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	C	24	Total	O	0	0
			24	24		
10	D	54	Total	O	0	0
			54	54		
10	A	55	Total	O	0	1
			56	56		
10	B	97	Total	O	0	2
			99	99		
10	E	17	Total	O	0	0
			17	17		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 13t1 Fab heavy chain




- Molecule 2: 13T1 Fab light chain




- Molecule 3: 13T5 Fab heavy chain



- Molecule 4: 13T5 Fab light chain



- Molecule 5: Ara h 2 allergen



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	136.30Å 136.79Å 96.82Å 90.00° 127.34° 90.00°	Depositor
Resolution (Å)	42.55 – 2.25 42.55 – 2.25	Depositor EDS
% Data completeness (in resolution range)	97.2 (42.55-2.25) 92.7 (42.55-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 2.24Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.189 , 0.217 0.186 , 0.213	Depositor DCC
R_{free} test set	2000 reflections (3.05%)	wwPDB-VP
Wilson B-factor (Å ²)	45.1	Xtrriage
Anisotropy	0.415	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14173	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NAG, EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.32	0/1650	0.60	0/2262
2	D	0.34	0/1611	0.60	0/2204
3	A	0.34	0/1712	0.65	0/2346
4	B	0.36	0/1659	0.65	0/2258
5	E	0.29	0/688	0.50	0/927
All	All	0.34	0/7320	0.62	0/9997

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1609	1482	1476	6	0
2	D	1573	1464	1462	3	0
3	A	1669	1577	1573	7	0
4	B	1627	1561	1553	3	0
5	E	674	574	559	1	0
6	A	8	12	12	0	0
6	B	12	18	18	0	0
6	C	4	6	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	3	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	2	0	0	0	0
8	B	5	0	0	0	0
8	C	5	0	0	0	0
8	D	5	0	0	0	0
8	E	5	0	0	0	0
9	A	14	12	13	1	0
10	A	56	0	0	1	0
10	B	99	0	0	0	0
10	C	24	0	0	1	0
10	D	54	0	0	1	0
10	E	17	0	0	0	0
All	All	7467	6706	6672	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:38:ARG:HB2	3:A:48:ILE:HD11	1.70	0.73
3:A:100[B]:ARG:NH2	5:E:140:CYS:O	2.27	0.68
3:A:155:LYS:NZ	3:A:183:GLN:OE1	2.30	0.61
3:A:38:ARG:NH2	3:A:89:ASP:OD1	2.35	0.60
2:D:189:GLU:O	2:D:213:ARG:NH2	2.36	0.59
1:C:50:ASP:OD2	2:D:98:HIS:NE2	2.38	0.57
3:A:18:LEU:CD1	3:A:121:VAL:HG11	2.34	0.57
3:A:18:LEU:HD12	3:A:121:VAL:HG11	1.92	0.51
1:C:216:THR:HG22	1:C:218:VAL:HG23	1.95	0.48
1:C:2:VAL:HG13	1:C:27:PHE:CD1	2.49	0.47
1:C:30:GLN:O	1:C:72:ARG:NH2	2.48	0.47
4:B:8:PRO:HG2	4:B:11:LEU:HD23	1.97	0.47
1:C:111:MET:HE2	10:C:514:HOH:O	2.15	0.46
3:A:100[B]:ARG:NH1	10:A:407:HOH:O	2.50	0.45
2:D:53:ARG:NH2	10:D:403:HOH:O	2.45	0.44
4:B:163:VAL:HG22	4:B:175:LEU:HD12	2.00	0.43
9:A:301:NAG:H82	4:B:95:PRO:HB2	2.01	0.43
1:C:12:VAL:HG21	1:C:18:LEU:HB2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	223/231 (96%)	219 (98%)	4 (2%)	0	100	100
2	D	210/216 (97%)	206 (98%)	4 (2%)	0	100	100
3	A	221/232 (95%)	217 (98%)	4 (2%)	0	100	100
4	B	212/214 (99%)	206 (97%)	5 (2%)	1 (0%)	29	29
5	E	91/135 (67%)	89 (98%)	2 (2%)	0	100	100
All	All	957/1028 (93%)	937 (98%)	19 (2%)	1 (0%)	51	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	B	31	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	164/192 (85%)	163 (99%)	1 (1%)	86	91
2	D	166/188 (88%)	166 (100%)	0	100	100
3	A	181/201 (90%)	181 (100%)	0	100	100
4	B	182/190 (96%)	180 (99%)	2 (1%)	73	82

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	E	62/121 (51%)	62 (100%)	0	100	100
All	All	755/892 (85%)	752 (100%)	3 (0%)	91	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	124	SER
4	B	30	SER
4	B	127	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 7 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EDO	B	604	-	3,3,3	0.53	0	2,2,2	0.15	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	C	401	-	3,3,3	0.53	0	2,2,2	0.09	0
8	SO4	B	605	-	4,4,4	0.14	0	6,6,6	0.23	0
8	SO4	E	201	-	4,4,4	0.15	0	6,6,6	0.09	0
6	EDO	B	602	-	3,3,3	0.50	0	2,2,2	0.33	0
8	SO4	C	403	-	4,4,4	0.26	0	6,6,6	0.28	0
6	EDO	A	303	-	3,3,3	0.49	0	2,2,2	0.27	0
8	SO4	D	303	-	4,4,4	0.14	0	6,6,6	0.22	0
6	EDO	B	601	-	3,3,3	0.56	0	2,2,2	0.24	0
9	NAG	A	301	3	14,14,15	0.25	0	17,19,21	0.68	1 (5%)
6	EDO	A	302	-	3,3,3	0.49	0	2,2,2	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	B	604	-	-	1/1/1/1	-
6	EDO	C	401	-	-	0/1/1/1	-
6	EDO	B	602	-	-	0/1/1/1	-
6	EDO	A	303	-	-	0/1/1/1	-
6	EDO	B	601	-	-	1/1/1/1	-
9	NAG	A	301	3	-	1/6/23/26	0/1/1/1
6	EDO	A	302	-	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	301	NAG	C1-O5-C5	2.21	115.19	112.19

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	601	EDO	O1-C1-C2-O2
6	B	604	EDO	O1-C1-C2-O2
9	A	301	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	301	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	225/231 (97%)	0.48	16 (7%) 16 17	41, 71, 109, 134	0
2	D	212/216 (98%)	0.12	8 (3%) 40 43	40, 58, 93, 116	0
3	A	222/232 (95%)	0.28	4 (1%) 68 71	36, 61, 93, 112	0
4	B	214/214 (100%)	0.04	2 (0%) 84 85	35, 50, 82, 103	0
5	E	94/135 (69%)	0.39	5 (5%) 26 29	43, 68, 109, 120	0
All	All	967/1028 (94%)	0.25	35 (3%) 42 44	35, 60, 97, 134	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	E	120	LEU	4.6
1	C	195	VAL	4.1
2	D	214	GLY	3.9
5	E	122	GLY	3.7
3	A	216	ASN	3.6
1	C	196	PRO	3.6
1	C	222	VAL	3.4
5	E	55	ARG	3.4
1	C	12	VAL	3.4
1	C	201	GLY	3.3
1	C	139	SER	3.2
1	C	142	THR	3.1
1	C	197	SER	3.0
1	C	15	GLY	3.0
3	A	63	LEU	2.9
2	D	25	ALA	2.8
1	C	144	GLY	2.7
2	D	5	THR	2.7
4	B	214	CYS	2.6
4	B	154	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	26	SER	2.5
5	E	56	ASP	2.4
2	D	71	PHE	2.4
1	C	84	SER	2.4
3	A	64	ASN	2.4
1	C	14	PRO	2.4
2	D	68	GLY	2.4
1	C	138	SER	2.3
1	C	146	THR	2.3
2	D	127	LEU	2.2
3	A	217	THR	2.2
2	D	4	MET	2.2
1	C	223	GLU	2.1
5	E	53	ILE	2.1
1	C	140	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
6	EDO	B	604	4/4	0.73	0.27	60,72,76,81	0
7	CL	A	305	1/1	0.76	0.31	84,84,84,84	0
6	EDO	A	302	4/4	0.77	0.21	76,91,102,102	0
8	SO4	C	403	5/5	0.78	0.16	66,73,92,95	0
7	CL	D	301	1/1	0.79	0.10	103,103,103,103	0
6	EDO	A	303	4/4	0.80	0.15	70,84,92,92	0
8	SO4	D	303	5/5	0.81	0.23	99,108,114,119	0
6	EDO	C	401	4/4	0.82	0.14	63,76,79,83	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	EDO	B	601	4/4	0.86	0.14	43,62,75,75	0
7	CL	A	306	1/1	0.86	0.07	80,80,80,80	0
6	EDO	B	602	4/4	0.86	0.24	62,75,79,87	0
7	CL	D	302	1/1	0.86	0.12	79,79,79,79	0
9	NAG	A	301	14/15	0.87	0.21	62,89,107,108	0
8	SO4	B	605	5/5	0.92	0.14	70,72,79,102	0
7	CL	A	304	1/1	0.92	0.19	78,78,78,78	0
8	SO4	E	201	5/5	0.93	0.22	79,80,96,105	0
7	CL	C	402	1/1	0.94	0.21	70,70,70,70	0
7	CL	B	603	1/1	0.96	0.12	75,75,75,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.