



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 14, 2022 – 03:57 AM EST

PDB ID : 1G5G
Title : FRAGMENT OF FUSION PROTEIN FROM NEWCASTLE DISEASE VIRUS
Authors : Lawrence, M.C.; Smith, B.J.
Deposited on : 2000-11-01
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	
1	B	481	
1	C	481	
1	D	481	
1	E	481	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	481	
2	G	2	
2	I	2	
2	J	2	
2	K	2	
2	L	2	
2	M	2	
2	N	2	
3	H	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	J	1	-	-	-	X
2	NAG	J	2	-	-	-	X
3	NAG	H	1	-	-	-	X
3	NAG	H	2	-	-	-	X
3	BMA	H	3	-	-	-	X
4	NAG	C	4471	-	-	X	X
4	NAG	D	1911	-	-	-	X
4	NAG	E	4471	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 16341 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fusion glycoprotein F0.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	357	2675	1679	440	538	18	0	0	0
1	B	357	2675	1679	440	538	18	0	0	0
1	C	357	2675	1679	440	538	18	0	0	0
1	D	357	2675	1679	440	538	18	0	0	0
1	E	357	2675	1679	440	538	18	0	0	0
1	F	357	2675	1679	440	538	18	0	0	0

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	500	SER	-	expression tag	UNP A9LSB1
A	501	ARG	-	expression tag	UNP A9LSB1
A	502	GLU	-	expression tag	UNP A9LSB1
A	503	GLN	-	expression tag	UNP A9LSB1
A	504	LYS	-	expression tag	UNP A9LSB1
A	505	LEU	-	expression tag	UNP A9LSB1
A	506	ILE	-	expression tag	UNP A9LSB1
A	507	SER	-	expression tag	UNP A9LSB1
A	508	GLU	-	expression tag	UNP A9LSB1
A	509	GLU	-	expression tag	UNP A9LSB1
A	510	ASP	-	expression tag	UNP A9LSB1
A	511	LEU	-	expression tag	UNP A9LSB1
A	512	ASN	-	expression tag	UNP A9LSB1
B	500	SER	-	expression tag	UNP A9LSB1
B	501	ARG	-	expression tag	UNP A9LSB1
B	502	GLU	-	expression tag	UNP A9LSB1
B	503	GLN	-	expression tag	UNP A9LSB1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	504	LYS	-	expression tag	UNP A9LSB1
B	505	LEU	-	expression tag	UNP A9LSB1
B	506	ILE	-	expression tag	UNP A9LSB1
B	507	SER	-	expression tag	UNP A9LSB1
B	508	GLU	-	expression tag	UNP A9LSB1
B	509	GLU	-	expression tag	UNP A9LSB1
B	510	ASP	-	expression tag	UNP A9LSB1
B	511	LEU	-	expression tag	UNP A9LSB1
B	512	ASN	-	expression tag	UNP A9LSB1
C	500	SER	-	expression tag	UNP A9LSB1
C	501	ARG	-	expression tag	UNP A9LSB1
C	502	GLU	-	expression tag	UNP A9LSB1
C	503	GLN	-	expression tag	UNP A9LSB1
C	504	LYS	-	expression tag	UNP A9LSB1
C	505	LEU	-	expression tag	UNP A9LSB1
C	506	ILE	-	expression tag	UNP A9LSB1
C	507	SER	-	expression tag	UNP A9LSB1
C	508	GLU	-	expression tag	UNP A9LSB1
C	509	GLU	-	expression tag	UNP A9LSB1
C	510	ASP	-	expression tag	UNP A9LSB1
C	511	LEU	-	expression tag	UNP A9LSB1
C	512	ASN	-	expression tag	UNP A9LSB1
D	500	SER	-	expression tag	UNP A9LSB1
D	501	ARG	-	expression tag	UNP A9LSB1
D	502	GLU	-	expression tag	UNP A9LSB1
D	503	GLN	-	expression tag	UNP A9LSB1
D	504	LYS	-	expression tag	UNP A9LSB1
D	505	LEU	-	expression tag	UNP A9LSB1
D	506	ILE	-	expression tag	UNP A9LSB1
D	507	SER	-	expression tag	UNP A9LSB1
D	508	GLU	-	expression tag	UNP A9LSB1
D	509	GLU	-	expression tag	UNP A9LSB1
D	510	ASP	-	expression tag	UNP A9LSB1
D	511	LEU	-	expression tag	UNP A9LSB1
D	512	ASN	-	expression tag	UNP A9LSB1
E	500	SER	-	expression tag	UNP A9LSB1
E	501	ARG	-	expression tag	UNP A9LSB1
E	502	GLU	-	expression tag	UNP A9LSB1
E	503	GLN	-	expression tag	UNP A9LSB1
E	504	LYS	-	expression tag	UNP A9LSB1
E	505	LEU	-	expression tag	UNP A9LSB1
E	506	ILE	-	expression tag	UNP A9LSB1

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
E	507	SER	-	expression tag	UNP A9LSB1
E	508	GLU	-	expression tag	UNP A9LSB1
E	509	GLU	-	expression tag	UNP A9LSB1
E	510	ASP	-	expression tag	UNP A9LSB1
E	511	LEU	-	expression tag	UNP A9LSB1
E	512	ASN	-	expression tag	UNP A9LSB1
F	500	SER	-	expression tag	UNP A9LSB1
F	501	ARG	-	expression tag	UNP A9LSB1
F	502	GLU	-	expression tag	UNP A9LSB1
F	503	GLN	-	expression tag	UNP A9LSB1
F	504	LYS	-	expression tag	UNP A9LSB1
F	505	LEU	-	expression tag	UNP A9LSB1
F	506	ILE	-	expression tag	UNP A9LSB1
F	507	SER	-	expression tag	UNP A9LSB1
F	508	GLU	-	expression tag	UNP A9LSB1
F	509	GLU	-	expression tag	UNP A9LSB1
F	510	ASP	-	expression tag	UNP A9LSB1
F	511	LEU	-	expression tag	UNP A9LSB1
F	512	ASN	-	expression tag	UNP A9LSB1

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	G	2	28	16	2	10	0	0	0
2	I	2	28	16	2	10	0	0	0
2	J	2	28	16	2	10	0	0	0
2	K	2	28	16	2	10	0	0	0
2	L	2	28	16	2	10	0	0	0
2	M	2	28	16	2	10	0	0	0
2	N	2	28	16	2	10	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	H	3	39	22	2	15	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	C	1	14	8	1	5	0	0
4	D	1	14	8	1	5	0	0
4	E	1	14	8	1	5	0	0
4	F	1	14	8	1	5	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fusion glycoprotein F0




- Molecule 1: Fusion glycoprotein F0







• Molecule 1: Fusion glycoprotein F0



• Molecule 1: Fusion glycoprotein F0



- Molecule 1: Fusion glycoprotein F0




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  50% 50%


MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  50% 50%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:  50% 50%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:  50% 50%

MAG1
MAG2

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  67% 33%

MAG1
MAG2
BMA3

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	134.39Å 308.33Å 243.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 3.30 24.84 – 3.29	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-3.30) 95.8 (24.84-3.29)	Depositor EDS
R_{merge}	0.22	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 3.30Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.224 , 0.273 0.225 , 0.266	Depositor DCC
R_{free} test set	3741 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	72.4	Xtrriage
Anisotropy	0.161	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 53.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	16341	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.87 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.3763e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2708	0.73	1/3683 (0.0%)
1	B	0.42	0/2708	0.73	0/3683
1	C	0.42	0/2708	0.74	1/3683 (0.0%)
1	D	0.44	0/2708	0.73	1/3683 (0.0%)
1	E	0.42	0/2708	0.74	0/3683
1	F	0.43	0/2708	0.73	0/3683
All	All	0.43	0/16248	0.73	3/22098 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	339	ILE	N-CA-C	-5.41	96.39	111.00
1	C	339	ILE	N-CA-C	-5.35	96.56	111.00
1	D	339	ILE	N-CA-C	-5.10	97.22	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2675	0	2716	311	0
1	B	2675	0	2717	313	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2675	0	2716	329	0
1	D	2675	0	2717	297	0
1	E	2675	0	2716	310	0
1	F	2675	0	2717	318	0
2	G	28	0	25	6	0
2	I	28	0	25	2	0
2	J	28	0	25	3	0
2	K	28	0	25	1	0
2	L	28	0	25	1	0
2	M	28	0	25	1	0
2	N	28	0	25	2	0
3	H	39	0	34	5	0
4	C	14	0	13	7	0
4	D	14	0	13	0	0
4	E	14	0	13	5	0
4	F	14	0	13	0	0
All	All	16341	0	16560	1521	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 46.

The worst 5 of 1521 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:VAL:HG13	1:A:292:VAL:HG11	1.23	1.19
1:D:255:VAL:HG13	1:D:292:VAL:HG11	1.27	1.10
1:E:314:PHE:HB3	1:E:374:LYS:HD3	1.32	1.10
1:C:405:PRO:CG	1:F:64:LYS:HD2	1.81	1.10
1:A:314:PHE:HB3	1:A:374:LYS:HD3	1.31	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	353/481 (73%)	304 (86%)	40 (11%)	9 (2%)	5	27
1	B	353/481 (73%)	299 (85%)	40 (11%)	14 (4%)	3	18
1	C	353/481 (73%)	290 (82%)	56 (16%)	7 (2%)	7	32
1	D	353/481 (73%)	299 (85%)	48 (14%)	6 (2%)	9	35
1	E	353/481 (73%)	302 (86%)	40 (11%)	11 (3%)	4	23
1	F	353/481 (73%)	289 (82%)	56 (16%)	8 (2%)	6	29
All	All	2118/2886 (73%)	1783 (84%)	280 (13%)	55 (3%)	5	27

5 of 55 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	72	ASP
1	B	403	ASP
1	B	412	TYR
1	C	403	ASP
1	D	310	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/406 (76%)	254 (83%)	53 (17%)	2	8
1	B	307/406 (76%)	256 (83%)	51 (17%)	2	10
1	C	307/406 (76%)	250 (81%)	57 (19%)	1	7
1	D	307/406 (76%)	249 (81%)	58 (19%)	1	6
1	E	307/406 (76%)	260 (85%)	47 (15%)	2	12
1	F	307/406 (76%)	251 (82%)	56 (18%)	1	7
All	All	1842/2436 (76%)	1520 (82%)	322 (18%)	2	8

5 of 322 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	192	LYS
1	F	231	THR
1	E	231	THR
1	E	430	ASP
1	F	355	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 67 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	184	GLN
1	F	205	GLN
1	F	422	GLN
1	C	57	GLN
1	B	451	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

17 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	G	1	1,2	14,14,15	0.56	0	17,19,21	0.77	1 (5%)
2	NAG	G	2	2	14,14,15	0.48	0	17,19,21	0.66	0
3	NAG	H	1	1,3	14,14,15	0.68	0	17,19,21	0.66	0
3	NAG	H	2	3	14,14,15	1.06	1 (7%)	17,19,21	1.28	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BMA	H	3	3	11,11,12	0.59	0	15,15,17	0.61	0
2	NAG	I	1	1,2	14,14,15	0.50	0	17,19,21	0.76	1 (5%)
2	NAG	I	2	2	14,14,15	0.48	0	17,19,21	0.89	1 (5%)
2	NAG	J	1	1,2	14,14,15	0.83	0	17,19,21	1.22	3 (17%)
2	NAG	J	2	2	14,14,15	0.80	0	17,19,21	0.71	0
2	NAG	K	1	1,2	14,14,15	0.54	0	17,19,21	0.67	0
2	NAG	K	2	2	14,14,15	0.46	0	17,19,21	0.83	1 (5%)
2	NAG	L	1	1,2	14,14,15	0.58	0	17,19,21	0.68	1 (5%)
2	NAG	L	2	2	14,14,15	0.66	0	17,19,21	0.68	0
2	NAG	M	1	1,2	14,14,15	0.66	0	17,19,21	0.63	0
2	NAG	M	2	2	14,14,15	0.56	0	17,19,21	0.74	1 (5%)
2	NAG	N	1	1,2	14,14,15	0.56	0	17,19,21	0.71	0
2	NAG	N	2	2	14,14,15	0.52	0	17,19,21	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	G	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	G	2	2	-	3/6/23/26	0/1/1/1
3	NAG	H	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	H	2	3	-	0/6/23/26	0/1/1/1
3	BMA	H	3	3	-	2/2/19/22	0/1/1/1
2	NAG	I	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	I	2	2	-	0/6/23/26	0/1/1/1
2	NAG	J	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	J	2	2	-	0/6/23/26	0/1/1/1
2	NAG	K	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	K	2	2	-	0/6/23/26	0/1/1/1
2	NAG	L	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	L	2	2	-	2/6/23/26	0/1/1/1
2	NAG	M	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	M	2	2	-	0/6/23/26	0/1/1/1
2	NAG	N	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	N	2	2	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	2	NAG	C1-C2	2.98	1.56	1.52

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	1	NAG	C1-O5-C5	2.72	115.88	112.19
3	H	2	NAG	O5-C1-C2	2.72	115.58	111.29
2	K	2	NAG	C2-N2-C7	-2.50	119.35	122.90
2	I	1	NAG	C2-N2-C7	-2.35	119.55	122.90
2	J	1	NAG	C4-C3-C2	-2.33	107.60	111.02

There are no chirality outliers.

5 of 19 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	G	1	NAG	O5-C5-C6-O6
3	H	3	BMA	O5-C5-C6-O6
3	H	3	BMA	C4-C5-C6-O6
3	H	1	NAG	O5-C5-C6-O6
3	H	1	NAG	C4-C5-C6-O6


There are no ring outliers.


13 monomers are involved in 21 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	L	2	NAG	1	0
3	H	1	NAG	2	0
3	H	3	BMA	3	0
2	J	1	NAG	3	0
2	N	1	NAG	2	0
3	H	2	NAG	2	0
2	M	2	NAG	1	0
2	I	2	NAG	1	0
2	G	2	NAG	5	0
2	J	2	NAG	3	0
2	I	1	NAG	1	0
2	G	1	NAG	2	0
2	K	2	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




















5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	F	1911	1	14,14,15	0.76	0	17,19,21	0.57	0
4	NAG	C	4471	1	14,14,15	0.53	0	17,19,21	0.65	1 (5%)
4	NAG	D	1911	1	14,14,15	0.55	0	17,19,21	0.66	0
4	NAG	E	4471	1	14,14,15	0.76	1 (7%)	17,19,21	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	F	1911	1	-	2/6/23/26	0/1/1/1
4	NAG	C	4471	1	-	2/6/23/26	0/1/1/1
4	NAG	D	1911	1	-	2/6/23/26	0/1/1/1
4	NAG	E	4471	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	4471	NAG	C1-C2	2.02	1.55	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	4471	NAG	C2-N2-C7	-2.03	120.02	122.90

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	4471	NAG	O5-C5-C6-O6
4	F	1911	NAG	O5-C5-C6-O6
4	D	1911	NAG	O5-C5-C6-O6
4	E	4471	NAG	C4-C5-C6-O6
4	C	4471	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	4471	NAG	7	0
4	E	4471	NAG	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	357/481 (74%)	-0.24	12 (3%) 45 43	7, 34, 87, 100	0
1	B	357/481 (74%)	-0.20	10 (2%) 53 51	3, 35, 88, 100	0
1	C	357/481 (74%)	-0.21	14 (3%) 39 37	7, 33, 90, 100	0
1	D	357/481 (74%)	-0.25	13 (3%) 42 40	6, 33, 88, 100	0
1	E	357/481 (74%)	-0.19	11 (3%) 49 48	7, 34, 86, 100	0
1	F	357/481 (74%)	-0.25	6 (1%) 70 68	5, 33, 87, 99	0
All	All	2142/2886 (74%)	-0.22	66 (3%) 49 48	3, 34, 88, 100	0

The worst 5 of 66 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	173	SER	11.0
1	B	173	SER	4.8
1	F	174	GLN	4.1
1	E	172	LEU	4.1
1	C	173	SER	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	H	1	14/15	0.41	0.77	90,95,98,98	0
2	NAG	J	2	14/15	0.47	0.91	93,100,100,100	0
3	NAG	H	2	14/15	0.52	0.99	97,100,100,100	0
3	BMA	H	3	11/12	0.72	0.81	98,100,100,100	0
2	NAG	J	1	14/15	0.76	0.65	84,90,95,96	0
2	NAG	L	2	14/15	0.85	0.29	66,72,73,75	0
2	NAG	N	2	14/15	0.87	0.32	48,49,54,57	0
2	NAG	I	2	14/15	0.87	0.32	55,60,63,68	0
2	NAG	G	2	14/15	0.89	0.43	61,66,75,77	0
2	NAG	M	2	14/15	0.89	0.29	66,70,75,79	0
2	NAG	L	1	14/15	0.89	0.24	48,59,71,73	0
2	NAG	M	1	14/15	0.90	0.16	49,55,64,64	0
2	NAG	G	1	14/15	0.90	0.39	49,55,64,65	0
2	NAG	K	2	14/15	0.90	0.43	40,43,48,53	0
2	NAG	K	1	14/15	0.92	0.26	34,40,49,52	0
2	NAG	I	1	14/15	0.94	0.13	46,53,56,59	0
2	NAG	N	1	14/15	0.94	0.18	38,45,53,56	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain J:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain L:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain M:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain N:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain H:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	D	1911	14/15	0.70	0.67	99,100,100,100	0
4	NAG	E	4471	14/15	0.74	0.55	84,90,100,100	0
4	NAG	C	4471	14/15	0.79	0.48	79,86,91,92	0
4	NAG	F	1911	14/15	0.81	0.46	91,95,100,100	0

6.5 Other polymers [i](#)

There are no such residues in this entry.