



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 08:47 pm GMT

PDB ID : 3G6V
Title : DNA synthesis across an abasic lesion by human DNA polymerase-iota
Authors : Nair, D.T.; Aggarwal, A.K.
Deposited on : 2009-02-09
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtrriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4343 (2.20-2.20)
Clashscore	122126	5027 (2.20-2.20)
Ramachandran outliers	120053	4952 (2.20-2.20)
Sidechain outliers	120020	4953 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	420	65% (green), 20% (yellow), 11% (grey), 4% (orange)
2	P	7	71% (yellow), 29% (orange)
3	T	12	8% (green), 50% (yellow), 17% (orange), 25% (grey)

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3390 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase iota.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	373	2876	1809	504	542	21	0	0	0


- Molecule 2 is a DNA chain called Primer DNA strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	P	7	139	67	29	37	6	0	0	0

- Molecule 3 is a DNA chain called Template DNA strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	T	9	172	83	27	54	8	0	0	0

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	30	10	5	12	3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	2	2	2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	156	156	156	0	0
6	P	3	3	3	0	0
6	T	12	12	12	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase iota



- Molecule 2: Primer DNA strand



- Molecule 3: Template DNA strand



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	98.15Å 98.15Å 203.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.22 – 2.20 39.21 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.0 (39.22-2.20) 97.0 (39.21-2.20)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.46 (at 2.20Å)	Xtrriage
Refinement program	CNS 1.1, REFMAC 5.2.0019	Depositor
R, R_{free}	0.211 , 0.251 0.273 , 0.308	Depositor DCC
R_{free} test set	2417 reflections (8.05%)	wwPDB-VP
Wilson B-factor (Å ²)	34.3	Xtrriage
Anisotropy	0.206	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3390	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: DOC, MG, ATP, 3DR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.10	5/2914 (0.2%)	1.18	15/3934 (0.4%)
2	P	1.81	0/136	3.27	20/208 (9.6%)
3	T	1.92	2/178 (1.1%)	2.92	24/271 (8.9%)
All	All	1.20	7/3228 (0.2%)	1.51	59/4413 (1.3%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	842	DG	C3'-O3'	-6.86	1.35	1.44
1	A	212	VAL	CB-CG1	6.83	1.67	1.52
3	T	839	DT	C1'-N1	6.72	1.57	1.49
1	A	122	ARG	CD-NE	-6.54	1.35	1.46
1	A	207	LYS	CE-NZ	-6.45	1.32	1.49
1	A	185	GLU	CB-CG	-5.73	1.41	1.52
1	A	122	ARG	CG-CD	5.04	1.64	1.51

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	122	ARG	NE-CZ-NH2	-27.27	106.66	120.30
1	A	184	ARG	NE-CZ-NH2	-15.63	112.48	120.30
2	P	870	DA	O4'-C1'-N9	15.62	118.94	108.00
1	A	122	ARG	NE-CZ-NH1	15.31	127.95	120.30
2	P	867	DA	O4'-C1'-N9	13.79	117.65	108.00
1	A	184	ARG	NE-CZ-NH1	13.74	127.17	120.30
2	P	867	DA	O4'-C4'-C3'	-11.62	99.03	106.00
2	P	870	DA	O5'-P-OP2	-11.56	95.30	105.70
3	T	841	DG	C2-N3-C4	11.46	117.63	111.90
2	P	869	DG	O4'-C1'-N9	11.15	115.80	108.00
3	T	841	DG	C5-C6-N1	10.96	116.98	111.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	871	DC	O5'-P-OP2	-10.54	96.21	105.70
1	A	126	ASP	CB-CG-OD1	-10.16	109.16	118.30
1	A	122	ARG	CD-NE-CZ	9.80	137.31	123.60
3	T	841	DG	O4'-C1'-N9	9.23	114.46	108.00
3	T	845	DC	P-O5'-C5'	8.87	135.10	120.90
2	P	870	DA	O4'-C4'-C3'	8.73	111.24	106.00
3	T	839	DT	O4'-C1'-N1	8.57	114.00	108.00
3	T	845	DC	P-O3'-C3'	8.49	129.89	119.70
3	T	841	DG	C5-C6-O6	-8.45	123.53	128.60
3	T	842	DG	O4'-C1'-C2'	8.08	112.37	105.90
2	P	868	DG	N1-C6-O6	7.83	124.60	119.90
2	P	869	DG	N9-C4-C5	7.46	108.39	105.40
3	T	846	DC	N1-C2-O2	7.36	123.31	118.90
3	T	842	DG	N1-C6-O6	-7.16	115.61	119.90
2	P	869	DG	C8-N9-C4	-7.06	103.58	106.40
1	A	122	ARG	CG-CD-NE	-7.05	96.98	111.80
3	T	842	DG	O4'-C1'-N9	-6.96	103.13	108.00
2	P	872	DC	O4'-C4'-C3'	-6.83	101.77	104.50
3	T	847	DT	C6-C5-C7	-6.83	118.80	122.90
3	T	847	DT	C4-C5-C7	6.64	122.98	119.00
3	T	843	DG	O4'-C4'-C3'	-6.43	101.93	104.50
2	P	871	DC	O4'-C4'-C3'	-6.32	101.97	104.50
3	T	841	DG	C6-N1-C2	-6.26	121.34	125.10
2	P	872	DC	C6-N1-C2	6.25	122.80	120.30
3	T	843	DG	O5'-P-OP2	-6.25	100.08	105.70
3	T	842	DG	N1-C2-N2	-6.20	110.62	116.20
2	P	869	DG	P-O3'-C3'	-6.15	112.32	119.70
1	A	331	ARG	NE-CZ-NH1	6.06	123.33	120.30
3	T	844	DT	C6-C5-C7	-5.99	119.30	122.90
2	P	871	DC	C4'-C3'-C2'	-5.99	97.71	103.10
3	T	847	DT	O4'-C1'-N1	-5.94	103.84	108.00
1	A	205	LEU	CB-CG-CD1	5.72	120.72	111.00
1	A	28	ARG	NE-CZ-NH2	-5.62	117.49	120.30
3	T	846	DC	N3-C4-N4	-5.50	114.15	118.00
2	P	868	DG	C5-C6-O6	-5.44	125.34	128.60
2	P	869	DG	C4-C5-N7	-5.38	108.65	110.80
2	P	871	DC	N1-C2-O2	-5.37	115.68	118.90
2	P	870	DA	C3'-C2'-C1'	5.32	108.89	102.50
1	A	337	ARG	NE-CZ-NH1	5.23	122.92	120.30
3	T	846	DC	C5-C4-N4	5.21	123.84	120.20
3	T	839	DT	C6-N1-C2	-5.18	118.71	121.30
3	T	842	DG	C5-C6-O6	5.13	131.68	128.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	869	DG	N1-C6-O6	-5.13	116.83	119.90
1	A	135	MET	CG-SD-CE	-5.12	92.01	100.20
3	T	844	DT	N1-C2-N3	5.11	117.67	114.60
1	A	347	ARG	NE-CZ-NH2	-5.08	117.76	120.30
1	A	103	ARG	NE-CZ-NH1	-5.03	117.79	120.30
1	A	140	LEU	CB-CG-CD1	-5.03	102.46	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2876	0	2899	65	0
2	P	139	0	79	1	1
3	T	172	0	102	1	1
4	A	30	0	10	0	0
5	A	2	0	0	0	0
6	A	156	0	0	3	0
6	P	3	0	0	0	0
6	T	12	0	0	0	0
All	All	3390	0	3090	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:LEU:CD1	1:A:387:LEU:HD13	1.81	1.10
1:A:309:LYS:H	1:A:309:LYS:CD	1.64	1.08
1:A:344:LEU:HD11	1:A:387:LEU:CD1	1.86	1.05
1:A:309:LYS:HD2	1:A:309:LYS:H	1.16	1.05
1:A:344:LEU:HD11	1:A:387:LEU:HD13	0.97	0.93
1:A:44:MET:CE	1:A:67:ASN:HD22	1.80	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:ARG:HD2	6:A:1014:HOH:O	1.70	0.92
1:A:164:ASN:H	1:A:170:HIS:HD2	1.20	0.88
1:A:344:LEU:HD13	1:A:390:LEU:HD12	1.60	0.84
1:A:309:LYS:N	1:A:309:LYS:CD	2.45	0.79
1:A:249:CYS:SG	1:A:273:LEU:HD21	2.27	0.73
1:A:44:MET:HE3	1:A:67:ASN:HD22	1.53	0.73
1:A:335:ASP:OD2	1:A:337:ARG:HD3	1.88	0.72
1:A:347:ARG:HD2	1:A:404:THR:OG1	1.89	0.71
1:A:80:ASN:HB3	6:A:1018:HOH:O	1.94	0.68
1:A:164:ASN:H	1:A:170:HIS:CD2	2.09	0.66
1:A:44:MET:CE	1:A:67:ASN:ND2	2.58	0.66
1:A:44:MET:HE1	1:A:67:ASN:HD22	1.60	0.65
1:A:48:PRO:O	1:A:51:LYS:HB2	1.98	0.64
1:A:343:ARG:HD2	1:A:345:ILE:HD11	1.80	0.64
1:A:335:ASP:CG	1:A:337:ARG:HD3	2.21	0.61
1:A:202:ASN:ND2	1:A:205:LEU:H	1.99	0.60
1:A:321:ILE:HD11	1:A:406:LEU:HD22	1.84	0.59
1:A:309:LYS:HD2	1:A:309:LYS:N	2.01	0.59
1:A:44:MET:HE2	1:A:51:LYS:O	2.03	0.59
1:A:360:ARG:HG2	1:A:394:MET:HG3	1.86	0.57
1:A:335:ASP:OD1	1:A:337:ARG:HD3	2.05	0.56
1:A:333:CYS:O	1:A:335:ASP:N	2.40	0.55
1:A:44:MET:HE3	1:A:67:ASN:ND2	2.19	0.55
1:A:365:PRO:HB2	1:A:368:VAL:HG13	1.89	0.53
1:A:309:LYS:HD3	1:A:311:CYS:SG	2.50	0.52
1:A:318:LYS:HB2	1:A:388:MET:SD	2.50	0.51
1:A:44:MET:HE1	1:A:67:ASN:ND2	2.23	0.51
1:A:309:LYS:H	1:A:309:LYS:HD3	1.67	0.51
1:A:184:ARG:HD2	1:A:195:GLY:O	2.12	0.49
1:A:112:LEU:HD23	1:A:112:LEU:C	2.35	0.47
1:A:122:ARG:HD3	6:A:957:HOH:O	2.14	0.47
3:T:846:DC:H1'	3:T:847:DT:H5'	1.96	0.47
1:A:308:PHE:HD1	1:A:311:CYS:H	1.62	0.47
1:A:202:ASN:C	1:A:202:ASN:HD22	2.18	0.46
1:A:329:LEU:HD13	1:A:380:MET:HG3	1.97	0.46
1:A:253:LEU:HD11	1:A:272:GLU:HG2	1.98	0.46
1:A:36:ASP:O	1:A:37:CYS:C	2.55	0.46
1:A:168:VAL:HG22	1:A:172:ARG:HD2	1.97	0.45
1:A:139:ARG:CZ	1:A:179:ILE:CD1	2.94	0.45
1:A:388:MET:O	1:A:391:PHE:HB3	2.18	0.44
1:A:196:CYS:HA	1:A:217:GLN:O	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:LEU:HD22	1:A:229:LEU:HD12	1.99	0.44
1:A:44:MET:HE2	1:A:51:LYS:HA	2.00	0.43
1:A:105:MET:CG	1:A:193:LEU:HD11	2.48	0.43
1:A:59:GLN:OE1	1:A:64:VAL:HG11	2.19	0.43
1:A:251:GLU:O	1:A:252:ALA:C	2.57	0.42
1:A:325:LEU:O	1:A:329:LEU:HD12	2.19	0.42
1:A:164:ASN:N	1:A:170:HIS:HD2	2.02	0.42
1:A:345:ILE:HG13	1:A:359:SER:HB3	2.02	0.42
1:A:309:LYS:N	1:A:309:LYS:HD3	2.31	0.42
1:A:364:ILE:HG23	1:A:368:VAL:HG22	2.01	0.42
1:A:344:LEU:O	1:A:359:SER:HA	2.20	0.42
1:A:347:ARG:HD2	1:A:404:THR:HG1	1.83	0.41
1:A:147:GLU:HG2	1:A:147:GLU:H	1.56	0.41
1:A:250:LEU:HB3	1:A:255:ILE:HB	2.01	0.41
2:P:872:DC:H2'	2:P:873:DOC:H6	2.03	0.41
1:A:359:SER:O	1:A:360:ARG:HD3	2.21	0.40
1:A:76:LYS:HA	1:A:76:LYS:HD3	1.84	0.40
1:A:150:ALA:O	1:A:152:THR:HG23	2.20	0.40
1:A:202:ASN:HD21	1:A:205:LEU:H	1.68	0.40
1:A:322:GLU:HG3	1:A:322:GLU:H	1.73	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:P:867:DA:O5'	3:T:847:DT:O3'[10_665]	1.84	0.36

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	367/420 (87%)	341 (93%)	18 (5%)	8 (2%)	7 4

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	310	LYS
1	A	315	VAL
1	A	333	CYS
1	A	334	GLN
1	A	354	HIS
1	A	313	SER
1	A	316	GLU
1	A	356	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	321/376 (85%)	291 (91%)	30 (9%)	10 9

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	LEU
1	A	48	PRO
1	A	51	LYS
1	A	52	ASP
1	A	62	LEU
1	A	87	LYS
1	A	90	GLN
1	A	123	LEU
1	A	126	ASP
1	A	142	GLN
1	A	145	SER
1	A	149	SER
1	A	170	HIS
1	A	202	ASN
1	A	204	LEU
1	A	205	LEU
1	A	209	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	229	LEU
1	A	236	ILE
1	A	267	LYS
1	A	295	LEU
1	A	307	SER
1	A	309	LYS
1	A	329	LEU
1	A	348	ARG
1	A	357	ARG
1	A	360	ARG
1	A	385	ASP
1	A	388	MET
1	A	413	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	GLN
1	A	67	ASN
1	A	170	HIS
1	A	202	ASN
1	A	262	GLN
1	A	300	GLN
1	A	412	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DOC	P	873	3,2	13,19,20	1.12	0	12,26,29	2.66	5 (41%)
3	3DR	T	840	3	8,11,12	1.51	1 (12%)	9,14,17	1.64	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DOC	P	873	3,2	-	0/3/18/19	0/2/2/2
3	3DR	T	840	3	-	0/3/15/16	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	840	3DR	O4'-C4'	-3.29	1.38	1.44

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	873	DOC	C2'-C1'-N1	-6.15	100.91	112.48
3	T	840	3DR	O3'-C3'-C2'	-2.73	105.17	111.62
3	T	840	3DR	C1'-C2'-C3'	-2.42	100.29	103.19
3	T	840	3DR	O4'-C1'-C2'	-2.33	101.66	106.41
2	P	873	DOC	O4'-C4'-C5'	2.05	112.92	109.59
2	P	873	DOC	O4'-C4'-C3'	2.96	109.73	104.80
2	P	873	DOC	O4'-C1'-N1	3.13	113.06	107.78
2	P	873	DOC	C3'-C2'-C1'	4.67	108.40	102.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	P	873	DOC	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ATP	A	875	5	27,32,33	1.87	9 (33%)	29,50,52	1.92	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	875	5	-	0/18/34/38	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	875	ATP	C5'-C4'	2.19	1.58	1.51
4	A	875	ATP	C2'-C1'	2.28	1.58	1.52
4	A	875	ATP	PG-O3B	2.32	1.63	1.60
4	A	875	ATP	C8-N9	2.38	1.39	1.36
4	A	875	ATP	C8-N7	2.47	1.39	1.34
4	A	875	ATP	C2-N3	2.53	1.36	1.32
4	A	875	ATP	C4-N3	3.30	1.40	1.35
4	A	875	ATP	O3'-C3'	3.57	1.51	1.43
4	A	875	ATP	C5-C4	4.25	1.50	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	875	ATP	N3-C2-N1	-6.32	123.46	128.86
4	A	875	ATP	C5-C6-N6	-2.36	115.67	120.47
4	A	875	ATP	O3'-C3'-C2'	2.13	118.57	110.86
4	A	875	ATP	C4-C5-N7	2.58	111.90	109.41
4	A	875	ATP	C2-N1-C6	2.89	123.67	118.75
4	A	875	ATP	N6-C6-N1	3.76	126.37	118.57

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.