



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 20, 2023 – 12:05 PM EDT

PDB ID : 8G65  
Title : Wildtype PTP1b in complex with DES4799  
Authors : Greisman, J.B.; Willmore, L.; Yeh, C.Y.; Giordanetto, F.; Shahamadtar, S.;  
Nisonoff, H.; Maragakis, P.; Shaw, D.E.  
Deposited on : 2023-02-14  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.32.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.32.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	
1	B	298	

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 5587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	Total 2500	C 1589	N 421	O 470	S 20	0	14	0
1	B	284	Total 2411	C 1538	N 406	O 447	S 20	0	19	0

- Molecule 2 is 4-(3,5-dimethyl-1H-pyrazol-1-yl)aniline (three-letter code: YW9) (formula: C<sub>11</sub>H<sub>13</sub>N<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
2	A	1	Total 14	C 11	N 3	0	0
2	B	1	Total 14	C 11	N 3	0	0


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Cl 1 1	0	0

- Molecule 5 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O S 4 2 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	323	Total O 323 323	0	0
6	B	319	Total O 319 319	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 1



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.44Å 89.44Å 165.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.32 – 1.45 63.24 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.8 (63.32-1.45) 99.8 (63.24-1.45)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.53 (at 1.45Å)	Xtrriage
Refinement program	REFMAC 5.8.0131	Depositor
R, $R_{free}$	0.134 , 0.188 0.142 , 0.195	Depositor DCC
$R_{free}$ test set	6003 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.2	Xtrriage
Anisotropy	1.081	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 40.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	5587	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.87% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, MG, CL, YW9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.46	10/2600 (0.4%)	1.31	22/3500 (0.6%)
1	B	1.54	18/2521 (0.7%)	1.40	27/3393 (0.8%)
All	All	1.50	28/5121 (0.5%)	1.35	49/6893 (0.7%)

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	2	GLU	CD-OE1	12.43	1.39	1.25
1	A	121	CYS	CB-SG	-10.93	1.63	1.82
1	A	297	GLU	CD-OE1	8.80	1.35	1.25
1	A	147	GLU	CD-OE1	-8.55	1.16	1.25
1	A	291	TRP	CB-CG	7.92	1.64	1.50
1	B	2	GLU	CD-OE2	7.12	1.33	1.25
1	B	132	GLU	CD-OE1	7.12	1.33	1.25
1	A	298	ASP	C-OXT	6.78	1.36	1.23
1	A	153	TYR	CE1-CZ	6.72	1.47	1.38
1	B	147	GLU	CD-OE2	-6.72	1.18	1.25
1	B	147	GLU	CB-CG	-6.47	1.39	1.52
1	A	153	TYR	CG-CD2	6.33	1.47	1.39
1	B	6	GLU	CD-OE2	6.08	1.32	1.25
1	A	289	ASP	CG-OD2	-5.93	1.11	1.25
1	B	170	GLU	CD-OE2	5.91	1.32	1.25
1	B	20	TYR	CG-CD1	5.67	1.46	1.39
1	B	193	ASN	CG-OD1	5.65	1.36	1.24
1	B	2	GLU	CG-CD	5.59	1.60	1.51
1	B	196	PHE	CG-CD2	-5.45	1.30	1.38
1	B	284	ASP	CB-CG	5.44	1.63	1.51
1	B	29	ASP	CB-CG	5.44	1.63	1.51
1	B	252	GLU	CD-OE2	5.36	1.31	1.25
1	A	9	GLN	CD-NE2	5.35	1.46	1.32
1	B	129	GLU	CG-CD	5.33	1.59	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	26	GLU	CD-OE1	5.26	1.31	1.25
1	A	9	GLN	CG-CD	5.14	1.62	1.51
1	B	130	GLU	CD-OE2	5.11	1.31	1.25
1	B	218	GLY	C-O	5.06	1.31	1.23

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	79	ARG	NE-CZ-NH1	15.79	128.19	120.30
1	A	43	ARG	NE-CZ-NH1	12.66	126.63	120.30
1	B	199	ARG	NE-CZ-NH1	12.45	126.52	120.30
1	B	257	ARG	NE-CZ-NH1	10.09	125.34	120.30
1	A	169	ARG	NE-CZ-NH2	-10.07	115.26	120.30
1	B	112[A]	ARG	NE-CZ-NH1	-10.03	115.29	120.30
1	B	112[B]	ARG	NE-CZ-NH1	-10.03	115.29	120.30
1	B	79	ARG	NE-CZ-NH2	-9.86	115.37	120.30
1	A	268	ARG	NE-CZ-NH1	9.83	125.22	120.30
1	B	235[A]	MET	CG-SD-CE	-9.78	84.56	100.20
1	B	235[B]	MET	CG-SD-CE	-9.78	84.56	100.20
1	A	79	ARG	NE-CZ-NH2	-9.20	115.70	120.30
1	B	236	ASP	CB-CG-OD2	-8.66	110.51	118.30
1	B	112[A]	ARG	NE-CZ-NH2	8.04	124.32	120.30
1	B	112[B]	ARG	NE-CZ-NH2	8.04	124.32	120.30
1	A	22	ASP	CB-CG-OD2	-7.88	111.20	118.30
1	A	268	ARG	NE-CZ-NH2	-7.84	116.38	120.30
1	A	253	MET	CG-SD-CE	-7.62	88.00	100.20
1	B	30	PHE	CB-CG-CD2	7.45	126.02	120.80
1	B	245	ASP	CB-CG-OD1	7.01	124.61	118.30
1	B	199	ARG	NE-CZ-NH2	-6.96	116.82	120.30
1	B	1	MET	CG-SD-CE	6.75	111.00	100.20
1	A	121	CYS	N-CA-CB	-6.72	98.50	110.60
1	A	112	ARG	NE-CZ-NH2	6.71	123.65	120.30
1	A	22	ASP	CB-CG-OD1	6.50	124.15	118.30
1	A	79	ARG	NE-CZ-NH1	6.36	123.48	120.30
1	A	33	ARG	NE-CZ-NH2	6.31	123.46	120.30
1	A	276	GLU	OE1-CD-OE2	6.31	130.87	123.30
1	B	257	ARG	NH1-CZ-NH2	-6.30	112.47	119.40
1	A	112	ARG	NE-CZ-NH1	-6.12	117.24	120.30
1	A	137	ASP	CB-CG-OD2	-6.09	112.82	118.30
1	B	11	ASP	CB-CG-OD2	-5.98	112.92	118.30
1	B	176	TYR	CB-CG-CD1	5.95	124.57	121.00
1	A	37	LEU	CB-CG-CD1	5.74	120.76	111.00

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47	ARG	NE-CZ-NH2	5.72	123.16	120.30
1	B	81	TYR	CB-CG-CD2	5.62	124.38	121.00
1	A	240	ASP	CB-CG-OD1	-5.59	113.26	118.30
1	B	268	ARG	NE-CZ-NH2	-5.58	117.51	120.30
1	B	215	CYS	CB-CA-C	5.57	121.54	110.40
1	B	56	ARG	NE-CZ-NH2	5.48	123.04	120.30
1	B	65	ASP	CB-CG-OD1	5.45	123.21	118.30
1	B	181	ASP	CB-CG-OD2	-5.43	113.42	118.30
1	A	112	ARG	CD-NE-CZ	5.40	131.16	123.60
1	B	43	ARG	NE-CZ-NH2	-5.35	117.63	120.30
1	B	81	TYR	CD1-CE1-CZ	5.26	124.53	119.80
1	B	181	ASP	CB-CG-OD1	5.14	122.92	118.30
1	A	289	ASP	CB-CG-OD2	5.11	122.90	118.30
1	A	167[A]	GLU	OE1-CD-OE2	-5.02	117.28	123.30
1	A	167[B]	GLU	OE1-CD-OE2	-5.02	117.28	123.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2500	0	2501	20	1
1	B	2411	0	2433	20	0
2	A	14	0	0	0	0
2	B	14	0	0	0	0
3	B	1	0	0	0	0
4	B	1	0	0	0	0
5	B	4	0	6	0	0
6	A	323	0	0	8	1
6	B	319	0	0	4	1
All	All	5587	0	4940	38	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:262[A]:GLN:OE1	6:B:403:HOH:O	1.68	1.10
1:A:5:LYS:HE3	6:A:652:HOH:O	1.62	0.98
1:B:3[A]:MET:HE1	1:B:246[A]:ILE:HD11	1.49	0.94
1:B:3[A]:MET:CE	1:B:246[A]:ILE:HD11	2.05	0.86
1:A:289:ASP:OD2	1:B:103:LYS:NZ	2.15	0.79
1:A:83:LEU:HD11	1:A:226[A]:CYS:SG	2.25	0.76
1:B:3[A]:MET:CE	1:B:246[A]:ILE:CD1	2.66	0.73
1:B:83:LEU:HD11	1:B:226[A]:CYS:SG	2.28	0.73
1:A:75[A]:GLU:OE1	6:A:401:HOH:O	2.07	0.73
1:A:62:GLU:O	1:A:63:ASP:HB2	1.91	0.69
1:B:3[A]:MET:HE1	1:B:246[A]:ILE:CD1	2.22	0.69
1:A:1:MET:HB2	6:A:698:HOH:O	1.95	0.66
1:B:3[A]:MET:HE3	1:B:246[A]:ILE:CD1	2.29	0.61
1:A:246[B]:ILE:CG2	1:A:271:TYR:CZ	2.84	0.61
1:A:181:ASP:HB2	6:A:559:HOH:O	2.01	0.59
1:B:75[B]:GLU:HG2	6:B:509:HOH:O	2.02	0.58
1:A:1:MET:CB	6:A:698:HOH:O	2.52	0.58
1:A:248:LYS:HE3	6:A:635:HOH:O	2.06	0.55
1:A:21:GLN:HE22	1:A:24:ARG:NH1	2.06	0.53
1:A:246[B]:ILE:HG21	1:A:271:TYR:CE1	2.46	0.50
1:B:246[B]:ILE:CG2	1:B:271:TYR:CZ	2.95	0.49
1:B:112[A]:ARG:NH1	1:B:181:ASP:OD1	2.45	0.49
1:A:246[B]:ILE:CG2	1:A:271:TYR:CE1	2.97	0.48
1:B:196:PHE:O	1:B:200[A]:GLU:HG3	2.15	0.46
1:A:63:ASP:N	6:A:408:HOH:O	2.44	0.46
1:A:246[B]:ILE:HG21	1:A:271:TYR:CZ	2.51	0.46
1:A:73[B]:LYS:HE3	6:A:612:HOH:O	2.18	0.44
1:B:242:SER:HB3	6:B:618:HOH:O	2.19	0.43
1:B:197:LYS:NZ	6:B:405:HOH:O	2.53	0.42
1:A:5:LYS:HB2	1:A:5:LYS:HE2	1.75	0.42
1:A:48:ASP:OD1	1:A:48:ASP:N	2.53	0.42
1:B:3[A]:MET:CE	1:B:246[A]:ILE:HD12	2.50	0.42
1:B:115:GLU:O	1:B:116:LYS:HG2	2.19	0.41
1:B:79:ARG:HD3	1:B:204:LEU:HD22	2.02	0.41
1:A:284:ASP:O	1:A:287:VAL:HG22	2.21	0.40
1:B:102:GLN:O	1:B:209:GLY:HA3	2.22	0.40
1:A:288:GLN:HB3	1:B:103:LYS:HE3	2.03	0.40
1:B:140:LEU:HD23	1:B:162:ASN:HA	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:445:HOH:O	6:A:688:HOH:O[6_555]	2.18	0.02
1:A:1:MET:CE	6:B:447:HOH:O[6_445]	2.19	0.01

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	311/298 (104%)	300 (96%)	9 (3%)	2 (1%)	25 7
1	B	301/298 (101%)	293 (97%)	7 (2%)	1 (0%)	41 18
All	All	612/596 (103%)	593 (97%)	16 (3%)	3 (0%)	29 9

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	63	ASP
1	A	261	ILE
1	B	261	ILE

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	286/271 (106%)	281 (98%)	5 (2%)	60 28
1	B	276/271 (102%)	273 (99%)	3 (1%)	73 48
All	All	562/542 (104%)	554 (99%)	8 (1%)	69 37

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118[A]	SER
1	A	118[B]	SER
1	A	197	LYS
1	A	205	SER
1	A	288	GLN
1	B	11	ASP
1	B	116	LYS
1	B	236	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	GLN
1	A	288	GLN
1	B	193	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	YW9	B	301	-	14,15,15	1.63	3 (21%)	17,21,21	1.15	2 (11%)
2	YW9	A	301	-	14,15,15	0.99	1 (7%)	17,21,21	1.33	4 (23%)
5	DMS	B	304	-	3,3,3	0.39	0	3,3,3	1.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	YW9	B	301	-	-	0/4/4/4	0/2/2/2
2	YW9	A	301	-	-	0/4/4/4	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	YW9	C11-N12	3.47	1.50	1.38
2	B	301	YW9	C7-C2	3.05	1.46	1.39
2	B	301	YW9	C6-C5	-2.48	1.44	1.49
2	A	301	YW9	C7-C2	2.16	1.44	1.39

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	YW9	C14-C8-N4	2.67	124.56	119.50
2	B	301	YW9	C2-N3-N4	2.43	108.47	105.66
2	A	301	YW9	C14-C13-C11	2.39	123.75	120.67
2	B	301	YW9	C5-N4-N3	-2.35	109.72	111.81
2	A	301	YW9	C9-C8-C14	-2.31	117.91	121.33
2	A	301	YW9	C10-C9-C8	2.15	122.19	119.07

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	298/298 (100%)	-0.51	5 (1%) 70 70	14, 22, 46, 73	0
1	B	284/298 (95%)	-0.50	0 100 100	11, 17, 38, 75	0
All	All	582/596 (97%)	-0.50	5 (0%) 84 86	11, 20, 42, 75	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	287	VAL	3.0
1	A	289	ASP	2.7
1	A	62	GLU	2.6
1	A	239	LYS	2.3
1	A	63	ASP	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	YW9	A	301	14/14	0.94	0.05	20,21,26,28	0
2	YW9	B	301	14/14	0.94	0.10	16,19,25,28	0
3	MG	B	302	1/1	0.97	0.18	39,39,39,39	0
5	DMS	B	304	4/4	0.98	0.16	28,29,29,30	0
4	CL	B	303	1/1	1.00	0.03	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.