



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 02:21 AM EDT

PDB ID : 6GCM  
Title : Escherichia coli DPS  
Authors : Kovalenko, V.V.; Loiko, N.G.; Tereshkin, E.V.; Tereshkina, K.B.; Chulichkov, A.L.; Popov, A.N.; Krupyanskii, Y.F.  
Deposited on : 2018-04-18  
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





















Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1544 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	96%
1	b	159	3% 96%
2	B	154	98%
2	C	154	96%
2	E	154	2% 97%
2	F	154	% 97%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	154	 99%
2	H	154	 3% 96%
2	I	154	 97%
2	J	154	 99%
2	K	154	 % 94% 6%
2	L	154	 97%
2	c	154	 97%
2	d	154	 98%
2	g	154	 97%
2	h	154	 96%
2	i	154	 97%
2	j	154	 97%
2	k	154	 97%
2	l	154	 97%
2	m	154	 97%
3	D	156	 % 97%
3	e	156	 % 96%
4	f	151	 97%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 30236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA protection during starvation protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	155	1226	771	214	238	3	0	0	0
1	b	159	1252	785	219	245	3	0	0	0

- Molecule 2 is a protein called DNA protection during starvation protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	154	1214	763	211	237	3	0	0	0
2	C	154	1218	766	212	237	3	0	0	0
2	E	154	1219	767	212	237	3	0	0	0
2	F	154	1219	767	212	237	3	0	0	0
2	G	154	1217	765	212	237	3	0	0	0
2	H	154	1219	767	212	237	3	0	0	0
2	I	154	1216	765	211	237	3	0	0	0
2	J	154	1220	768	212	237	3	0	0	0
2	K	154	1221	768	213	237	3	0	0	0
2	L	154	1217	766	211	237	3	0	0	0
2	c	154	1214	763	211	237	3	0	0	0
2	d	154	1218	766	212	237	3	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	g	154	Total	C	N	O	S	0	0	0
			1219	767	212	237	3			
2	h	154	Total	C	N	O	S	0	0	0
			1217	765	212	237	3			
2	i	154	Total	C	N	O	S	0	0	0
			1219	767	212	237	3			
2	j	154	Total	C	N	O	S	0	0	0
			1216	765	211	237	3			
2	k	154	Total	C	N	O	S	0	0	0
			1220	768	212	237	3			
2	l	154	Total	C	N	O	S	0	0	0
			1221	768	213	237	3			
2	m	154	Total	C	N	O	S	0	0	0
			1217	766	211	237	3			

- Molecule 3 is a protein called DNA protection during starvation protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	156	Total	C	N	O	S	0	0	0
			1236	776	216	241	3			
3	e	156	Total	C	N	O	S	0	0	0
			1236	776	216	241	3			

- Molecule 4 is a protein called DNA protection during starvation protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	f	151	Total	C	N	O	S	0	0	0
			1191	746	209	233	3			

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	40	Total	O	0	0
			40	40		
5	B	37	Total	O	0	0
			37	37		
5	C	36	Total	O	0	0
			36	36		
5	D	41	Total	O	0	0
			41	41		
5	E	47	Total	O	0	0
			47	47		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	F	26	Total O 26 26	0	0
5	G	43	Total O 43 43	0	0
5	H	40	Total O 40 40	0	0
5	I	34	Total O 34 34	0	0
5	J	42	Total O 42 42	0	0
5	K	34	Total O 34 34	0	0
5	L	38	Total O 38 38	0	0
5	b	52	Total O 52 52	0	0
5	c	39	Total O 39 39	0	0
5	d	40	Total O 40 40	0	0
5	e	24	Total O 24 24	0	0
5	f	42	Total O 42 42	0	0
5	g	28	Total O 28 28	0	0
5	h	42	Total O 42 42	0	0
5	i	45	Total O 45 45	0	0
5	j	48	Total O 48 48	0	0
5	k	44	Total O 44 44	0	0
5	l	42	Total O 42 42	0	0
5	m	50	Total O 50 50	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA protection during starvation protein

Chain A:  96%



- Molecule 1: DNA protection during starvation protein

Chain b:  96%




- Molecule 2: DNA protection during starvation protein

Chain B:  98%




- Molecule 2: DNA protection during starvation protein

Chain C:  96%



- Molecule 2: DNA protection during starvation protein


Chain E:  97%



- Molecule 2: DNA protection during starvation protein



- Molecule 2: DNA protection during starvation protein



- Molecule 2: DNA protection during starvation protein



- Molecule 2: DNA protection during starvation protein




- Molecule 2: DNA protection during starvation protein



- Molecule 2: DNA protection during starvation protein



- Molecule 2: DNA protection during starvation protein



- Molecule 2: DNA protection during starvation protein



Chain c:  97%



- Molecule 2: DNA protection during starvation protein

Chain d:  98%



- Molecule 2: DNA protection during starvation protein

Chain g:  97%



- Molecule 2: DNA protection during starvation protein

Chain h:  96%



- Molecule 2: DNA protection during starvation protein

Chain i:  97%



- Molecule 2: DNA protection during starvation protein

Chain j:  97%



- Molecule 2: DNA protection during starvation protein

Chain k:  97%



- Molecule 2: DNA protection during starvation protein

Chain l:  97%




- Molecule 2: DNA protection during starvation protein

Chain m:  97%



- Molecule 3: DNA protection during starvation protein

Chain D:  97%



- Molecule 3: DNA protection during starvation protein

Chain e:  96%



- Molecule 4: DNA protection during starvation protein

Chain f:  97%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	150.15Å 89.42Å 150.65Å 90.00° 90.24° 90.00°	Depositor
Resolution (Å)	100.02 – 2.45 47.68 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.8 (100.02-2.45) 99.8 (47.68-2.45)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.33 (at 2.45Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.203 , 0.258 0.210 , 0.264	Depositor DCC
$R_{free}$ test set	7407 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.4	Xtrriage
Anisotropy	0.051	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 19.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.014 for l,k,-h 0.027 for h,-k,-l 0.022 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	30236	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.73	0/1244	0.80	0/1684
1	b	0.73	0/1270	0.84	0/1720
2	B	0.70	0/1232	0.82	0/1670
2	C	0.70	0/1236	0.83	0/1674
2	E	0.71	0/1237	0.80	0/1675
2	F	0.65	0/1237	0.78	0/1675
2	G	0.68	0/1234	0.79	0/1672
2	H	0.63	0/1237	0.81	0/1675
2	I	0.66	0/1234	0.78	0/1672
2	J	0.66	0/1238	0.77	0/1676
2	K	0.65	0/1239	0.81	0/1677
2	L	0.64	0/1235	0.80	0/1673
2	c	0.68	0/1232	0.77	0/1670
2	d	0.68	0/1236	0.84	0/1674
2	g	0.65	0/1237	0.78	0/1675
2	h	0.69	0/1234	0.81	0/1672
2	i	0.67	0/1237	0.85	0/1675
2	j	0.74	0/1234	0.83	0/1672
2	k	0.69	0/1238	0.83	0/1676
2	l	0.71	0/1239	0.82	0/1677
2	m	0.67	0/1235	0.79	0/1673
3	D	0.66	0/1254	0.78	0/1698
3	e	0.70	0/1254	0.82	0/1698
4	f	0.72	0/1208	0.80	0/1635
All	All	0.68	0/29711	0.81	0/40238

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	153/159 (96%)	149 (97%)	4 (3%)	0	100	100
1	b	157/159 (99%)	153 (98%)	4 (2%)	0	100	100
2	B	152/154 (99%)	148 (97%)	4 (3%)	0	100	100
2	C	152/154 (99%)	148 (97%)	4 (3%)	0	100	100
2	E	152/154 (99%)	146 (96%)	6 (4%)	0	100	100
2	F	152/154 (99%)	145 (95%)	7 (5%)	0	100	100
2	G	152/154 (99%)	151 (99%)	1 (1%)	0	100	100
2	H	152/154 (99%)	147 (97%)	3 (2%)	2 (1%)	12	11
2	I	152/154 (99%)	145 (95%)	7 (5%)	0	100	100
2	J	152/154 (99%)	146 (96%)	6 (4%)	0	100	100
2	K	152/154 (99%)	146 (96%)	6 (4%)	0	100	100
2	L	152/154 (99%)	152 (100%)	0	0	100	100
2	c	152/154 (99%)	148 (97%)	4 (3%)	0	100	100
2	d	152/154 (99%)	148 (97%)	4 (3%)	0	100	100
2	g	152/154 (99%)	147 (97%)	5 (3%)	0	100	100
2	h	152/154 (99%)	151 (99%)	1 (1%)	0	100	100
2	i	152/154 (99%)	150 (99%)	2 (1%)	0	100	100
2	j	152/154 (99%)	148 (97%)	4 (3%)	0	100	100
2	k	152/154 (99%)	150 (99%)	2 (1%)	0	100	100
2	l	152/154 (99%)	146 (96%)	6 (4%)	0	100	100
2	m	152/154 (99%)	148 (97%)	4 (3%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	D	154/156 (99%)	150 (97%)	4 (3%)	0	100	100
3	e	154/156 (99%)	152 (99%)	2 (1%)	0	100	100
4	f	149/151 (99%)	144 (97%)	5 (3%)	0	100	100
All	All	3655/3707 (99%)	3558 (97%)	95 (3%)	2 (0%)	51	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	23	ASP
2	H	24	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/136 (97%)	130 (98%)	2 (2%)	65	76
1	b	135/136 (99%)	129 (96%)	6 (4%)	28	37
2	B	130/132 (98%)	127 (98%)	3 (2%)	50	63
2	C	131/132 (99%)	125 (95%)	6 (5%)	27	35
2	E	131/132 (99%)	127 (97%)	4 (3%)	40	52
2	F	131/132 (99%)	126 (96%)	5 (4%)	33	43
2	G	130/132 (98%)	128 (98%)	2 (2%)	65	76
2	H	131/132 (99%)	127 (97%)	4 (3%)	40	52
2	I	130/132 (98%)	126 (97%)	4 (3%)	40	52
2	J	131/132 (99%)	130 (99%)	1 (1%)	81	88
2	K	132/132 (100%)	123 (93%)	9 (7%)	16	19
2	L	130/132 (98%)	126 (97%)	4 (3%)	40	52
2	c	130/132 (98%)	126 (97%)	4 (3%)	40	52
2	d	131/132 (99%)	128 (98%)	3 (2%)	50	63
2	g	131/132 (99%)	127 (97%)	4 (3%)	40	52

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	h	130/132 (98%)	124 (95%)	6 (5%)	27	35
2	i	131/132 (99%)	127 (97%)	4 (3%)	40	52
2	j	130/132 (98%)	125 (96%)	5 (4%)	33	43
2	k	131/132 (99%)	126 (96%)	5 (4%)	33	43
2	l	132/132 (100%)	128 (97%)	4 (3%)	41	52
2	m	130/132 (98%)	126 (97%)	4 (3%)	40	52
3	D	134/134 (100%)	130 (97%)	4 (3%)	41	52
3	e	134/134 (100%)	128 (96%)	6 (4%)	27	36
4	f	128/129 (99%)	124 (97%)	4 (3%)	40	52
All	All	3146/3177 (99%)	3043 (97%)	103 (3%)	38	49

All (103) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	HIS
1	A	116	ASP
2	B	51	HIS
2	B	77	LEU
2	B	95	THR
2	C	24	SER
2	C	51	HIS
2	C	85	VAL
2	C	119	LYS
2	C	132	VAL
2	C	141	ASP
3	D	51	HIS
3	D	85	VAL
3	D	100	SER
3	D	101	LYS
2	E	20	ASP
2	E	24	SER
2	E	51	HIS
2	E	113	ASN
2	F	32	LEU
2	F	51	HIS
2	F	55	ARG
2	F	85	VAL
2	F	163	GLU
2	G	77	LEU

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
2	G	140	LYS
2	H	51	HIS
2	H	100	SER
2	H	106	SER
2	H	134	LYS
2	I	24	SER
2	I	25	GLU
2	I	92	LEU
2	I	134	LYS
2	J	95	THR
2	K	14	LEU
2	K	32	LEU
2	K	51	HIS
2	K	77	LEU
2	K	85	VAL
2	K	100	SER
2	K	106	SER
2	K	124	ARG
2	K	134	LYS
2	L	14	LEU
2	L	31	GLU
2	L	85	VAL
2	L	138	GLU
1	b	13	ASN
1	b	15	LEU
1	b	17	THR
1	b	24	SER
1	b	51	HIS
1	b	55	ARG
2	c	24	SER
2	c	51	HIS
2	c	85	VAL
2	c	95	THR
2	d	24	SER
2	d	100	SER
2	d	106	SER
3	e	27	LYS
3	e	32	LEU
3	e	35	ARG
3	e	85	VAL
3	e	100	SER
3	e	155	LEU

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
4	f	24	SER
4	f	31	GLU
4	f	51	HIS
4	f	113	ASN
2	g	32	LEU
2	g	51	HIS
2	g	85	VAL
2	g	106	SER
2	h	17	THR
2	h	24	SER
2	h	51	HIS
2	h	85	VAL
2	h	95	THR
2	h	113	ASN
2	i	32	LEU
2	i	51	HIS
2	i	105	LYS
2	i	113	ASN
2	j	20	ASP
2	j	51	HIS
2	j	77	LEU
2	j	100	SER
2	j	119	LYS
2	k	55	ARG
2	k	85	VAL
2	k	100	SER
2	k	119	LYS
2	k	134	LYS
2	l	14	LEU
2	l	31	GLU
2	l	32	LEU
2	l	51	HIS
2	m	23	ASP
2	m	24	SER
2	m	51	HIS
2	m	77	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (84) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	112	HIS
1	A	113	ASN

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	115	GLN
1	A	117	HIS
2	B	99	ASN
2	B	113	ASN
2	B	115	GLN
2	B	117	HIS
2	C	112	HIS
2	C	115	GLN
2	C	117	HIS
3	D	13	ASN
3	D	113	ASN
3	D	115	GLN
3	D	117	HIS
2	E	99	ASN
2	E	112	HIS
2	E	113	ASN
2	E	115	GLN
2	E	117	HIS
2	F	99	ASN
2	F	113	ASN
2	F	115	GLN
2	F	117	HIS
2	G	99	ASN
2	G	112	HIS
2	G	113	ASN
2	G	115	GLN
2	G	117	HIS
2	H	115	GLN
2	H	117	HIS
2	I	113	ASN
2	I	115	GLN
2	I	117	HIS
2	J	112	HIS
2	J	113	ASN
2	J	115	GLN
2	J	117	HIS
2	K	113	ASN
2	K	115	GLN
2	K	117	HIS
2	L	99	ASN
2	L	112	HIS
2	L	113	ASN

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
2	L	115	GLN
2	L	117	HIS
1	b	13	ASN
1	b	99	ASN
1	b	115	GLN
1	b	117	HIS
2	c	113	ASN
2	c	115	GLN
2	c	117	HIS
2	d	115	GLN
2	d	117	HIS
3	e	99	ASN
3	e	113	ASN
3	e	115	GLN
3	e	117	HIS
4	f	99	ASN
4	f	113	ASN
4	f	115	GLN
4	f	117	HIS
2	g	112	HIS
2	g	113	ASN
2	g	115	GLN
2	g	117	HIS
2	h	115	GLN
2	h	117	HIS
2	i	99	ASN
2	i	113	ASN
2	i	115	GLN
2	i	117	HIS
2	j	113	ASN
2	j	115	GLN
2	j	117	HIS
2	k	113	ASN
2	k	115	GLN
2	k	117	HIS
2	l	115	GLN
2	l	117	HIS
2	m	113	ASN
2	m	115	GLN
2	m	117	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	155/159 (97%)	-0.42	0 100 100	23, 34, 53, 66	0
1	b	159/159 (100%)	-0.28	4 (2%) 57 53	24, 32, 55, 162	0
2	B	154/154 (100%)	-0.46	0 100 100	27, 35, 49, 66	0
2	C	154/154 (100%)	-0.42	0 100 100	23, 35, 52, 68	0
2	E	154/154 (100%)	-0.43	3 (1%) 66 64	23, 30, 45, 63	0
2	F	154/154 (100%)	-0.31	1 (0%) 89 89	30, 39, 53, 69	0
2	G	154/154 (100%)	-0.52	0 100 100	21, 32, 46, 72	0
2	H	154/154 (100%)	-0.33	4 (2%) 56 52	29, 39, 54, 67	0
2	I	154/154 (100%)	-0.43	0 100 100	24, 34, 51, 65	0
2	J	154/154 (100%)	-0.40	0 100 100	30, 38, 53, 65	0
2	K	154/154 (100%)	-0.41	1 (0%) 89 89	27, 38, 54, 69	0
2	L	154/154 (100%)	-0.44	0 100 100	26, 35, 48, 68	0
2	c	154/154 (100%)	-0.44	0 100 100	27, 36, 50, 70	0
2	d	154/154 (100%)	-0.53	0 100 100	24, 32, 44, 56	0
2	g	154/154 (100%)	-0.44	0 100 100	28, 36, 51, 66	0
2	h	154/154 (100%)	-0.49	0 100 100	25, 34, 48, 63	0
2	i	154/154 (100%)	-0.48	0 100 100	24, 33, 46, 57	0
2	j	154/154 (100%)	-0.45	0 100 100	24, 31, 43, 57	0
2	k	154/154 (100%)	-0.51	0 100 100	25, 33, 46, 57	0
2	l	154/154 (100%)	-0.51	0 100 100	24, 32, 50, 65	0
2	m	154/154 (100%)	-0.53	0 100 100	23, 33, 47, 66	0
3	D	156/156 (100%)	-0.47	1 (0%) 89 89	25, 36, 54, 102	0
3	e	156/156 (100%)	-0.42	1 (0%) 89 89	26, 37, 56, 109	0
4	f	151/151 (100%)	-0.53	0 100 100	24, 33, 47, 56	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
All	All	3703/3707 (99%)	-0.44	15 (0%) 92 93	21, 34, 51, 162	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	b	11	ALA	10.9
3	e	12	THR	10.3
1	b	12	THR	8.6
2	H	22	SER	7.0
1	b	9	SER	6.7
2	E	20	ASP	4.2
1	b	10	LYS	3.5
2	H	21	VAL	3.2
3	D	12	THR	3.1
2	E	21	VAL	3.1
2	H	137	GLY	2.7
2	F	15	LEU	2.6
2	H	23	ASP	2.4
2	K	77	LEU	2.3
2	E	32	LEU	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.