



Full wwPDB EM Validation Report ⓘ

Nov 3, 2024 – 04:52 PM EST

PDB ID : 8GEB
EMDB ID : EMD-29964
Title : CryoEM structure of beta-2-adrenergic receptor in complex with nucleotide-free Gs heterotrimer (#12 of 20)
Authors : Papasergi-Scott, M.M.; Skiniotis, G.
Deposited on : 2023-03-06
Resolution : 3.60 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev113
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	58% 39%
2	B	343	80% 16%
3	G	71	65% 6% 30%
4	R	460	54% 7% 39%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6223 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide-binding protein G(s) subunit alpha isoforms short.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	233	1650	1064	299	281	6	0	0

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	329	2301	1457	418	408	18	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP P62873
B	-1	SER	-	expression tag	UNP P62873
B	0	SER	-	expression tag	UNP P62873
B	1	GLY	-	expression tag	UNP P62873

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	G	50	319	203	57	58	1	0	0

- Molecule 4 is a protein called Beta-2 adrenergic receptor.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	R	281	1938	1282	316	329	11	0	0

There are 54 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
R	-30	MET	-	initiating methionine	UNP P07550
R	-29	LYS	-	expression tag	UNP P07550
R	-28	THR	-	expression tag	UNP P07550
R	-27	ILE	-	expression tag	UNP P07550
R	-26	ILE	-	expression tag	UNP P07550
R	-25	ALA	-	expression tag	UNP P07550
R	-24	LEU	-	expression tag	UNP P07550
R	-23	SER	-	expression tag	UNP P07550
R	-22	TYR	-	expression tag	UNP P07550
R	-21	ILE	-	expression tag	UNP P07550
R	-20	PHE	-	expression tag	UNP P07550
R	-19	CYS	-	expression tag	UNP P07550
R	-18	LEU	-	expression tag	UNP P07550
R	-17	VAL	-	expression tag	UNP P07550
R	-16	PHE	-	expression tag	UNP P07550
R	-15	ALA	-	expression tag	UNP P07550
R	-14	ASP	-	expression tag	UNP P07550
R	-13	TYR	-	expression tag	UNP P07550
R	-12	LYS	-	expression tag	UNP P07550
R	-11	ASP	-	expression tag	UNP P07550
R	-10	ASP	-	expression tag	UNP P07550
R	-9	ASP	-	expression tag	UNP P07550
R	-8	ASP	-	expression tag	UNP P07550
R	-7	ALA	-	expression tag	UNP P07550
R	9	ARG	GLY	variant	UNP P07550
R	18	GLU	-	insertion	UNP P07550
R	19	ASN	-	insertion	UNP P07550
R	20	LEU	-	insertion	UNP P07550
R	21	TYR	-	insertion	UNP P07550
R	22	PHE	-	insertion	UNP P07550
R	23	GLN	-	insertion	UNP P07550
R	24	GLY	-	insertion	UNP P07550
R	27	GLN	GLU	variant	UNP P07550
R	96	THR	MET	conflict	UNP P07550
R	98	THR	MET	conflict	UNP P07550
R	187	GLU	ASN	conflict	UNP P07550
R	366	LEU	-	insertion	UNP P07550
R	367	GLU	-	insertion	UNP P07550
R	368	VAL	-	insertion	UNP P07550
R	369	LEU	-	insertion	UNP P07550
R	370	PHE	-	insertion	UNP P07550
R	371	GLN	-	insertion	UNP P07550
R	372	GLY	-	insertion	UNP P07550

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	373	PRO	-	insertion	UNP P07550
R	386	ALA	CYS	conflict	UNP P07550
R	414	ALA	CYS	conflict	UNP P07550
R	422	GLU	-	expression tag	UNP P07550
R	423	THR	-	expression tag	UNP P07550
R	424	SER	-	expression tag	UNP P07550
R	425	GLN	-	expression tag	UNP P07550
R	426	VAL	-	expression tag	UNP P07550
R	427	ALA	-	expression tag	UNP P07550
R	428	PRO	-	expression tag	UNP P07550
R	429	ALA	-	expression tag	UNP P07550

- Molecule 5 is (5R,6R)-6-(methylamino)-5,6,7,8-tetrahydronaphthalene-1,2,5-triol (three-letter code: G1I) (formula: C₁₁H₁₅NO₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
5	R	1	15	11	1	3	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Guanine nucleotide-binding protein G(s) subunit alpha isoforms short



- Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1



- Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2



- Molecule 4: Beta-2 adrenergic receptor



MET
 LYS
 THR
 ILE
 ILE
 ILE
 ALA
 LEU
 SER
 TYR
 ILE
 PHE
 CYS
 LEU
 VAL
 PHE
 ALA
 ASP
 TYR
 LYS
 ASP
 ASP
 ASP
 ASP
 ALA
 MET
 GLY
 GLN
 PRO
 GLY
 ASN
 GLY
 SER
 ALA
 PHE
 LEU
 LEU
 ALA
 PRO
 ASN
 ARG
 SER
 HIS
 HIS
 ALA
 PRO
 ASP
 HIS
 ASP
 VAL
 GLU
 ASN
 LEU
 TYR
 PHE
 GLN
 GLY
 THR
 GLN
 GLN
 ARG
 ASP

GLU
 V51
 N51
 N54
 V54
 I58
 C77
 L80
 V81
 M82
 V86
 W99
 W109
 D113
 V114
 L115
 C116
 C125
 Y132
 K149
 G162
 I169
 Y174
 T189
 I201
 I205
 Y209
 V210
 P211
 V218
 R239
 PHE
 HIS
 VAL
 GLN
 ASN
 LEU
 SER
 GLN
 VAL

GLU
 GLN
 ASP
 GLY
 THR
 GLY
 HIS
 GLY
 LEU
 ARG
 ARG
 ARG
 SER
 SER
 LYS
 PHE
 CYS
 LEU
 LYS
 GLU
 H269
 K270
 A271
 T281
 P288
 N293
 H296
 L311
 I325
 Y326
 L340
 CYS
 LEU
 ARG
 ARG
 SER
 SER
 SER
 LEU
 LYS
 ALA
 TYR
 GLY
 ASN
 GLY
 TYR
 SER
 SER
 ASN
 ASN
 GLY
 ASN
 THR
 THR
 ASN
 SER
 THR
 LEU
 THR
 GLY
 GLU

GLN
 SER
 GLY
 LEU
 GLU
 VAL
 LEU
 PHE
 GLN
 GLY
 PRO
 TYR
 HIS
 VAL
 GLU
 GLN
 GLU
 LYS
 GLU
 ASN
 LYS
 LEU
 LEU
 ALA
 GLU
 ASP
 LEU
 PRO
 GLY
 THR
 GLU
 ASP
 PHE
 VAL
 GLY
 HIS
 GLN
 GLY
 THR
 VAL
 PRO
 SER
 ASP
 ASN
 ILE
 ASP
 SER
 GLN
 GLY
 ARG
 ASN
 ALA
 ALA
 THR
 ASN
 ASP
 SER
 LEU
 LEU
 GLU

THR
 SER
 GLN
 VAL
 ALA
 PRO
 ALA

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	105147	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	52	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	1.548	Depositor
Minimum map value	-0.895	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.028	Depositor
Recommended contour level	0.23	Depositor
Map size (\AA)	317.99997, 317.99997, 317.99997	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.06, 1.06, 1.06	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: G1I

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1683	0.44	0/2296
2	B	0.27	0/2347	0.51	0/3201
3	G	0.24	0/325	0.40	0/451
4	R	0.26	0/1987	0.41	0/2739
All	All	0.26	0/6342	0.46	0/8687

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1650	0	1431	8	0
2	B	2301	0	2066	42	0
3	G	319	0	258	4	0
4	R	1938	0	1648	22	0
5	R	15	0	0	1	0
All	All	6223	0	5403	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:200:VAL:HG23	2:B:209:LYS:O	1.69	0.92
2:B:200:VAL:HG12	2:B:234:PHE:HE1	1.52	0.75
2:B:58:ILE:HD11	2:B:330:GLY:HA3	1.80	0.63
4:R:293:ASN:HA	4:R:296:HIS:CE1	2.33	0.63
4:R:325:ILE:HG13	4:R:326:TYR:HD1	1.64	0.61
2:B:315:VAL:HA	2:B:331:SER:HA	1.83	0.61
2:B:149:CYS:HB3	2:B:157:ILE:HD11	1.86	0.58
2:B:200:VAL:HG23	2:B:209:LYS:C	2.22	0.58
2:B:137:ARG:NH1	2:B:171:ILE:O	2.37	0.57
1:A:283:ARG:O	1:A:356:ARG:NH1	2.38	0.56
2:B:73:ALA:HB2	2:B:79:LEU:CD1	2.35	0.56
2:B:81:ILE:HD12	2:B:91:HIS:HB2	1.88	0.55
2:B:82:TRP:NE1	2:B:89:LYS:HG2	2.23	0.54
2:B:274:THR:HG21	2:B:316:SER:HA	1.90	0.53
2:B:142:HIS:HE2	2:B:159:THR:HG1	1.57	0.53
2:B:200:VAL:HG12	2:B:234:PHE:CE1	2.41	0.52
2:B:149:CYS:O	2:B:150:ARG:NH1	2.43	0.51
1:A:56:ILE:HD11	1:A:292:ASN:HD21	1.75	0.51
2:B:200:VAL:HG22	2:B:201:SER:N	2.26	0.51
2:B:225:HIS:CE1	2:B:251:ARG:HD2	2.46	0.50
2:B:82:TRP:HE1	2:B:89:LYS:HG2	1.76	0.50
2:B:249:THR:HG22	2:B:265:SER:HB3	1.94	0.50
2:B:30:LEU:HD13	3:G:34:ALA:HB1	1.94	0.49
2:B:198:LEU:HD22	2:B:210:LEU:HD21	1.94	0.49
4:R:51:ASN:HB3	4:R:80:LEU:HB2	1.95	0.49
2:B:82:TRP:CZ3	2:B:336:LEU:HD12	2.46	0.49
2:B:308:LEU:HD23	2:B:339:TRP:CE3	2.48	0.49
2:B:73:ALA:HB2	2:B:79:LEU:HD13	1.95	0.48
2:B:145:TYR:O	2:B:162:GLY:N	2.44	0.48
2:B:82:TRP:CE3	2:B:336:LEU:HD12	2.49	0.47
4:R:82:MET:HB2	4:R:116:CYS:HB3	1.96	0.47
2:B:211:TRP:HE1	2:B:218:CYS:HB3	1.79	0.47
2:B:49:ARG:NH2	3:G:61:PHE:O	2.48	0.47
1:A:273:PHE:HE1	1:A:287:VAL:HG11	1.80	0.46
2:B:82:TRP:CD1	2:B:89:LYS:HG2	2.51	0.46
2:B:232:ILE:HG12	2:B:243:THR:HG22	1.97	0.46
4:R:54:VAL:O	4:R:58:ILE:HD12	2.16	0.46
4:R:281:THR:HG21	4:R:325:ILE:HD13	1.97	0.46
2:B:279:SER:OG	3:G:48:ASP:OD2	2.34	0.46
4:R:125:CYS:HB2	4:R:211:PRO:HB3	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:R:132:TYR:HB2	4:R:218:VAL:HB	1.98	0.45
2:B:211:TRP:HD1	2:B:218:CYS:HA	1.82	0.45
2:B:301:LYS:HA	2:B:301:LYS:HD2	1.74	0.45
4:R:205:ILE:HA	4:R:209:TYR:HB2	1.98	0.45
4:R:169:ILE:HA	4:R:174:TYR:HD2	1.81	0.45
4:R:293:ASN:HA	4:R:296:HIS:HE1	1.77	0.45
1:A:274:LYS:HE3	1:A:274:LYS:HB2	1.69	0.44
4:R:86:VAL:HG12	4:R:109:TRP:CD1	2.51	0.44
4:R:201:ILE:O	4:R:205:ILE:HG22	2.17	0.44
1:A:295:ASP:OD1	1:A:296:LEU:N	2.51	0.44
2:B:27:ASP:H	3:G:30:VAL:HG12	1.83	0.44
4:R:169:ILE:HA	4:R:174:TYR:CD2	2.53	0.44
1:A:393:LEU:O	4:R:271:ALA:HB2	2.18	0.43
4:R:288:PRO:HB2	4:R:311:LEU:HD13	2.00	0.43
2:B:266:HIS:HB3	2:B:269:ILE:HG12	2.00	0.43
4:R:82:MET:HB2	4:R:116:CYS:CB	2.48	0.43
2:B:180:PHE:CE1	2:B:216:GLY:HA2	2.54	0.43
4:R:113:ASP:OD1	5:R:501:GII:OAM	2.37	0.42
2:B:264:TYR:CE2	2:B:285:LEU:HD13	2.54	0.42
2:B:34:THR:HG21	2:B:300:LEU:O	2.19	0.42
1:A:327:GLU:O	1:A:329:GLY:N	2.49	0.42
2:B:283:ARG:NE	2:B:298:ASP:OD1	2.49	0.42
4:R:99:TRP:HB3	4:R:189:THR:O	2.20	0.42
2:B:155:ASN:O	2:B:171:ILE:HG12	2.20	0.41
1:A:346:LEU:HD23	1:A:346:LEU:HA	1.93	0.41
2:B:58:ILE:HA	2:B:74:SER:HA	2.01	0.41
2:B:137:ARG:HH12	2:B:172:GLU:HA	1.86	0.41
4:R:77:CYS:O	4:R:81:VAL:HG23	2.20	0.41
2:B:200:VAL:CG2	2:B:209:LYS:O	2.56	0.40
4:R:115:LEU:HD13	4:R:162:GLY:HA2	2.03	0.40
4:R:149:LYS:HE2	4:R:149:LYS:HB2	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/380 (59%)	223 (99%)	2 (1%)	0	100	100
2	B	325/343 (95%)	311 (96%)	14 (4%)	0	100	100
3	G	48/71 (68%)	48 (100%)	0	0	100	100
4	R	277/460 (60%)	272 (98%)	5 (2%)	0	100	100
All	All	875/1254 (70%)	854 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/341 (39%)	132 (100%)	0	100	100
2	B	196/284 (69%)	196 (100%)	0	100	100
3	G	21/58 (36%)	21 (100%)	0	100	100
4	R	151/399 (38%)	151 (100%)	0	100	100
All	All	500/1082 (46%)	500 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	G1I	R	501	-	16,16,16	2.30	5 (31%)	19,23,23	1.50	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	G1I	R	501	-	-	2/2/15/15	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	R	501	G1I	CAJ-CAI	5.21	1.58	1.52
5	R	501	G1I	CAC-CAD	4.65	1.45	1.40
5	R	501	G1I	CAD-CAE	3.07	1.44	1.40
5	R	501	G1I	CAF-CAJ	2.60	1.55	1.51
5	R	501	G1I	CAH-CAG	2.20	1.56	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	R	501	G1I	CAG-CAH-CAI	3.39	117.06	110.85
5	R	501	G1I	CAB-CAC-CAD	-3.07	116.91	120.09
5	R	501	G1I	CAH-CAG-CAE	2.42	117.70	112.92

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	R	501	G1I	CAJ-CAI-NAN-CAO
5	R	501	G1I	CAH-CAI-NAN-CAO

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	R	501	G1I	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-29964. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150

6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 149




Y Index: 162




Z Index: 169


6.3.2 Raw map



X Index: 142



Y Index: 162




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map




6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.23. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_29964_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 52 nm³; this corresponds to an approximate mass of 47 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.278 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.278 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.60	-	-
Author-provided FSC curve	3.57	4.04	3.63
Unmasked-calculated*	4.36	7.49	4.45

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.36 differs from the reported value 3.6 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-29964 and PDB model 8GEB. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.23 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.23).











9.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.23) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8460	 0.4870
A	 0.8520	 0.4880
B	 0.8580	 0.5040
G	 0.8640	 0.4660
R	 0.8260	 0.4700

