



wwPDB NMR Structure Validation Summary Report

Sep 16, 2024 – 01:30 pm BST

PDB ID : 9GE9
BMRB ID : 34950
Title : Trp-cage fortified Tc5b-Exenatide chimera (Ex-4-Tc5bQR) at 277K
Authors : Horvath, D.
Deposited on : 2024-08-07

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 53%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	25	

2 Ensemble composition and analysis

This entry contains 10 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:6-A:24 (19)	0.42	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 4, 5, 6, 7, 8, 9
2	3, 10
Single-model clusters	1

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 390 atoms, of which 193 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Trp-cage fortified Tc5b-Exenatide chimera (Ex-4-Tc5bQR).


Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
1	A	25	390	124	193	35	38	0

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Trp-cage fortified Tc5b-Exenatide chimera (Ex-4-Tc5bQR)



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 5. Colouring as in section 4.1 above.

- Molecule 1: Trp-cage fortified Tc5b-Exenatide chimera (Ex-4-Tc5bQR)



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 20 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CcpNmr Analysis Assign	refinement	2.4.1.
ARIA2alpha	structure calculation	2.3.1.

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	168
Number of shifts mapped to atoms	168
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	53%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.92±0.04	0±0/158 (0.0± 0.0%)	1.14±0.06	1±1/216 (0.4± 0.2%)
All	All	0.92	0/1580 (0.0%)	1.14	9/2160 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.6±0.5
All	All	0	6

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	6	ARG	NE-CZ-NH2	-9.20	115.70	120.30	9	8
1	A	12	LEU	N-CA-CB	-5.10	100.20	110.40	1	1

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	6	ARG	Sidechain	6

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	151	154	154	20±3
All	All	1510	1540	1540	195

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 64.

5 of 78 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:11:TRP:CZ2	1:A:17:PRO:HD3	1.08	1.83	4	1
1:A:11:TRP:CH2	1:A:17:PRO:HD3	0.85	2.06	4	2
1:A:7:LEU:HD13	1:A:8:TYR:N	0.84	1.88	9	9
1:A:9:ILE:O	1:A:13:LYS:HG2	0.73	1.83	8	7
1:A:12:LEU:HD23	1:A:12:LEU:H	0.69	1.47	9	5

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	19/25 (76%)	16±1 (83±5%)	1±1 (6±5%)	2±1 (11±4%)	1	9
All	All	190/250 (76%)	158 (83%)	12 (6%)	20 (11%)	1	9

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	16	GLY	9
1	A	18	SER	9
1	A	24	PRO	1
1	A	17	PRO	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	16/21 (76%)	12±1 (75±9%)	4±1 (25±9%)	2	23
All	All	160/210 (76%)	120 (75%)	40 (25%)	2	23

5 of 9 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	6	ARG	9
1	A	7	LEU	9
1	A	12	LEU	6
1	A	10	GLN	5
1	A	14	GLN	5

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 53% for the well-defined parts and 50% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *nef_chemical_shift_list_ShiftList_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	168
Number of shifts mapped to atoms	168
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 53%, i.e. 142 atoms were assigned a chemical shift out of a possible 267. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	37/90 (41%)	37/37 (100%)	0/38 (0%)	0/15 (0%)
Sidechain	95/156 (61%)	95/101 (94%)	0/46 (0%)	0/9 (0%)
Aromatic	10/21 (48%)	10/10 (100%)	0/10 (0%)	0/1 (0%)
Overall	142/267 (53%)	142/148 (96%)	0/94 (0%)	0/25 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	16	GLY	HA2	1.14	2.15 – 5.77	-7.8
1	A	23	PRO	HA	2.71	2.78 – 6.00	-5.2

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1707
Intra-residue ($ i-j =0$)	924
Sequential ($ i-j =1$)	414
Medium range ($ i-j >1$ and $ i-j <5$)	225
Long range ($ i-j \geq 5$)	144
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	68.3
Number of long range restraints per residue ¹	5.8

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	84.7	0.2
0.2-0.5 (Medium)	161.5	0.5
>0.5 (Large)	185.0	4.93

8.2.2 Average number of dihedral-angle violations per model

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis [i](#)


9.1 Summary of distance violations [i](#)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	924	54.1	344	37.2	20.2	102	11.0	6.0
Backbone-Backbone	75	4.4	10	13.3	0.6	6	8.0	0.4
Backbone-Sidechain	381	22.3	173	45.4	10.1	27	7.1	1.6
Sidechain-Sidechain	468	27.4	161	34.4	9.4	69	14.7	4.0
Sequential ($i-j =1$)	414	24.3	213	51.4	12.5	42	10.1	2.5
Backbone-Backbone	108	6.3	23	21.3	1.3	5	4.6	0.3
Backbone-Sidechain	228	13.4	128	56.1	7.5	25	11.0	1.5
Sidechain-Sidechain	78	4.6	62	79.5	3.6	12	15.4	0.7
Medium range ($i-j >1$ & $i-j <5$)	225	13.2	131	58.2	7.7	36	16.0	2.1
Backbone-Backbone	57	3.3	11	19.3	0.6	1	1.8	0.1
Backbone-Sidechain	102	6.0	68	66.7	4.0	18	17.6	1.1
Sidechain-Sidechain	66	3.9	52	78.8	3.0	17	25.8	1.0
Long range ($i-j \geq 5$)	144	8.4	63	43.8	3.7	20	13.9	1.2
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	33	1.9	14	42.4	0.8	3	9.1	0.2
Sidechain-Sidechain	111	6.5	49	44.1	2.9	17	15.3	1.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1707	100.0	751	44.0	44.0	200	11.7	11.7
Backbone-Backbone	240	14.1	44	18.3	2.6	12	5.0	0.7
Backbone-Sidechain	744	43.6	383	51.5	22.4	73	9.8	4.3
Sidechain-Sidechain	723	42.4	324	44.8	19.0	115	15.9	6.7

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis


9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	201	136	78	34	0	449	0.64	3.77	0.61	0.39
2	201	135	78	34	0	448	0.63	4.27	0.61	0.41
3	205	136	78	35	0	454	0.65	4.68	0.68	0.42
4	196	129	79	35	0	439	0.67	4.52	0.75	0.39
5	203	128	73	37	0	441	0.68	4.46	0.71	0.4
6	198	115	69	30	0	412	0.69	4.65	0.69	0.43
7	205	129	70	34	0	438	0.7	4.2	0.74	0.45
8	207	129	72	42	0	450	0.7	3.93	0.71	0.44
9	206	124	70	33	0	433	0.7	4.93	0.71	0.42
10	192	112	72	33	0	409	0.66	4.61	0.66	0.42

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 956(IR:580, SQ:201, MR:94, LR:81, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
72	38	20	25	0	155	1	10.0
28	16	16	0	0	60	2	20.0
19	14	12	2	0	47	3	30.0


Continued on next page...

Continued from previous page...

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
17	12	9	0	0	38	4	40.0
30	14	10	3	0	57	5	50.0
16	11	6	3	0	36	6	60.0
14	11	11	1	0	37	7	70.0
17	15	7	5	0	44	8	80.0
29	40	4	4	0	77	9	90.0
102	42	36	20	0	200	10	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶ Number of models with violations

9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)




9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models

in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(2,535)	1:3:A:GLU:HG3	1:7:A:LEU:H	10	3.75	0.99	4.05
(4,59)	1:3:A:GLU:HG3	1:7:A:LEU:H	10	3.75	0.99	4.05
(1,535)	1:3:A:GLU:HG3	1:7:A:LEU:H	10	3.24	0.99	3.54
(2,539)	1:3:A:GLU:HB3	1:6:A:ARG:H	10	2.72	0.39	2.82
(4,62)	1:3:A:GLU:HB3	1:6:A:ARG:H	10	2.72	0.39	2.82
(2,514)	1:7:A:LEU:HD11	1:10:A:GLN:HG3	10	2.67	1.29	3.1
(2,514)	1:7:A:LEU:HD12	1:10:A:GLN:HG3	10	2.67	1.29	3.1
(2,514)	1:7:A:LEU:HD13	1:10:A:GLN:HG3	10	2.67	1.29	3.1
(4,54)	1:7:A:LEU:HD11	1:10:A:GLN:HG3	10	2.67	1.29	3.1
(4,54)	1:7:A:LEU:HD12	1:10:A:GLN:HG3	10	2.67	1.29	3.1
(4,54)	1:7:A:LEU:HD13	1:10:A:GLN:HG3	10	2.67	1.29	3.1
(2,538)	1:10:A:GLN:HE21	1:14:A:GLN:HG2	10	2.35	0.98	2.58
(4,61)	1:10:A:GLN:HE21	1:14:A:GLN:HG2	10	2.35	0.98	2.58
(2,421)	1:3:A:GLU:HG2	1:5:A:VAL:HG11	10	2.18	0.64	2.32

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(4,42)	1:6:A:ARG:HH11	1:7:A:LEU:HD21	9	4.93
(4,42)	1:6:A:ARG:HH11	1:7:A:LEU:HD22	9	4.93
(4,42)	1:6:A:ARG:HH11	1:7:A:LEU:HD23	9	4.93
(2,443)	1:6:A:ARG:HH11	1:7:A:LEU:HD21	9	4.93
(2,443)	1:6:A:ARG:HH11	1:7:A:LEU:HD22	9	4.93
(2,443)	1:6:A:ARG:HH11	1:7:A:LEU:HD23	9	4.93
(4,59)	1:3:A:GLU:HG3	1:7:A:LEU:H	3	4.68
(2,535)	1:3:A:GLU:HG3	1:7:A:LEU:H	3	4.68

Continued on next page...

Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(4,42)	1:6:A:ARG:HH11	1:7:A:LEU:HD21	6	4.65
(4,42)	1:6:A:ARG:HH11	1:7:A:LEU:HD22	6	4.65
(4,42)	1:6:A:ARG:HH11	1:7:A:LEU:HD23	6	4.65
(2,443)	1:6:A:ARG:HH11	1:7:A:LEU:HD21	6	4.65
(2,443)	1:6:A:ARG:HH11	1:7:A:LEU:HD22	6	4.65
(2,443)	1:6:A:ARG:HH11	1:7:A:LEU:HD23	6	4.65
(4,59)	1:3:A:GLU:HG3	1:7:A:LEU:H	10	4.61
(2,535)	1:3:A:GLU:HG3	1:7:A:LEU:H	10	4.61
(4,59)	1:3:A:GLU:HG3	1:7:A:LEU:H	4	4.52
(2,535)	1:3:A:GLU:HG3	1:7:A:LEU:H	4	4.52

10 Dihedral-angle violation analysis

No dihedral-angle restraints found