



Full wwPDB X-ray Structure Validation Report i

Aug 26, 2023 – 05:30 PM EDT

PDB ID : 3GFK
Title : Crystal structure of Bacillus subtilis Spx/RNA polymerase alpha subunit C-terminal domain complex
Authors : Lamour, V.; Westblade, L.F.; Campbell, E.A.; Darst, S.A.
Deposited on : 2009-02-27
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see references ①) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	A	131	56%	34%	• 10%		
2	B	79	56%	34%	• • .		

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1843 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Regulatory protein spx.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	118	Total	C 979	N 618	O 174	S 182	5	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	76	Total	C 607	N 373	O 108	S 121	5	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	236	GLY	-	expression tag	UNP P20429
B	237	PRO	-	expression tag	UNP P20429
B	238	HIS	-	expression tag	UNP P20429
B	239	MET	-	expression tag	UNP P20429


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	149	Total O 149 149	0	0
3	B	108	Total O 108 108	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Regulatory protein spx



- Molecule 2: DNA-directed RNA polymerase subunit alpha



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	41.90 Å 44.30 Å 117.20 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 39.45 – 0.93	Depositor EDS
% Data completeness (in resolution range)	98.9 (20.00-2.30) 7.6 (39.45-0.93)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	CNS	Depositor
R , R_{free}	0.229 , 0.282 0.285 , 0.243	Depositor DCC
R_{free} test set	569 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	(Not available)	Xtriage
Anisotropy	(Not available)	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 45.7	EDS
L-test for twinning ¹	$\langle L \rangle =$ (Not available), $\langle L^2 \rangle =$ (Not available)	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.77	EDS
Total number of atoms	1843	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *(Not available)*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/996	0.62	0/1344
2	B	0.40	0/609	0.86	2/809 (0.2%)
All	All	0.37	0/1605	0.72	2/2153 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	240	LYS	N-CA-C	-9.74	84.71	111.00
2	B	245	LYS	N-CA-C	-7.06	91.93	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	979	0	1001	40	0
2	B	607	0	629	54	0
3	A	149	0	0	19	0
3	B	108	0	0	11	0
All	All	1843	0	1630	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:243:ASP:OD1	2:B:244:GLN:N	1.85	1.09
2:B:238:HIS:CD2	2:B:239:MET:H	1.72	1.06
2:B:238:HIS:CG	2:B:239:MET:H	1.75	1.04
2:B:243:ASP:CG	2:B:244:GLN:H	1.79	0.86
1:A:5:TYR:CE2	1:A:37:LEU:HD21	2.14	0.82
2:B:238:HIS:CG	2:B:239:MET:N	2.46	0.81
1:A:10:CYS:HG	1:A:13:CYS:HG	1.00	0.81
2:B:242:GLU:O	2:B:243:ASP:HB3	1.83	0.77
2:B:240:LYS:O	2:B:242:GLU:OE1	2.01	0.77
2:B:251:MET:HE3	2:B:306:LEU:HD13	1.68	0.75
2:B:236:GLY:N	2:B:237:PRO:HD3	2.00	0.75
2:B:238:HIS:CD2	2:B:239:MET:N	2.54	0.75
2:B:294:LYS:HG3	2:B:295:SER:N	2.05	0.72
2:B:244:GLN:H	2:B:244:GLN:HE21	1.38	0.72
1:A:1:MET:HA	3:A:189:HOH:O	1.90	0.72
2:B:276:GLU:HB2	3:B:166:HOH:O	1.90	0.71
2:B:239:MET:HG3	2:B:241:GLU:CD	2.13	0.69
2:B:243:ASP:CG	2:B:244:GLN:HE21	1.94	0.69
1:A:49:THR:HB	3:A:274:HOH:O	1.91	0.69
1:A:95:ILE:HD12	1:A:102:GLN:HG2	1.74	0.69
2:B:244:GLN:N	2:B:244:GLN:HE21	1.90	0.68
1:A:4:LEU:HD23	3:A:219:HOH:O	1.92	0.67
2:B:242:GLU:OE1	2:B:242:GLU:N	2.27	0.67
2:B:282:GLU:HA	3:B:100:HOH:O	1.93	0.67
2:B:248:VAL:O	2:B:251:MET:HE2	1.95	0.66
2:B:239:MET:HA	2:B:241:GLU:HG3	1.79	0.65
1:A:72:GLU:N	3:A:199:HOH:O	2.28	0.64
2:B:236:GLY:N	2:B:237:PRO:CD	2.61	0.63
1:A:48:MET:HG3	1:A:96:ILE:HD11	1.80	0.63
1:A:5:TYR:CZ	1:A:37:LEU:HD21	2.34	0.62
2:B:257:ASP:O	2:B:302:LYS:HE3	2.00	0.61
1:A:70:ASN:CG	3:A:199:HOH:O	2.38	0.61
2:B:296:LEU:HD21	3:B:100:HOH:O	2.00	0.61
2:B:239:MET:HG3	2:B:241:GLU:CG	2.31	0.60
2:B:289:ARG:HD2	3:B:161:HOH:O	2.02	0.59
2:B:251:MET:CE	2:B:306:LEU:HD13	2.33	0.59
1:A:55:GLU:HB2	3:A:274:HOH:O	2.02	0.58
1:A:91:ARG:HD3	3:A:192:HOH:O	2.02	0.58
1:A:17:ARG:HG3	3:A:219:HOH:O	2.03	0.58
1:A:63:VAL:HG13	1:A:89:LEU:HD23	1.86	0.58
1:A:100:ARG:HH11	1:A:100:ARG:HB3	1.67	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34:SER:C	1:A:36:PRO:HD3	2.25	0.57
2:B:237:PRO:O	2:B:238:HIS:C	2.43	0.56
2:B:285:MET:HB2	3:B:100:HOH:O	2.06	0.56
1:A:52:GLY:N	3:A:274:HOH:O	2.40	0.55
2:B:286:MET:SD	2:B:293:ARG:HG2	2.47	0.55
2:B:310:LEU:O	2:B:311:ARG:HB3	2.07	0.55
2:B:251:MET:HE3	2:B:306:LEU:CD1	2.36	0.54
2:B:267:LYS:HA	3:B:200:HOH:O	2.08	0.54
2:B:238:HIS:ND1	2:B:239:MET:SD	2.81	0.54
2:B:289:ARG:HB3	3:B:45:HOH:O	2.07	0.54
2:B:244:GLN:N	2:B:244:GLN:NE2	2.55	0.53
1:A:83:ILE:HG23	3:A:171:HOH:O	2.09	0.53
1:A:24:GLU:HA	3:A:159:HOH:O	2.09	0.52
3:A:178:HOH:O	2:B:260:VAL:N	2.36	0.52
2:B:311:ARG:HD3	3:B:190:HOH:O	2.10	0.51
1:A:27:PHE:HE1	3:A:219:HOH:O	1.94	0.50
1:A:7:SER:OG	1:A:92:ARG:NE	2.45	0.50
1:A:62:LYS:O	1:A:66:LYS:HG3	2.11	0.50
1:A:44:GLN:O	1:A:48:MET:HG2	2.11	0.49
2:B:294:LYS:HG2	3:B:20:HOH:O	2.11	0.49
2:B:296:LEU:HD11	3:B:100:HOH:O	2.12	0.49
2:B:243:ASP:CG	2:B:244:GLN:N	2.48	0.48
1:A:58:SER:HB3	1:A:91:ARG:HG2	1.95	0.47
2:B:294:LYS:CG	2:B:295:SER:N	2.77	0.47
1:A:62:LYS:HD2	3:A:196:HOH:O	2.14	0.47
2:B:268:ARG:NH1	3:B:328:HOH:O	2.47	0.47
3:A:178:HOH:O	2:B:259:SER:HB2	2.15	0.47
1:A:1:MET:HB3	3:A:187:HOH:O	2.14	0.46
1:A:37:LEU:HD22	1:A:41:GLU:CB	2.46	0.46
2:B:239:MET:HB3	2:B:241:GLU:H	1.79	0.45
2:B:237:PRO:O	2:B:239:MET:O	2.35	0.45
2:B:239:MET:O	2:B:240:LYS:HB2	2.16	0.45
1:A:35:GLU:N	1:A:36:PRO:HD3	2.31	0.44
1:A:4:LEU:HG	1:A:6:THR:HG23	2.00	0.44
1:A:61:SER:OG	1:A:64:PHE:HB2	2.18	0.43
2:B:239:MET:O	2:B:240:LYS:CB	2.66	0.43
1:A:19:TRP:O	1:A:23:HIS:HD2	2.02	0.43
1:A:48:MET:HB2	1:A:101:LEU:HD22	2.00	0.43
2:B:244:GLN:H	2:B:244:GLN:NE2	2.09	0.42
2:B:247:LYS:HB2	2:B:250:GLU:HG2	2.01	0.42
1:A:72:GLU:HG2	3:A:255:HOH:O	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:239:MET:HB3	2:B:240:LYS:H	1.29	0.42
2:B:244:GLN:NE2	2:B:244:GLN:CA	2.83	0.42
1:A:26:PRO:O	3:A:189:HOH:O	2.22	0.41
1:A:51:ASP:HB3	2:B:268:ARG:CZ	2.50	0.41
2:B:243:ASP:CG	2:B:244:GLN:NE2	2.67	0.41
1:A:14:ARG:HB2	3:A:233:HOH:O	2.21	0.40
1:A:37:LEU:HD22	1:A:41:GLU:HB2	2.04	0.40
1:A:38:SER:OG	1:A:41:GLU:HG3	2.21	0.40
1:A:58:SER:HB3	1:A:91:ARG:CG	2.50	0.40
2:B:240:LYS:C	2:B:242:GLU:OE1	2.59	0.40
1:A:10:CYS:CB	1:A:13:CYS:HG	2.31	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	116/131 (88%)	113 (97%)	3 (3%)	0	100 100
2	B	74/79 (94%)	65 (88%)	5 (7%)	4 (5%)	2 1
All	All	190/210 (90%)	178 (94%)	8 (4%)	4 (2%)	7 5

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	238	HIS
2	B	240	LYS
2	B	237	PRO
2	B	243	ASP

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	113/124 (91%)	110 (97%)	3 (3%)	44 61
2	B	68/71 (96%)	66 (97%)	2 (3%)	42 58
All	All	181/195 (93%)	176 (97%)	5 (3%)	43 60

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	MET
1	A	100	ARG
1	A	118	VAL
2	B	238	HIS
2	B	242	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	HIS
1	A	44	GLN
1	A	70	ASN
2	B	244	GLN
2	B	275	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.