



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 10:02 pm BST

PDB ID : 6GGY
Title : Paenibacillus sp. YM1 laminaribiose phosphorylase with sulphate bound
Authors : Kuhaudomlarp, S.; Walpole, S.; Stevenson, C.E.M.; Nepogodiev, S.A.; Lawson, D.M.; Angulo, J.; Field, R.A.
Deposited on : 2018-05-04
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	913	
1	B	913	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15063 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Laminaribiose phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	908	6976	4421	1211	1321	23	0	7	0
1	B	905	7087	4489	1241	1333	24	0	7	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP D7UT17
A	0	PRO	-	expression tag	UNP D7UT17
B	-1	GLY	-	expression tag	UNP D7UT17
B	0	PRO	-	expression tag	UNP D7UT17

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Cl	0	0
			2	2		
4	A	2	Total	Cl	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	329	Total	O	0	19
			348	348		
5	B	563	Total	O	0	25
			588	588		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Laminaribiose phosphorylase



- Molecule 1: Laminaribiose phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	146.70Å 146.70Å 222.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	103.74 – 1.95 88.76 – 1.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (103.74-1.95) 100.0 (88.76-1.95)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 1.95Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.202 , 0.223 0.208 , 0.230	Depositor DCC
R_{free} test set	8729 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	24.0	Xtrriage
Anisotropy	0.536	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15063	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.94 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2028e-03.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/7142	0.74	4/9710 (0.0%)
1	B	0.58	0/7252	0.76	5/9837 (0.1%)
All	All	0.55	0/14394	0.75	9/19547 (0.0%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	ASP	CB-CG-OD1	7.56	125.11	118.30
1	B	704	ARG	NE-CZ-NH1	7.28	123.94	120.30
1	B	354	ARG	NE-CZ-NH1	5.57	123.09	120.30
1	B	849	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	A	775	ARG	NE-CZ-NH1	5.30	122.95	120.30
1	B	301	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	B	470	ARG	NE-CZ-NH1	5.22	122.91	120.30
1	A	775	ARG	NE-CZ-NH2	-5.19	117.70	120.30
1	A	488	ASP	CB-CG-OD1	5.08	122.87	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6976	0	6524	19	0
1	B	7087	0	6786	12	0
2	A	10	0	0	1	0
2	B	10	0	0	0	0
3	A	16	0	24	0	0
3	B	24	0	36	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	348	0	0	0	0
5	B	588	0	0	1	0
All	All	15063	0	13370	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:404:MET:HE1	1:A:460:TYR:CE2	2.32	0.63
1:B:557:GLU:HG2	1:B:817:PHE:CZ	2.44	0.51
1:B:311[B]:ARG:NH2	5:B:1553[B]:HOH:O	2.44	0.49
1:A:877:CYS:HB3	1:A:884:CYS:HB2	1.95	0.49
1:A:186:ILE:HG21	1:A:230:PRO:HB2	1.97	0.47
1:A:551:LEU:HB3	1:A:623:LEU:HD13	1.96	0.47
1:B:186:ILE:HG21	1:B:230:PRO:HB2	1.97	0.47
1:B:864:ARG:NH1	1:B:866:ASP:OD1	2.49	0.46
1:A:865:VAL:O	1:A:865:VAL:HG23	2.15	0.46
1:A:754:VAL:HG21	1:A:860:HIS:CE1	2.52	0.45
1:A:859:GLU:H	1:A:862:GLN:HE21	1.64	0.45
1:A:305:ASP:O	1:A:311[A]:ARG:NE	2.50	0.45
1:A:135:VAL:CG1	1:A:135:VAL:O	2.65	0.45
1:B:551:LEU:HB3	1:B:623:LEU:HD13	1.98	0.44
1:A:263[A]:GLU:HG3	1:B:727:ALA:HB1	1.99	0.44
1:B:20:PRO:HG3	1:B:107:VAL:HG23	2.00	0.43
1:A:20:PRO:HG3	1:A:107:VAL:HG23	1.99	0.43
1:A:497:LEU:HD23	1:A:567:LEU:HD22	2.01	0.43
1:B:754:VAL:HG21	1:B:860:HIS:CE1	2.54	0.43
1:B:436:TRP:N	1:B:437:PRO:CD	2.82	0.43
1:B:497:LEU:HD23	1:B:567:LEU:HD22	2.01	0.42
1:A:436:TRP:N	1:A:437:PRO:CD	2.82	0.42
1:A:404:MET:HE3	1:A:484:LEU:CD2	2.49	0.42
1:A:128:ARG:NH1	2:A:1002:SO4:O3	2.53	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:505[A]:ILE:HG23	1:B:626:LYS:HE2	2.03	0.41
1:A:404:MET:CE	1:A:460:TYR:CD2	3.03	0.41
1:A:505:ILE:HG23	1:A:626:LYS:HE2	2.03	0.41
1:A:135:VAL:O	1:A:135:VAL:HG12	2.21	0.40
1:A:404:MET:HE1	1:A:460:TYR:CD2	2.56	0.40
1:B:385:VAL:HG11	1:B:807:LEU:HD23	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	913/913 (100%)	886 (97%)	25 (3%)	2 (0%)	47 38
1	B	910/913 (100%)	881 (97%)	27 (3%)	2 (0%)	47 38
All	All	1823/1826 (100%)	1767 (97%)	52 (3%)	4 (0%)	47 38

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	522	ALA
1	B	522	ALA
1	A	882	ALA
1	B	882	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	682/740 (92%)	677 (99%)	5 (1%)	84	82
1	B	715/740 (97%)	704 (98%)	11 (2%)	65	60
All	All	1397/1480 (94%)	1381 (99%)	16 (1%)	71	71

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	MET
1	A	184	HIS
1	A	462	ARG
1	A	543	PHE
1	A	597	ASP
1	B	4	LYS
1	B	36	MET
1	B	108	GLU
1	B	184	HIS
1	B	471	GLU
1	B	543	PHE
1	B	597	ASP
1	B	750	LYS
1	B	814	LYS
1	B	856	GLN
1	B	884	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	378	GLN
1	A	862	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	1001	-	4,4,4	0.36	0	6,6,6	0.50	0
3	EDO	B	1004	-	3,3,3	0.54	0	2,2,2	0.07	0
3	EDO	A	1006	-	3,3,3	0.45	0	2,2,2	0.47	0
3	EDO	B	1008	-	3,3,3	0.39	0	2,2,2	0.38	0
2	SO4	B	1002	-	4,4,4	0.34	0	6,6,6	0.27	0
2	SO4	A	1002	-	4,4,4	0.39	0	6,6,6	0.24	0
3	EDO	A	1005	-	3,3,3	0.59	0	2,2,2	0.43	0
3	EDO	B	1005	-	3,3,3	0.46	0	2,2,2	0.57	0
3	EDO	A	1004	-	3,3,3	0.48	0	2,2,2	0.52	0
2	SO4	A	1001	-	4,4,4	0.35	0	6,6,6	0.43	0
3	EDO	B	1003	-	3,3,3	0.63	0	2,2,2	0.25	0
3	EDO	B	1006	-	3,3,3	0.48	0	2,2,2	0.41	0
3	EDO	A	1003	-	3,3,3	0.55	0	2,2,2	0.09	0
3	EDO	B	1007	-	3,3,3	0.58	0	2,2,2	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	1004	-	-	0/1/1/1	-
3	EDO	A	1006	-	-	0/1/1/1	-
3	EDO	B	1008	-	-	0/1/1/1	-
3	EDO	A	1005	-	-	1/1/1/1	-
3	EDO	B	1005	-	-	0/1/1/1	-
3	EDO	A	1004	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	1003	-	-	0/1/1/1	-
3	EDO	B	1006	-	-	1/1/1/1	-
3	EDO	A	1003	-	-	0/1/1/1	-
3	EDO	B	1007	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1005	EDO	O1-C1-C2-O2
3	A	1004	EDO	O1-C1-C2-O2
3	B	1007	EDO	O1-C1-C2-O2
3	B	1006	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1002	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	908/913 (99%)	0.64	69 (7%) 13 21	18, 44, 85, 100	0
1	B	905/913 (99%)	0.20	18 (1%) 65 73	15, 26, 63, 92	0
All	All	1813/1826 (99%)	0.42	87 (4%) 30 40	15, 35, 79, 100	0

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	414	ALA	6.6
1	A	881	GLY	5.2
1	B	424	ASN	5.2
1	A	487	ALA	5.2
1	A	563	GLY	5.1
1	A	882	ALA	4.8
1	A	877	CYS	4.7
1	A	872	GLY	4.7
1	A	416	PRO	4.7
1	A	460	TYR	4.6
1	A	558	LEU	4.6
1	A	562	THR	4.5
1	B	881	GLY	4.4
1	A	880	ASP	4.4
1	A	879	ASN	4.4
1	B	880	ASP	4.2
1	A	876	ASP	4.1
1	B	882	ALA	4.0
1	A	419	PHE	3.9
1	B	562	THR	3.9
1	B	416	PRO	3.8
1	A	567	LEU	3.7
1	B	474	ALA	3.7
1	B	879	ASN	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	614	LEU	3.6
1	A	871	ASN	3.5
1	A	868	VAL	3.5
1	A	455	PHE	3.4
1	B	613	LEU	3.4
1	A	309	ALA	3.4
1	A	499	HIS	3.4
1	A	884	CYS	3.3
1	B	414	ALA	3.3
1	A	559	GLN	3.2
1	B	872	GLY	3.2
1	A	442	LEU	3.1
1	A	441	THR	3.0
1	A	574	ALA	3.0
1	B	479	GLU	2.9
1	A	911	SER	2.9
1	A	617	ARG	2.9
1	A	396	LEU	2.8
1	B	476	TRP	2.8
1	A	556	LEU	2.7
1	A	613	LEU	2.7
1	A	616	ILE	2.7
1	B	426	ILE	2.7
1	A	476[A]	TRP	2.7
1	A	619	VAL	2.7
1	B	871	ASN	2.7
1	A	449	GLY	2.7
1	A	445	LEU	2.7
1	A	604	THR	2.7
1	A	595	LEU	2.6
1	A	321	ALA	2.6
1	A	505	ILE	2.6
1	A	482	ASN	2.6
1	A	496	ILE	2.6
1	A	497	LEU	2.5
1	A	492	TYR	2.5
1	A	577	LEU	2.5
1	A	584	ILE	2.5
1	A	878	GLN	2.5
1	A	614	LEU	2.4
1	A	817	PHE	2.4
1	A	891	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	457	PRO	2.4
1	A	485	LEU	2.3
1	A	612	LEU	2.3
1	B	477	THR	2.3
1	A	909	ASN	2.2
1	A	462	ARG	2.2
1	A	461	PHE	2.2
1	A	600	TYR	2.2
1	A	580	LEU	2.2
1	A	902	LEU	2.1
1	A	867	SER	2.1
1	A	399	TYR	2.1
1	A	488	ASP	2.1
1	A	620	ALA	2.1
1	A	456	GLN	2.1
1	A	486	THR	2.1
1	A	154	PRO	2.1
1	A	415	GLY	2.0
1	B	563	GLY	2.0
1	A	603	VAL	2.0
1	A	607	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	B	1005	4/4	0.82	0.19	39,40,40,41	0
3	EDO	A	1004	4/4	0.82	0.12	44,44,46,46	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	B	1004	4/4	0.84	0.28	49,49,49,49	0
4	CL	B	1010	1/1	0.88	0.10	65,65,65,65	0
3	EDO	B	1006	4/4	0.89	0.11	45,48,48,50	0
3	EDO	B	1007	4/4	0.90	0.22	38,42,42,45	0
3	EDO	A	1005	4/4	0.91	0.13	35,36,36,36	0
4	CL	A	1008	1/1	0.91	0.09	65,65,65,65	0
2	SO4	A	1002	5/5	0.92	0.18	56,57,60,64	0
3	EDO	B	1008	4/4	0.92	0.18	34,37,39,40	0
4	CL	A	1007	1/1	0.93	0.09	53,53,53,53	0
4	CL	B	1009	1/1	0.94	0.13	54,54,54,54	0
3	EDO	B	1003	4/4	0.94	0.13	22,22,24,24	0
3	EDO	A	1006	4/4	0.95	0.18	35,36,37,39	0
2	SO4	A	1001	5/5	0.96	0.14	32,33,36,37	0
3	EDO	A	1003	4/4	0.98	0.13	23,25,25,26	0
2	SO4	B	1002	5/5	0.98	0.09	35,38,40,41	0
2	SO4	B	1001	5/5	0.98	0.13	24,26,28,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.