



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 11, 2021 – 12:33 PM EDT

PDB ID : 2GHM
Title : Mutated MAP kinase P38 (Mus Musculus) in complex with Inhibitor PG-895449
Authors : Walter, R.L.; Mekel, M.J.; Evdokimov, A.G.; Pokross, M.E.; Maier, J.A.
Deposited on : 2006-03-27
Resolution : 2.35 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2867 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mitogen-activated protein kinase 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	336	2703	1735	463	494	11	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	180	ALA	THR	engineered mutation	UNP P47811
A	182	PHE	TYR	engineered mutation	UNP P47811

- Molecule 2 is 3-(2-CHLOROBENZYL)-1-(2-{{(1S)-2-HYDROXY-1,2-DIMETHYLPROPYL}AMINO}PYRIMIDIN-4-YL)-1-(4-METHOXYPHENYL)UREA (three-letter code: LIC) (formula: C₂₄H₂₈ClN₅O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
2	A	1	33	24	1	5	3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	131	Total 131	O 131	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 14



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.17Å 74.86Å 77.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.35 34.54 – 2.30	Depositor EDS
% Data completeness (in resolution range)	90.6 (50.00-2.35) 81.5 (34.54-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	-2.77 (at 2.29Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.227 , 0.323 0.231 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	43.3	Xtrriage
Anisotropy	0.339	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.022 for -h,l,k	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2867	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LIC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.92	1/2766 (0.0%)	1.01	9/3755 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	213	MET	SD-CE	-5.14	1.49	1.77

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	205	ASP	CB-CG-OD2	7.49	125.04	118.30
1	A	343	ASP	CB-CG-OD2	6.89	124.50	118.30
1	A	88	ASP	CB-CG-OD2	6.75	124.37	118.30
1	A	335	ASP	CB-CG-OD2	6.67	124.30	118.30
1	A	125	ASP	CB-CG-OD2	6.39	124.05	118.30
1	A	227	ASP	CB-CG-OD2	5.80	123.52	118.30
1	A	124	ASP	CB-CG-OD2	5.70	123.43	118.30
1	A	324	ASP	CB-CG-OD2	5.68	123.41	118.30
1	A	296	ARG	NE-CZ-NH2	-5.02	117.79	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	167	LEU	Peptide
1	A	168	ASP	Peptide
1	A	194	MET	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2703	0	2689	93	0
2	A	33	0	28	3	0
3	A	131	0	0	30	0
All	All	2867	0	2717	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:LEU:HB3	3:A:478:HOH:O	1.59	1.00
1:A:131:ILE:HG21	1:A:213:MET:CE	1.98	0.94
1:A:131:ILE:HG21	1:A:213:MET:HE3	1.56	0.88
1:A:80:HIS:HD2	1:A:82:ASN:H	1.25	0.83
1:A:77:HIS:CE1	1:A:323:TYR:HB2	2.20	0.76
1:A:64:HIS:CD2	3:A:395:HOH:O	2.42	0.72
1:A:80:HIS:CD2	1:A:82:ASN:H	2.08	0.71
1:A:59:PHE:O	1:A:334:ILE:HD11	1.89	0.70
1:A:127:VAL:HG13	3:A:469:HOH:O	1.91	0.70
1:A:77:HIS:CD2	1:A:77:HIS:O	2.46	0.69
1:A:296:ARG:NH1	3:A:451:HOH:O	2.26	0.68
1:A:5:ARG:N	3:A:468:HOH:O	2.24	0.68
1:A:296:ARG:CZ	3:A:451:HOH:O	2.42	0.68
1:A:77:HIS:O	1:A:77:HIS:HD2	1.81	0.63
1:A:216:LEU:HB2	3:A:469:HOH:O	1.99	0.63
1:A:148:HIS:ND1	1:A:169:PHE:HA	2.13	0.61
1:A:106:THR:HB	2:A:353:LIC:H312	1.82	0.61
1:A:63:ILE:HB	3:A:446:HOH:O	2.01	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:MET:HE1	1:A:284:LEU:HD23	1.84	0.59
1:A:75:LEU:HB3	1:A:86:LEU:HG	1.84	0.59
1:A:116:ILE:HG22	1:A:116:ILE:O	2.03	0.58
1:A:77:HIS:CD2	3:A:390:HOH:O	2.57	0.58
1:A:192:GLU:OE2	1:A:296:ARG:NH2	2.31	0.58
1:A:123:THR:HG22	3:A:400:HOH:O	2.04	0.57
1:A:150:ASP:OD2	1:A:185:THR:HG23	2.05	0.57
1:A:235:ILE:O	3:A:478:HOH:O	2.18	0.56
1:A:237:ARG:NH2	3:A:384:HOH:O	2.38	0.55
1:A:283:ASP:OD2	1:A:287:LYS:NZ	2.37	0.55
1:A:115:ASN:C	3:A:435:HOH:O	2.45	0.55
1:A:35:TYR:CE1	1:A:171:LEU:HD22	2.41	0.55
1:A:62:ILE:HD11	3:A:366:HOH:O	2.06	0.55
1:A:148:HIS:O	1:A:149:ARG:HB2	2.07	0.55
1:A:157:ALA:HB2	1:A:167:LEU:HD11	1.90	0.54
1:A:214:ALA:O	1:A:218:THR:HB	2.07	0.54
1:A:87:LEU:HD21	1:A:107:HIS:HE1	1.73	0.54
1:A:327:PHE:O	1:A:330:ARG:HB2	2.09	0.52
1:A:127:VAL:CG1	3:A:469:HOH:O	2.52	0.52
1:A:64:HIS:ND1	1:A:67:ARG:NH2	2.57	0.52
1:A:14:ASN:HB3	3:A:463:HOH:O	2.11	0.51
1:A:213:MET:HA	3:A:469:HOH:O	2.10	0.51
1:A:64:HIS:CE1	1:A:67:ARG:NH2	2.79	0.50
1:A:183:VAL:O	1:A:187:TRP:CE2	2.64	0.50
1:A:80:HIS:HD2	1:A:82:ASN:N	2.02	0.50
1:A:218:THR:HG22	1:A:220:ARG:H	1.77	0.50
1:A:126:HIS:NE2	1:A:162:CYS:SG	2.83	0.49
1:A:12:GLU:HB3	3:A:418:HOH:O	2.11	0.49
1:A:199:HIS:CD2	1:A:199:HIS:N	2.80	0.49
1:A:116:ILE:HG23	1:A:122:LEU:HD21	1.95	0.48
1:A:113:LEU:O	1:A:116:ILE:HG12	2.14	0.48
1:A:184:ALA:HB3	3:A:445:HOH:O	2.14	0.48
1:A:310:GLN:H	1:A:310:GLN:CD	2.17	0.48
1:A:148:HIS:ND1	1:A:169:PHE:CA	2.77	0.47
1:A:268:MET:HE2	3:A:430:HOH:O	2.14	0.47
1:A:139:LYS:NZ	1:A:317:GLU:O	2.35	0.47
1:A:131:ILE:CG2	1:A:213:MET:HE3	2.37	0.46
1:A:148:HIS:CD2	1:A:151:LEU:HG	2.51	0.46
1:A:94:ARG:NE	3:A:459:HOH:O	2.49	0.46
1:A:148:HIS:HD2	1:A:151:LEU:N	2.13	0.46
1:A:193:ILE:O	1:A:194:MET:CG	2.64	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:LYS:NZ	3:A:481:HOH:O	2.49	0.46
1:A:64:HIS:CE1	1:A:67:ARG:HH22	2.34	0.46
1:A:116:ILE:HG21	1:A:122:LEU:HG	1.96	0.46
1:A:123:THR:O	1:A:127:VAL:HG23	2.16	0.46
1:A:60:GLN:HB3	1:A:64:HIS:HD2	1.80	0.45
1:A:83:VAL:HG13	1:A:169:PHE:CE2	2.51	0.45
1:A:86:LEU:HD23	2:A:353:LIC:H311	1.97	0.45
1:A:291:LEU:HA	3:A:483:HOH:O	2.16	0.45
1:A:335:ASP:HB2	3:A:440:HOH:O	2.17	0.45
1:A:290:VAL:O	1:A:296:ARG:HD2	2.17	0.45
1:A:277:ALA:O	1:A:278:ASN:C	2.55	0.44
1:A:77:HIS:ND1	1:A:323:TYR:HB2	2.33	0.44
1:A:195:LEU:HG	1:A:232:LEU:HD11	1.99	0.44
1:A:23:ARG:HD3	1:A:24:TYR:CZ	2.53	0.44
1:A:168:ASP:OD1	1:A:168:ASP:N	2.50	0.44
1:A:235:ILE:HA	3:A:478:HOH:O	2.17	0.43
1:A:38:VAL:HG22	1:A:53:LYS:HB2	2.00	0.43
1:A:283:ASP:OD1	1:A:287:LYS:NZ	2.52	0.43
1:A:289:LEU:HD23	1:A:289:LEU:HA	1.75	0.43
1:A:5:ARG:CA	3:A:468:HOH:O	2.66	0.42
1:A:189:ARG:HD3	3:A:465:HOH:O	2.20	0.42
1:A:74:LEU:O	1:A:78:MET:HG2	2.20	0.42
1:A:270:PHE:CD1	1:A:286:GLU:HG3	2.55	0.42
1:A:158:VAL:CG2	3:A:414:HOH:O	2.67	0.42
1:A:203:THR:HG21	3:A:398:HOH:O	2.19	0.42
1:A:34:ALA:N	3:A:370:HOH:O	2.52	0.41
1:A:183:VAL:O	1:A:187:TRP:CD2	2.74	0.41
1:A:77:HIS:CD2	1:A:77:HIS:C	2.93	0.41
1:A:30:VAL:HG11	2:A:353:LIC:H453	2.02	0.41
1:A:194:MET:CE	1:A:231:GLN:HG2	2.51	0.41
1:A:138:LEU:HD12	1:A:138:LEU:HA	1.91	0.41
1:A:169:PHE:CD2	1:A:169:PHE:N	2.87	0.41
1:A:148:HIS:CE1	1:A:169:PHE:N	2.90	0.40
1:A:80:HIS:CD2	1:A:81:GLU:N	2.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	330/348 (95%)	304 (92%)	19 (6%)	7 (2%)	7 4

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	169	PHE
1	A	168	ASP
1	A	194	MET
1	A	56	SER
1	A	15	LYS
1	A	26	ASN
1	A	116	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	294/306 (96%)	252 (86%)	42 (14%)	3 3

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ARG
1	A	13	LEU
1	A	15	LYS
1	A	27	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	28	SER
1	A	30	VAL
1	A	62	ILE
1	A	72	LEU
1	A	74	LEU
1	A	84	ILE
1	A	86	LEU
1	A	94	ARG
1	A	96	LEU
1	A	97	GLU
1	A	100	ASN
1	A	107	HIS
1	A	114	ASN
1	A	117	VAL
1	A	121	LYS
1	A	163	GLU
1	A	168	ASP
1	A	189	ARG
1	A	192	GLU
1	A	194	MET
1	A	199	HIS
1	A	200	TYR
1	A	218	THR
1	A	220	ARG
1	A	221	THR
1	A	227	ASP
1	A	228	HIS
1	A	232	LEU
1	A	241	THR
1	A	245	GLU
1	A	256	ARG
1	A	257	ASN
1	A	259	ILE
1	A	264	GLN
1	A	275	ILE
1	A	310	GLN
1	A	333	LEU
1	A	339	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	HIS
1	A	77	HIS
1	A	80	HIS
1	A	107	HIS
1	A	199	HIS
1	A	231	GLN
1	A	257	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LIC	A	353	-	33,35,35	0.88	2 (6%)	43,49,49	2.38	10 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LIC	A	353	-	-	3/29/29/29	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	353	LIC	C2-N1	-3.26	1.39	1.44
2	A	353	LIC	C12-CL16	2.61	1.79	1.73

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	353	LIC	N39-C41-N42	-7.59	119.36	126.55
2	A	353	LIC	C37-N39-C41	7.51	122.11	115.45
2	A	353	LIC	C41-N36-C43	-5.59	119.94	124.53
2	A	353	LIC	N36-C41-N42	4.34	123.70	117.18
2	A	353	LIC	C35-C37-N39	-3.88	119.14	123.96
2	A	353	LIC	C35-C3-N42	-3.05	118.55	123.53
2	A	353	LIC	C22-C2-N1	-2.91	116.61	119.74
2	A	353	LIC	C9-C7-N5	-2.47	107.83	113.03
2	A	353	LIC	C51-C47-C43	2.36	114.90	110.95
2	A	353	LIC	C41-N42-C3	2.13	121.68	114.64

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	353	LIC	C25-C21-O30-C31
2	A	353	LIC	C24-C21-O30-C31
2	A	353	LIC	N42-C3-N1-C1


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	353	LIC	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	336/348 (96%)	0.14	11 (3%) 46 59	28, 45, 70, 85	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	119	CYS	11.7
1	A	198	MET	4.9
1	A	263	ALA	3.5
1	A	197	TRP	3.5
1	A	183	VAL	3.4
1	A	34	ALA	3.4
1	A	171	LEU	3.1
1	A	14	ASN	2.9
1	A	185	THR	2.6
1	A	184	ALA	2.6
1	A	35	TYR	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	LIG	A	353	33/33	0.93	0.14	31,41,50,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.