



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 11:07 pm BST

PDB ID : 5GHR  
Title : DNA replication protein  
Authors : Oyama, T.  
Deposited on : 2016-06-20  
Resolution : 2.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	477	<p>9% 87% 9% .</p>
1	C	477	<p>2% 66% 6% . 28%</p>
2	B	80	<p>4% 55% 13% . 31%</p>
2	D	80	<p>3% 64% 6% 30%</p>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7174 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SsDNA-specific exonuclease.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	457	Total 3556	C 2246	N 627	O 671	S 1	Se 11	0	0	0
1	C	344	Total 2688	C 1694	N 472	O 512	S 1	Se 9	0	0	0

- Molecule 2 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
2	B	55	Total 417	C 270	N 70	O 74	Se 3	0	0	0
2	D	56	Total 428	C 276	N 74	O 75	Se 3	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	109	MSE	-	expression tag	UNP Q5JF31
B	110	GLY	-	expression tag	UNP Q5JF31
B	111	SER	-	expression tag	UNP Q5JF31
B	112	SER	-	expression tag	UNP Q5JF31
B	113	HIS	-	expression tag	UNP Q5JF31
B	114	HIS	-	expression tag	UNP Q5JF31
B	115	HIS	-	expression tag	UNP Q5JF31
B	116	HIS	-	expression tag	UNP Q5JF31
B	117	HIS	-	expression tag	UNP Q5JF31
B	118	HIS	-	expression tag	UNP Q5JF31
B	119	SER	-	expression tag	UNP Q5JF31
B	120	SER	-	expression tag	UNP Q5JF31
B	121	GLY	-	expression tag	UNP Q5JF31
B	122	GLU	-	expression tag	UNP Q5JF31
B	123	ASN	-	expression tag	UNP Q5JF31
B	124	LEU	-	expression tag	UNP Q5JF31

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	125	TYR	-	expression tag	UNP Q5JF31
B	126	PHE	-	expression tag	UNP Q5JF31
B	127	GLN	-	expression tag	UNP Q5JF31
B	128	GLY	-	expression tag	UNP Q5JF31
B	129	HIS	-	expression tag	UNP Q5JF31
B	130	MSE	-	expression tag	UNP Q5JF31
D	109	MSE	-	expression tag	UNP Q5JF31
D	110	GLY	-	expression tag	UNP Q5JF31
D	111	SER	-	expression tag	UNP Q5JF31
D	112	SER	-	expression tag	UNP Q5JF31
D	113	HIS	-	expression tag	UNP Q5JF31
D	114	HIS	-	expression tag	UNP Q5JF31
D	115	HIS	-	expression tag	UNP Q5JF31
D	116	HIS	-	expression tag	UNP Q5JF31
D	117	HIS	-	expression tag	UNP Q5JF31
D	118	HIS	-	expression tag	UNP Q5JF31
D	119	SER	-	expression tag	UNP Q5JF31
D	120	SER	-	expression tag	UNP Q5JF31
D	121	GLY	-	expression tag	UNP Q5JF31
D	122	GLU	-	expression tag	UNP Q5JF31
D	123	ASN	-	expression tag	UNP Q5JF31
D	124	LEU	-	expression tag	UNP Q5JF31
D	125	TYR	-	expression tag	UNP Q5JF31
D	126	PHE	-	expression tag	UNP Q5JF31
D	127	GLN	-	expression tag	UNP Q5JF31
D	128	GLY	-	expression tag	UNP Q5JF31
D	129	HIS	-	expression tag	UNP Q5JF31
D	130	MSE	-	expression tag	UNP Q5JF31

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	33	Total	O	0	0
			33	33		
4	B	2	Total	O	0	0
			2	2		
4	C	22	Total	O	0	0
			22	22		
4	D	3	Total	O	0	0
			3	3		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SsDNA-specific exonuclease




- Molecule 1: SsDNA-specific exonuclease



- Molecule 2: Putative uncharacterized protein



- Molecule 2: Putative uncharacterized protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.10Å 116.00Å 234.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.04 – 2.51 25.04 – 2.51	Depositor EDS
% Data completeness (in resolution range)	98.2 (25.04-2.51) 98.6 (25.04-2.51)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.51 (at 2.50Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.212 , 0.249 0.216 , 0.251	Depositor DCC
$R_{free}$ test set	2293 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.3	Xtrriage
Anisotropy	0.905	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 42.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7174	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.19	0/3605	0.35	0/4841
1	C	0.20	0/2725	0.35	0/3662
2	B	0.19	0/421	0.38	0/565
2	D	0.19	0/432	0.37	0/579
All	All	0.20	0/7183	0.35	0/9647

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3556	0	3542	22	0
1	C	2688	0	2668	14	0
2	B	417	0	434	5	0
2	D	428	0	447	4	0
3	A	15	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	1	0
4	A	33	0	0	0	0
4	B	2	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	22	0	0	0	0
4	D	3	0	0	0	0
All	All	7174	0	7091	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:VAL:HG12	1:A:148:MSE:HE2	1.76	0.68
1:A:89:ILE:HD12	1:A:111:LYS:H	1.59	0.67
1:C:239:ASP:HB3	1:C:242:MSE:HE2	1.81	0.62
1:A:50:ARG:NH1	1:A:152:ASN:OD1	2.34	0.61
1:A:349:ASN:HB3	1:A:352:MSE:HE2	1.82	0.61

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	453/477 (95%)	437 (96%)	16 (4%)	0	100	100
1	C	342/477 (72%)	335 (98%)	7 (2%)	0	100	100
2	B	53/80 (66%)	52 (98%)	1 (2%)	0	100	100
2	D	54/80 (68%)	52 (96%)	2 (4%)	0	100	100
All	All	902/1114 (81%)	876 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	363/372 (98%)	354 (98%)	9 (2%)	47 73
1	C	278/372 (75%)	269 (97%)	9 (3%)	39 65
2	B	42/61 (69%)	38 (90%)	4 (10%)	8 17
2	D	43/61 (70%)	42 (98%)	1 (2%)	50 76
All	All	726/866 (84%)	703 (97%)	23 (3%)	39 65

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	177	GLU
2	B	185	ARG
1	C	334	ASP
2	B	184	VAL
1	C	33	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	D	201	-	4,4,4	0.15	0	6,6,6	0.06	0
3	SO4	A	503	-	4,4,4	0.10	0	6,6,6	0.19	0
3	SO4	A	501	-	4,4,4	0.14	0	6,6,6	0.06	0
3	SO4	C	501	-	4,4,4	0.15	0	6,6,6	0.05	0
3	SO4	A	502	-	4,4,4	0.14	0	6,6,6	0.04	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	201	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	446/477 (93%)	0.45	45 (10%) <b>7</b> <b>6</b>	18, 38, 86, 105	0
1	C	335/477 (70%)	0.15	9 (2%) 54 58	18, 34, 72, 94	0
2	B	52/80 (65%)	0.08	3 (5%) <b>23</b> <b>24</b>	22, 34, 50, 66	0
2	D	53/80 (66%)	0.23	2 (3%) 40 43	25, 39, 63, 71	0
All	All	886/1114 (79%)	0.30	59 (6%) <b>17</b> <b>18</b>	18, 36, 82, 105	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	419	GLY	9.7
1	C	342	ALA	9.0
2	D	188	LEU	5.6
1	C	335	TYR	5.5
1	A	442	ILE	5.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	A	503	5/5	0.80	0.35	86,99,111,115	0
3	SO4	A	502	5/5	0.89	0.33	65,75,84,92	0
3	SO4	D	201	5/5	0.90	0.23	56,63,81,83	0
3	SO4	C	501	5/5	0.98	0.16	39,39,47,48	0
3	SO4	A	501	5/5	0.99	0.11	38,38,48,52	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.