

# wwPDB X-ray Structure Validation Summary Report (i)

### Feb 6, 2024 – 11:03 AM EST

PDB ID : 2GJV

Title : Crystal Structure of a Protein of Unknown Function from Salmonella ty-

phimurium

Authors: Minasov, G.; Shuvalova, L.; Brunzelle, J.S.; Vorontsov, I.I.; Collart, F.R.;

Joachimiak, A.; Anderson, W.F.; Midwest Center for Structural Genomics

(MCSG)

Deposited on : 2006-03-31

Resolution : 2.39 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Xtriage (Phenix) : 1.13 EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of o	chain	
1	A	175	65%	12% •	22%
1	В	175	71%	7%	22%
1	С	175	70%	8%	22%
1	D	175	70%	7% •	22%
1	Е	175	71%	6% •	22%



Mol	Chain	Length	Quality of chain		
1	17	1 77	% •		
1	F	175	68%	10%	22%



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6971 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called putative cytoplasmic protein.

Mol	Chain	Residues		Atoms		ZeroOcc	AltConf	Trace		
1	A	136	Total	С	N	О	S	0	0	0
1	Λ	150	1058	663	182	207	6	0	U	
1	В	136	Total	С	N	О	S	0	0	0
1	D	130	1058	663	182	207	6		U	
1	С	136	Total	С	N	О	S	0	0	0
1		130	1058	663	182	207	6	0	U	
1	D	136	Total	С	N	О	S	0	0	0
1	D	130	1058	663	182	207	6	0	U	
1	Е	136	Total	С	N	О	S	0	0	0
1	12	130	1058	663	182	207	6	0	U	0
1	F	136	Total	С	N	О	S	0	0	0
1	Г	190	1058	663	182	207	6	U	0	

There are 144 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	cloning artifact	UNP Q8ZKJ0
A	-22	HIS	-	expression tag	UNP Q8ZKJ0
A	-21	HIS	-	expression tag	UNP Q8ZKJ0
A	-20	HIS	-	expression tag	UNP Q8ZKJ0
A	-19	HIS	-	expression tag	UNP Q8ZKJ0
A	-18	HIS	-	expression tag	UNP Q8ZKJ0
A	-17	HIS	-	expression tag	UNP Q8ZKJ0
A	-16	SER	-	cloning artifact	UNP Q8ZKJ0
A	-15	SER	-	cloning artifact	UNP Q8ZKJ0
A	-14	GLY	-	cloning artifact	UNP Q8ZKJ0
A	-13	VAL	-	cloning artifact	UNP Q8ZKJ0
A	-12	ASP	-	cloning artifact	UNP Q8ZKJ0
A	-11	LEU	-	cloning artifact	UNP Q8ZKJ0
A	-10	GLY	-	cloning artifact	UNP Q8ZKJ0
A	-9	THR	-	cloning artifact	UNP Q8ZKJ0
A	-8	GLU	-	cloning artifact	UNP Q8ZKJ0
A	-7	ASN	-	cloning artifact	UNP Q8ZKJ0



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
A	-6	LEU	-	cloning artifact	UNP Q8ZKJ0
A	-5	TYR	-	cloning artifact	UNP Q8ZKJ0
A	-4	PHE	-	cloning artifact	UNP Q8ZKJ0
A	-3	GLN	-	cloning artifact	UNP Q8ZKJ0
A	-2	SER	-	cloning artifact	UNP Q8ZKJ0
A	-1	ASN	-	cloning artifact	UNP Q8ZKJ0
A	0	ALA	-	cloning artifact	UNP Q8ZKJ0
В	-23	MET	-	cloning artifact	UNP Q8ZKJ0
В	-22	HIS	-	expression tag	UNP Q8ZKJ0
В	-21	HIS	-	expression tag	UNP Q8ZKJ0
В	-20	HIS	-	expression tag	UNP Q8ZKJ0
В	-19	HIS	-	expression tag	UNP Q8ZKJ0
В	-18	HIS	-	expression tag	UNP Q8ZKJ0
В	-17	HIS	-	expression tag	UNP Q8ZKJ0
В	-16	SER	-	cloning artifact	UNP Q8ZKJ0
В	-15	SER	_	cloning artifact	UNP Q8ZKJ0
В	-14	GLY	-	cloning artifact	UNP Q8ZKJ0
В	-13	VAL	-	cloning artifact	UNP Q8ZKJ0
В	-12	ASP	-	cloning artifact	UNP Q8ZKJ0
В	-11	LEU	-	cloning artifact	UNP Q8ZKJ0
В	-10	GLY	_	cloning artifact	UNP Q8ZKJ0
В	-9	THR	-	cloning artifact	UNP Q8ZKJ0
В	-8	GLU	-	cloning artifact	UNP Q8ZKJ0
В	-7	ASN	-	cloning artifact	UNP Q8ZKJ0
В	-6	LEU	-	cloning artifact	UNP Q8ZKJ0
В	-5	TYR	-	cloning artifact	UNP Q8ZKJ0
В	-4	PHE	-	cloning artifact	UNP Q8ZKJ0
В	-3	GLN	-	cloning artifact	UNP Q8ZKJ0
В	-2	SER	-	cloning artifact	UNP Q8ZKJ0
В	-1	ASN	-	cloning artifact	UNP Q8ZKJ0
В	0	ALA	-	cloning artifact	UNP Q8ZKJ0
С	-23	MET	-	cloning artifact	UNP Q8ZKJ0
С	-22	HIS	-	expression tag	UNP Q8ZKJ0
С	-21	HIS	-	expression tag	UNP Q8ZKJ0
С	-20	HIS	_	expression tag	UNP Q8ZKJ0
С	-19	HIS	-	expression tag	UNP Q8ZKJ0
С	-18	HIS	-	expression tag	UNP Q8ZKJ0
С	-17	HIS	-	expression tag	UNP Q8ZKJ0
С	-16	SER	-	cloning artifact	UNP Q8ZKJ0
С	-15	SER	-	cloning artifact	UNP Q8ZKJ0
С	-14	GLY	-	cloning artifact	UNP Q8ZKJ0
С	-13	VAL	-	cloning artifact	UNP Q8ZKJ0



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
С	-12	ASP	-	cloning artifact	UNP Q8ZKJ0
С	-11	LEU	-	cloning artifact	UNP Q8ZKJ0
С	-10	GLY	-	cloning artifact	UNP Q8ZKJ0
С	-9	THR	-	cloning artifact	UNP Q8ZKJ0
С	-8	GLU	-	cloning artifact	UNP Q8ZKJ0
С	-7	ASN	-	cloning artifact	UNP Q8ZKJ0
С	-6	LEU	-	cloning artifact	UNP Q8ZKJ0
С	-5	TYR	-	cloning artifact	UNP Q8ZKJ0
С	-4	PHE	-	cloning artifact	UNP Q8ZKJ0
С	-3	GLN	-	cloning artifact	UNP Q8ZKJ0
С	-2	SER	-	cloning artifact	UNP Q8ZKJ0
С	-1	ASN	-	cloning artifact	UNP Q8ZKJ0
С	0	ALA	-	cloning artifact	UNP Q8ZKJ0
D	-23	MET	-	cloning artifact	UNP Q8ZKJ0
D	-22	HIS	-	expression tag	UNP Q8ZKJ0
D	-21	HIS	-	expression tag	UNP Q8ZKJ0
D	-20	HIS	-	expression tag	UNP Q8ZKJ0
D	-19	HIS	-	expression tag	UNP Q8ZKJ0
D	-18	HIS	-	expression tag	UNP Q8ZKJ0
D	-17	HIS	-	expression tag	UNP Q8ZKJ0
D	-16	SER	-	cloning artifact	UNP Q8ZKJ0
D	-15	SER	_	cloning artifact	UNP Q8ZKJ0
D	-14	GLY	-	cloning artifact	UNP Q8ZKJ0
D	-13	VAL	-	cloning artifact	UNP Q8ZKJ0
D	-12	ASP	_	cloning artifact	UNP Q8ZKJ0
D	-11	LEU	-	cloning artifact	UNP Q8ZKJ0
D	-10	GLY	-	cloning artifact	UNP Q8ZKJ0
D	-9	THR	-	cloning artifact	UNP Q8ZKJ0
D	-8	GLU	-	cloning artifact	UNP Q8ZKJ0
D	-7	ASN	-	cloning artifact	UNP Q8ZKJ0
D	-6	LEU	-	cloning artifact	UNP Q8ZKJ0
D	-5	TYR	-	cloning artifact	UNP Q8ZKJ0
D	-4	PHE	-	cloning artifact	UNP Q8ZKJ0
D	-3	GLN	-	cloning artifact	UNP Q8ZKJ0
D	-2	SER	-	cloning artifact	UNP Q8ZKJ0
D	-1	ASN	-	cloning artifact	UNP Q8ZKJ0
D	0	ALA	-	cloning artifact	UNP Q8ZKJ0
Е	-23	MET	-	cloning artifact	UNP Q8ZKJ0
Е	-22	HIS	-	expression tag	UNP Q8ZKJ0
E	-21	HIS	-	expression tag	UNP Q8ZKJ0
Е	-20	HIS	-	expression tag	UNP Q8ZKJ0
E	-19	HIS	-	expression tag	UNP Q8ZKJ0



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
Е	-18	HIS	-	expression tag	UNP Q8ZKJ0
Е	-17	HIS	-	expression tag	UNP Q8ZKJ0
Е	-16	SER	-	cloning artifact	UNP Q8ZKJ0
Е	-15	SER	-	cloning artifact	UNP Q8ZKJ0
Е	-14	GLY	-	cloning artifact	UNP Q8ZKJ0
Е	-13	VAL	-	cloning artifact	UNP Q8ZKJ0
Е	-12	ASP	-	cloning artifact	UNP Q8ZKJ0
Е	-11	LEU	-	cloning artifact	UNP Q8ZKJ0
Е	-10	GLY	-	cloning artifact	UNP Q8ZKJ0
Е	-9	THR	-	cloning artifact	UNP Q8ZKJ0
Е	-8	GLU	-	cloning artifact	UNP Q8ZKJ0
Е	-7	ASN	-	cloning artifact	UNP Q8ZKJ0
Е	-6	LEU	_	cloning artifact	UNP Q8ZKJ0
Е	-5	TYR	-	cloning artifact	UNP Q8ZKJ0
Е	-4	PHE	-	cloning artifact	UNP Q8ZKJ0
Е	-3	GLN	-	cloning artifact	UNP Q8ZKJ0
Е	-2	SER	-	cloning artifact	UNP Q8ZKJ0
Е	-1	ASN	_	cloning artifact	UNP Q8ZKJ0
Е	0	ALA	-	cloning artifact	UNP Q8ZKJ0
F	-23	MET	-	cloning artifact	UNP Q8ZKJ0
F	-22	HIS	-	expression tag	UNP Q8ZKJ0
F	-21	HIS	-	expression tag	UNP Q8ZKJ0
F	-20	HIS	-	expression tag	UNP Q8ZKJ0
F	-19	HIS	-	expression tag	UNP Q8ZKJ0
F	-18	HIS	-	expression tag	UNP Q8ZKJ0
F	-17	HIS	-	expression tag	UNP Q8ZKJ0
F	-16	SER	-	cloning artifact	UNP Q8ZKJ0
F	-15	SER	-	cloning artifact	UNP Q8ZKJ0
F	-14	GLY	-	cloning artifact	UNP Q8ZKJ0
F	-13	VAL	-	cloning artifact	UNP Q8ZKJ0
F	-12	ASP	-	cloning artifact	UNP Q8ZKJ0
F	-11	LEU	-	cloning artifact	UNP Q8ZKJ0
F	-10	GLY	-	cloning artifact	UNP Q8ZKJ0
F	-9	THR	-	cloning artifact	UNP Q8ZKJ0
F	-8	GLU	-	cloning artifact	UNP Q8ZKJ0
F	-7	ASN	-	cloning artifact	UNP Q8ZKJ0
F	-6	LEU	-	cloning artifact	UNP Q8ZKJ0
F	-5	TYR	-	cloning artifact	UNP Q8ZKJ0
F	-4	PHE	-	cloning artifact	UNP Q8ZKJ0
F	-3	GLN	-	cloning artifact	UNP Q8ZKJ0
F	-2	SER	-	cloning artifact	UNP Q8ZKJ0
F	-1	ASN	-	cloning artifact	UNP Q8ZKJ0



Chain	Residue	Modelled	Actual	Comment	Reference
F	0	ALA	-	cloning artifact	UNP Q8ZKJ0

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	В	1	Total Cl 1 1	0	0
2	С	1	Total Cl 1 1	0	0
2	D	1	Total Cl 1 1	0	0
2	E	1	Total Cl 1 1	0	0
2	F	1	Total Cl 1 1	0	0

• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Ca 2 2	0	0
3	В	1	Total Ca 1 1	0	0
3	С	1	Total Ca 1 1	0	0
3	D	2	Total Ca 2 2	0	0
3	E	1	Total Ca 1 1	0	0
3	F	1	Total Ca 1 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	110	Total O 110 110	0	0
4	В	116	Total O 116 116	0	0




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	92	Total O 92 92	0	0
4	D	108	Total O 108 108	0	0
4	Е	86	Total O 86 86	0	0
4	F	97	Total O 97 97	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: putative cytoplasmic protein






#### SER PRO LLEU LLEU THR ILE VAL ASN GLU GLU

• Molecule 1: putative cytoplasmic protein



• Molecule 1: putative cytoplasmic protein





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	134.14Å 77.00Å 85.11Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $109.25^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	27.90 - 2.39	Depositor
rtesolution (A)	27.90 - 2.39	EDS
% Data completeness	93.5 (27.90-2.39)	Depositor
(in resolution range)	93.5 (27.90-2.39)	EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	7.83  (at  2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
P. P.	0.168 , 0.249	Depositor
$R, R_{free}$	0.167 , $0.248$	DCC
$R_{free}$ test set	1520 reflections $(5.01\%)$	wwPDB-VP
Wilson B-factor $(\mathring{A}^2)$	27.2	Xtriage
Anisotropy	0.270	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.31, 33.2	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6971	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.08% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	Bond lengths		nd angles
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.54	0/1075	0.75	0/1464
1	В	0.51	0/1075	0.75	0/1464
1	С	0.52	0/1075	0.78	1/1464 (0.1%)
1	D	0.50	0/1075	0.75	0/1464
1	Е	0.54	1/1075~(0.1%)	0.76	0/1464
1	F	0.48	0/1075	0.76	0/1464
All	All	0.52	1/6450 (0.0%)	0.76	1/8784 (0.0%)

#### All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}( ext{\AA})$
1	Е	118	CYS	CB-SG	-5.54	1.72	1.81

#### All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	С	25	ILE	N-CA-C	-5.00	97.50	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1058	0	1057	12	0
1	В	1058	0	1057	7	0
1	С	1058	0	1057	7	0
1	D	1058	0	1057	10	0
1	Е	1058	0	1057	9	0
1	F	1058	0	1057	11	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	1	0
3	A	2	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	2	0	0	0	0
3	Ε	1	0	0	0	0
3	F	1	0	0	0	0
4	A	110	0	0	1	0
4	В	116	0	0	0	0
4	С	92	0	0	0	0
4	D	108	0	0	2	0
4	E	86	0	0	0	0
4	F	97	0	0	0	0
All	All	6971	0	6342	47	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 4.

The worst 5 of 47 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} &  ext{Interatomic} \ &  ext{distance} \ &  ext{(Å)} \end{aligned}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:C:62:GLN:HE22	1:D:105:LEU:H	1.23	0.86
1:E:62:GLN:HE22	1:F:105:LEU:H	1.28	0.82
1:D:62:GLN:HE22	1:E:105:LEU:H	1.28	0.77
1:F:14:LEU:HB3	1:F:23:ILE:HD13	1.72	0.71
1:B:88:ARG:HD3	1:B:108:GLU:OE2	1.90	0.71

There are no symmetry-related clashes.



### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	134/175~(77%)	134 (100%)	0	0	100	100
1	В	134/175 (77%)	132 (98%)	2 (2%)	0	100	100
1	С	134/175 (77%)	132 (98%)	1 (1%)	1 (1%)	22	32
1	D	134/175 (77%)	131 (98%)	2 (2%)	1 (1%)	22	32
1	E	134/175 (77%)	132 (98%)	2 (2%)	0	100	100
1	F	134/175 (77%)	129 (96%)	5 (4%)	0	100	100
All	All	804/1050 (77%)	790 (98%)	12 (2%)	2 (0%)	47	62

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	55	GLU
1	С	55	GLU

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	118/155 (76%)	115 (98%)	3 (2%)	47	67
1	В	118/155 (76%)	118 (100%)	0	100	100
1	С	118/155 (76%)	116 (98%)	2 (2%)	60	78
1	D	118/155 (76%)	116 (98%)	2 (2%)	60	78
1	Е	118/155 (76%)	117 (99%)	1 (1%)	81	91



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	F	118/155 (76%)	118 (100%)	0	100	100
All	All	708/930 (76%)	700 (99%)	8 (1%)	73	87

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	88	ARG
1	D	86	ARG
1	С	134	GLN
1	С	-1	ASN
1	D	13	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	62	GLN
1	Е	116	ASN
1	F	134	GLN
1	Е	134	GLN
1	Е	76	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

# 5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.



There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(A^2)$	Q < 0.9
1	A	$136/175 \ (77\%)$	-0.25	1 (0%) 87 86	21, 28, 35, 39	0
1	В	136/175 (77%)	-0.24	2 (1%) 73 72	22, 28, 35, 38	0
1	С	136/175 (77%)	-0.22	1 (0%) 87 86	20, 28, 35, 36	0
1	D	136/175 (77%)	-0.19	4 (2%) 51 50	21, 29, 34, 36	0
1	E	136/175 (77%)	-0.06	7 (5%) 28 26	21, 28, 35, 40	0
1	F	136/175 (77%)	-0.09	2 (1%) 73 72	21, 28, 35, 38	0
All	All	816/1050 (77%)	-0.18	17 (2%) 63 61	20, 28, 35, 40	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	56	ASN	3.8
1	Е	58	ASP	3.6
1	Е	38	GLN	3.6
1	Е	134	GLN	3.0
1	A	134	GLN	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	CA	D	154	1/1	0.90	0.17	56,56,56,56	1
3	CA	В	153	1/1	0.91	0.22	31,31,31,31	1
3	CA	F	153	1/1	0.91	0.19	29,29,29,29	1
3	CA	A	154	1/1	0.94	0.12	44,44,44	1
3	CA	С	153	1/1	0.94	0.12	27,27,27,27	1
2	CL	Ε	152	1/1	0.96	0.20	38,38,38,38	0
2	CL	F	152	1/1	0.96	0.24	54,54,54,54	0
3	CA	A	153	1/1	0.96	0.08	32,32,32,32	1
2	CL	D	152	1/1	0.96	0.25	48,48,48,48	0
2	CL	A	152	1/1	0.97	0.10	40,40,40,40	0
2	CL	В	152	1/1	0.98	0.25	44,44,44,44	0
3	CA	Е	153	1/1	0.98	0.12	48,48,48,48	0
3	CA	D	153	1/1	0.98	0.08	58,58,58,58	0
2	CL	С	152	1/1	0.99	0.15	35,35,35,35	0

# 6.5 Other polymers (i)

There are no such residues in this entry.

