



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 28, 2023 – 04:42 PM EST

PDB ID : 7GKM  
Title : Group deposition SARS-CoV-2 main protease in complex with inhibitors from the COVID Moonshot – Crystal Structure of SARS-CoV-2 main protease in complex with ALP-POS-e0fe77e5-13 (Mpro-P0950)  
Authors : Fearon, D.; Aimon, A.; Aschenbrenner, J.C.; Balcomb, B.H.; Bertram, F.K.R.; Brandao-Neto, J.; Dias, A.; Douangamath, A.; Dunnett, L.; Godoy, A.S.; Gorrie-Stone, T.J.; Koekemoer, L.; Krojer, T.; Lithgo, R.M.; Lukacik, P.; Marples, P.G.; Mikolajek, H.; Nelson, E.; Owen, C.D.; Powell, A.J.; Rangel, V.L.; Skyner, R.; Strain-Damerell, C.M.; Thompson, W.; Tomlinson, C.W.E.; Wild, C.; Walsh, M.A.; von Delft, F.  
Deposited on : 2023-08-11  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	
1	B	306	

Refmac : 5.8.0158  
 CCP4 : 7.0.044 (Gargrove)  
 Ideal geometry (proteins) : Engh & Huber (2001)  
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.36

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	302	Total 2333	C 1476	N 397	O 438	S 22	0	0	0
1	B	299	Total 2312	C 1467	N 394	O 430	S 21	0	0	0

- Molecule 2 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
2	A	1	Total 4	C 2	O 1	S 1	0	0
2	A	1	Total 4	C 2	O 1	S 1	0	0
2	A	1	Total 4	C 2	O 1	S 1	0	0
2	A	1	Total 4	C 2	O 1	S 1	0	0

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	S	0	0
			4	2	1	1		
2	A	1	Total	C	O	S	0	0
			4	2	1	1		
2	A	1	Total	C	O	S	0	0
			4	2	1	1		
2	B	1	Total	C	O	S	0	0
			4	2	1	1		
2	B	1	Total	C	O	S	0	0
			4	2	1	1		
2	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is (3R)-6'-chloro-1-(isoquinolin-4-yl)-2',3'-dihydro-1'H-spiro[piperidine-3,4'-quinolin]-2-one (three-letter code: QY6) (formula: C<sub>22</sub>H<sub>20</sub>ClN<sub>3</sub>O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			27	22	1	3	1		
4	B	1	Total	C	Cl	N	O	0	0
			27	22	1	3	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	208	Total	O	0	0
			208	208		
5	B	133	Total	O	0	0
			133	133		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3C-like proteinase



- Molecule 1: 3C-like proteinase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.26Å 99.12Å 103.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.22 – 1.95 37.22 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.22-1.95) 99.8 (37.22-1.95)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 1.95Å)	Xtrriage
Refinement program	BUSTER 2.10.4 (20-APR-2021)	Depositor
R, $R_{free}$	0.232 , 0.280 0.223 , 0.265	Depositor DCC
$R_{free}$ test set	2407 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.7	Xtrriage
Anisotropy	0.523	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 40.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.012 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5085	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, CL, QY6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/2385	0.62	0/3241
1	B	0.41	0/2364	0.60	0/3212
All	All	0.42	0/4749	0.61	0/6453

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2333	0	2281	19	0
1	B	2312	0	2263	17	0
2	A	28	0	42	3	0
2	B	16	0	24	1	0
3	A	1	0	0	0	0
4	A	27	0	0	2	0
4	B	27	0	0	0	0
5	A	208	0	0	1	0
5	B	133	0	0	1	0
All	All	5085	0	4610	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:ILE:HD11	1:B:54:TYR:HD1	1.65	0.61
1:A:298:ARG:NH1	2:A:403:DMS:O	2.34	0.61
1:B:44:CYS:SG	1:B:54:TYR:CE1	2.97	0.57
1:A:298:ARG:HD3	2:A:403:DMS:C1	2.36	0.55
1:B:298:ARG:HG3	1:B:303:VAL:HB	1.88	0.55
1:A:138:GLY:O	1:B:2:GLY:HA3	2.07	0.55
1:A:298:ARG:HD3	2:A:403:DMS:H11	1.88	0.54
1:A:123:SER:HB2	1:B:303:VAL:O	2.08	0.54
1:A:44:CYS:SG	1:A:54:TYR:CE1	3.01	0.53
1:B:5:LYS:HD3	5:B:506:HOH:O	2.07	0.53
1:B:40:ARG:O	1:B:43:ILE:HG12	2.08	0.53
1:B:298:ARG:HD2	2:B:403:DMS:H12	1.90	0.53
1:A:5:LYS:HD3	5:A:509:HOH:O	2.09	0.51
1:A:86:VAL:HG13	1:A:179:GLY:HA2	1.94	0.48
1:A:165:MET:HB3	4:A:407:QY6:C	2.43	0.48
1:A:95:ASN:HB3	1:A:98:THR:OG1	2.13	0.48
1:B:44:CYS:SG	1:B:54:TYR:HE1	2.37	0.47
1:A:163:HIS:NE2	4:A:407:QY6:N2	2.63	0.46
1:B:95:ASN:HB3	1:B:98:THR:OG1	2.16	0.46
1:B:298:ARG:HD3	1:B:305:PHE:HZ	1.81	0.46
1:A:5:LYS:HG2	1:A:127:GLN:HB3	1.97	0.45
1:A:47:GLU:H	1:A:47:GLU:HG3	1.52	0.44
1:A:233:VAL:HG21	1:A:269:LYS:HE3	1.99	0.44
1:B:113:SER:O	1:B:149:GLY:HA2	2.18	0.44
1:A:113:SER:O	1:A:149:GLY:HA2	2.18	0.44
1:A:109:GLY:HA2	1:A:200:ILE:HD13	2.01	0.43
1:B:233:VAL:HG11	1:B:269:LYS:HG3	2.00	0.43
1:B:109:GLY:HA2	1:B:200:ILE:HD13	1.99	0.43
1:A:186:VAL:HG23	1:A:188:ARG:HB2	2.01	0.42
1:A:123:SER:HB3	1:B:304:THR:HG22	2.02	0.41
1:A:62:SER:OG	1:A:64:HIS:CE1	2.74	0.41
1:B:266:ALA:O	1:B:270:GLU:HG2	2.22	0.40
1:B:164:HIS:CE1	1:B:175:THR:HG23	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	300/306 (98%)	293 (98%)	7 (2%)	0	100	100
1	B	295/306 (96%)	287 (97%)	8 (3%)	0	100	100
All	All	595/612 (97%)	580 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/263 (98%)	253 (98%)	6 (2%)	50	42
1	B	256/263 (97%)	255 (100%)	1 (0%)	91	90
All	All	515/526 (98%)	508 (99%)	7 (1%)	67	62

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	GLU
1	A	153	ASP
1	A	189	GLN
1	A	216	ASP
1	A	235	MET
1	A	263	ASP
1	B	216	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	64	HIS
1	A	164	HIS
1	A	244	GLN
1	B	19	GLN
1	B	164	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 1 is monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	DMS	A	404	-	3,3,3	0.30	0	3,3,3	0.23	0
4	QY6	A	407	-	31,31,31	0.31	0	37,46,46	0.85	1 (2%)
2	DMS	A	408	-	3,3,3	0.30	0	3,3,3	0.36	0
2	DMS	B	401	-	3,3,3	0.28	0	3,3,3	0.23	0
2	DMS	A	409	-	3,3,3	0.30	0	3,3,3	0.26	0
2	DMS	A	402	-	3,3,3	0.30	0	3,3,3	0.43	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DMS	A	403	-	3,3,3	0.25	0	3,3,3	0.86	0
2	DMS	B	403	-	3,3,3	0.04	0	3,3,3	0.32	0
4	QY6	B	405	-	31,31,31	0.30	0	37,46,46	0.95	1 (2%)
2	DMS	A	405	-	3,3,3	0.30	0	3,3,3	0.27	0
2	DMS	B	404	-	3,3,3	0.30	0	3,3,3	0.34	0
2	DMS	B	402	-	3,3,3	0.31	0	3,3,3	0.45	0
2	DMS	A	401	-	3,3,3	0.29	0	3,3,3	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	QY6	A	407	-	-	1/4/34/34	0/5/5/5
4	QY6	B	405	-	-	1/4/34/34	0/5/5/5

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	405	QY6	C9-N1-C11	-4.34	110.10	117.28
4	A	407	QY6	C9-N1-C11	-3.73	111.10	117.28

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	407	QY6	C19-C11-N1-C9
4	B	405	QY6	C19-C11-N1-C9


There are no ring outliers.


3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	407	QY6	2	0
2	A	403	DMS	3	0
2	B	403	DMS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	302/306 (98%)	0.47	22 (7%) 15 23	32, 44, 62, 80	0
1	B	299/306 (97%)	0.62	31 (10%) 6 10	33, 49, 77, 95	0
All	All	601/612 (98%)	0.55	53 (8%) 10 16	32, 45, 69, 95	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	154	TYR	9.0
1	A	302	GLY	7.4
1	B	51	ASN	7.4
1	B	154	TYR	6.0
1	B	60	ARG	5.8
1	B	56	ASP	5.2
1	B	44	CYS	5.1
1	B	189	GLN	4.4
1	B	24	THR	4.2
1	B	57	LEU	3.6
1	B	305	PHE	3.5
1	A	114	VAL	3.4
1	B	232	LEU	3.3
1	B	52	PRO	3.2
1	B	302	GLY	3.2
1	A	72	ASN	3.0
1	B	119	ASN	3.0
1	A	73	VAL	3.0
1	B	235	MET	2.9
1	A	125	VAL	2.8
1	A	115	LEU	2.8
1	A	196	THR	2.8
1	A	112	PHE	2.8
1	B	73	VAL	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	128	CYS	2.8
1	B	76	ARG	2.7
1	B	143	GLY	2.7
1	A	153	ASP	2.7
1	B	59	ILE	2.6
1	B	72	ASN	2.6
1	A	129	ALA	2.5
1	A	126	TYR	2.5
1	B	301	SER	2.5
1	B	153	ASP	2.5
1	B	54	TYR	2.5
1	A	111	THR	2.5
1	A	106	ILE	2.4
1	B	227	LEU	2.4
1	B	86	VAL	2.4
1	B	303	VAL	2.4
1	B	294	PHE	2.3
1	A	7	ALA	2.2
1	B	53	ASN	2.2
1	B	142	ASN	2.2
1	A	150	PHE	2.2
1	A	148	VAL	2.2
1	B	61	LYS	2.2
1	B	278	GLY	2.1
1	B	69	GLN	2.1
1	A	155	ASP	2.1
1	A	161	TYR	2.1
1	A	142	ASN	2.1
1	A	46	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	DMS	B	401	4/4	0.63	0.38	109,109,109,109	0
2	DMS	A	405	4/4	0.80	0.32	98,98,98,98	0
2	DMS	B	404	4/4	0.83	0.23	97,97,97,97	0
2	DMS	A	401	4/4	0.84	0.19	97,97,97,97	0
2	DMS	A	409	4/4	0.84	0.27	115,115,115,115	0
4	QY6	B	405	27/27	0.84	0.18	50,52,52,52	27
2	DMS	A	408	4/4	0.85	0.34	97,97,97,97	0
2	DMS	B	402	4/4	0.88	0.32	78,78,78,78	0
4	QY6	A	407	27/27	0.93	0.11	46,48,51,51	27
2	DMS	A	403	4/4	0.95	0.16	65,65,65,66	0
2	DMS	A	402	4/4	0.96	0.16	80,80,80,80	0
2	DMS	B	403	4/4	0.98	0.13	51,51,51,52	0
2	DMS	A	404	4/4	0.98	0.22	74,74,74,74	0
3	CL	A	406	1/1	0.99	0.13	39,39,39,39	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around QY6 B 405:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around QY6 A 407:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.