



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 23, 2018 – 08:42 PM EDT

PDB ID : 4GLL
Title : Crystal structure of human UDP-xylose synthase.
Authors : Wood, Z.A.; Polizzi, S.J.
Deposited on : 2012-08-14
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : rb-20031021
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20031021


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4155 (2.50-2.50)
Clashscore	122126	4827 (2.50-2.50)
Ramachandran outliers	120053	4735 (2.50-2.50)
Sidechain outliers	120020	4737 (2.50-2.50)
RSRZ outliers	108989	4058 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	 .% 75% 14% • 9%
1	B	337	 4% 74% 15% • 9%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5079 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called UDP-glucuronic acid decarboxylase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	307	Total 2423	C 1537	N 423	O 451	S 12	0	0	0
1	B	306	Total 2414	C 1530	N 422	O 451	S 11	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	84	MET	-	EXPRESSION TAG	UNP Q8NBZ7
B	84	MET	-	EXPRESSION TAG	UNP Q8NBZ7

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 44	C 21	N 7	O 14	P 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	B	1	44	21	7	14	2	0	0

- Molecule 3 is GUANIDINE (three-letter code: GAI) (formula: CH₅N₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
3	A	1	4	1	3	0	0
3	B	1	4	1	3	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	70	Total	O	0	0
			70	70		
5	B	66	Total	O	0	0
			66	66		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UDP-glucuronic acid decarboxylase 1



- Molecule 1: UDP-glucuronic acid decarboxylase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	125.61Å 125.61Å 98.92Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.00 – 2.50 45.02 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.00-2.50) 99.9 (45.02-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.09 (at 2.51Å)	Xtrriage
Refinement program	CNS, REFMAC 5.5.0044	Depositor
R, R_{free}	0.178 , 0.223 0.180 , 0.221	Depositor DCC
R_{free} test set	1589 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtrriage
Anisotropy	0.029	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 64.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.031 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5079	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAD, GAI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/2479	0.66	0/3360
1	B	0.69	0/2467	0.66	0/3341
All	All	0.68	0/4946	0.66	0/6701

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2423	0	2395	37	0
1	B	2414	0	2388	36	0
2	A	44	0	26	0	0
2	B	44	0	26	1	0
3	A	4	0	4	1	0
3	B	4	0	4	1	0
4	A	5	0	0	1	0
4	B	5	0	0	0	0
5	A	70	0	0	1	0
5	B	66	0	0	1	0
All	All	5079	0	4843	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:247:LYS:HE3	1:A:247:LYS:HA	1.19	1.10
1:A:90:ARG:HH11	1:A:90:ARG:HG3	1.27	0.98
1:A:247:LYS:HA	1:A:247:LYS:CE	1.89	0.95
1:B:150:TYR:HE1	1:B:192:ARG:NH1	1.81	0.79
1:A:247:LYS:HE3	1:A:247:LYS:CA	2.06	0.78
1:B:353:GLU:OE2	1:B:353:GLU:N	2.22	0.73
1:A:213:PRO:HB3	1:A:362:LYS:HD2	1.70	0.72
1:B:126:LYS:HE3	1:B:133:ILE:CD1	2.21	0.70
1:B:264:PRO:O	1:B:265:ARG:HB2	1.91	0.70
1:B:150:TYR:HE1	1:B:192:ARG:HH11	1.40	0.70
1:A:90:ARG:NH1	1:A:90:ARG:HG3	2.03	0.70
1:B:263:GLY:O	1:B:266:MET:HG2	1.93	0.69
1:A:169:MET:HE3	1:A:228:ARG:HG3	1.77	0.66
1:B:390:TYR:CE2	1:B:394:GLU:HG3	2.30	0.66
1:A:88:ARG:CZ	1:A:88:ARG:HB3	2.24	0.66
1:B:176:LEU:HG	1:B:230:CYS:HB2	1.78	0.65
1:B:126:LYS:HE3	1:B:133:ILE:HD13	1.79	0.64
1:A:290:TYR:H	3:A:801:GAI:HN22	1.43	0.63
1:B:163:ALA:HB2	2:B:501:NAD:H2D	1.80	0.62
1:B:204:GLU:HG3	1:B:231:TYR:HE2	1.66	0.60
1:A:169:MET:HE3	1:A:228:ARG:CG	2.32	0.59
1:A:274:VAL:HG13	5:A:953:HOH:O	2.01	0.59
1:A:172:PRO:HG2	1:B:248:GLN:HE21	1.68	0.59
1:B:96:GLY:HA3	1:B:117:VAL:HG13	1.84	0.58
1:A:264:PRO:O	1:A:265:ARG:HB2	2.05	0.57
1:B:88:ARG:HH11	1:B:88:ARG:HA	1.70	0.57
1:A:169:MET:CE	1:A:228:ARG:HG3	2.36	0.55
1:A:136:GLU:CD	1:A:136:GLU:H	2.08	0.55
1:A:90:ARG:HH21	1:A:152:GLU:HG2	1.72	0.55
1:B:109:MET:CE	1:B:138:PHE:HB2	2.38	0.54
1:A:287:LEU:N	1:A:287:LEU:HD12	2.23	0.54
1:B:290:TYR:H	3:B:502:GAI:HN21	1.56	0.53
1:B:126:LYS:CE	1:B:133:ILE:CD1	2.88	0.51
1:B:327:GLU:HG3	1:B:329:HIS:CE1	2.46	0.50
1:A:118:VAL:HG21	1:A:149:LEU:HD21	1.93	0.50
1:A:295:GLN:HA	1:A:295:GLN:NE2	2.27	0.49
1:B:390:TYR:CZ	1:B:394:GLU:HG3	2.47	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:164:SER:HA	1:B:165:PRO:HA	1.56	0.48
1:A:127:ARG:H	1:A:127:ARG:NH1	2.11	0.48
1:A:144:ASP:OD1	1:A:146:VAL:HG13	2.13	0.48
1:A:259:PHE:CZ	1:A:363:PRO:HB3	2.49	0.48
1:B:298:ALA:O	1:B:299:PHE:CD2	2.68	0.47
1:B:131:HIS:CG	1:B:131:HIS:O	2.68	0.47
1:A:125:ARG:HB2	1:A:127:ARG:NE	2.30	0.46
1:A:270:ASP:OD1	1:A:275:SER:OG	2.32	0.46
1:A:329:HIS:CD2	1:A:380:LEU:HD22	2.51	0.45
1:A:287:LEU:CD1	1:A:287:LEU:N	2.79	0.45
1:A:89:LYS:HE2	1:A:89:LYS:HB2	1.62	0.45
1:B:191:LYS:C	1:B:191:LYS:HD3	2.37	0.45
1:B:150:TYR:CE1	1:B:192:ARG:NH1	2.72	0.45
1:A:198:LEU:HD23	1:A:198:LEU:C	2.37	0.45
1:A:90:ARG:NH1	1:A:154:ASP:OD2	2.50	0.45
1:A:261:THR:HA	1:A:300:GLN:O	2.16	0.44
1:A:164:SER:HA	1:A:165:PRO:HD3	1.85	0.44
1:B:273:VAL:HG12	1:B:274:VAL:N	2.32	0.43
1:A:162:PRO:HB2	1:A:168:TYR:HB3	2.00	0.43
1:B:348:ILE:HG22	1:B:349:GLN:N	2.34	0.43
1:B:270:ASP:HB3	1:B:272:ARG:H	1.84	0.43
1:A:165:PRO:HB2	1:A:166:PRO:HD3	2.01	0.42
1:B:222:VAL:HG21	1:B:236:ARG:HD2	2.01	0.42
1:A:246:MET:CE	1:A:247:LYS:HD2	2.49	0.42
1:B:289:VAL:CG1	1:B:293:GLY:HA3	2.50	0.42
1:B:88:ARG:HD2	1:B:88:ARG:HA	1.89	0.42
1:B:191:LYS:HD3	1:B:191:LYS:O	2.19	0.42
1:B:279:LEU:HA	1:B:279:LEU:HD23	1.88	0.41
1:B:208:ASP:HB2	5:B:653:HOH:O	2.21	0.41
1:A:203:SER:OG	4:A:802:SO4:O4	2.24	0.41
1:B:94:THR:HB	1:B:160:ALA:HB2	2.02	0.41
1:B:348:ILE:HG22	1:B:349:GLN:H	1.86	0.40
1:A:169:MET:HE3	1:A:228:ARG:NH1	2.35	0.40
1:A:317:VAL:HG21	1:A:371:MET:HB3	2.02	0.40
1:B:109:MET:HE2	1:B:138:PHE:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	305/337 (90%)	296 (97%)	9 (3%)	0	100	100
1	B	302/337 (90%)	287 (95%)	14 (5%)	1 (0%)	43	64
All	All	607/674 (90%)	583 (96%)	23 (4%)	1 (0%)	49	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	343	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/291 (91%)	242 (92%)	22 (8%)	12	23
1	B	263/291 (90%)	250 (95%)	13 (5%)	27	50
All	All	527/582 (90%)	492 (93%)	35 (7%)	18	34

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	88	ARG
1	A	89	LYS
1	A	90	ARG
1	A	126	LYS
1	A	127	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	136	GLU
1	A	204	GLU
1	A	211	VAL
1	A	247	LYS
1	A	260	ASN
1	A	273	VAL
1	A	274	VAL
1	A	275	SER
1	A	282	LEU
1	A	294	SER
1	A	328	GLU
1	A	339	LYS
1	A	346	SER
1	A	360	LYS
1	A	361	ARG
1	A	362	LYS
1	A	393	LYS
1	B	110	MET
1	B	164	SER
1	B	191	LYS
1	B	204	GLU
1	B	273	VAL
1	B	282	LEU
1	B	328	GLU
1	B	337	LEU
1	B	344	SER
1	B	361	ARG
1	B	385	ASN
1	B	393	LYS
1	B	396	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	283	GLN
1	A	295	GLN
1	A	329	HIS
1	B	131	HIS
1	B	135	HIS
1	B	212	HIS
1	B	221	HIS
1	B	248	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	329	HIS
1	B	359	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	A	800	-	40,48,48	1.64	7 (17%)	44,73,73	1.45	6 (13%)
3	GAI	A	801	-	3,3,3	1.52	0	3,3,3	1.08	0
4	SO4	A	802	-	4,4,4	0.29	0	6,6,6	0.18	0
2	NAD	B	501	-	40,48,48	1.68	3 (7%)	44,73,73	1.82	5 (11%)
3	GAI	B	502	-	3,3,3	1.53	0	3,3,3	1.10	0
4	SO4	B	503	-	4,4,4	0.15	0	6,6,6	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '–' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	800	-	-	0/22/62/62	0/5/5/5
3	GAI	A	801	-	-	0/0/0/0	0/0/0/0
4	SO4	A	802	-	-	0/0/0/0	0/0/0/0
2	NAD	B	501	-	-	0/22/62/62	0/5/5/5
3	GAI	B	502	-	-	0/0/0/0	0/0/0/0
4	SO4	B	503	-	-	0/0/0/0	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	800	NAD	C8A-N9A	-4.18	1.31	1.36
2	A	800	NAD	O7N-C7N	-2.82	1.18	1.24
2	A	800	NAD	PA-O2A	-2.73	1.41	1.55
2	A	800	NAD	PN-O2N	-2.47	1.43	1.55
2	A	800	NAD	O4D-C4D	-2.42	1.39	1.45
2	A	800	NAD	C5A-N7A	-2.16	1.32	1.39
2	A	800	NAD	PN-O1N	-2.16	1.43	1.50
2	B	501	NAD	C2A-N1A	2.48	1.38	1.33
2	B	501	NAD	C2A-N3A	4.28	1.39	1.32
2	B	501	NAD	O7N-C7N	8.09	1.40	1.24

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	NAD	N3A-C2A-N1A	-9.89	120.40	128.86
2	A	800	NAD	N3A-C2A-N1A	-5.13	124.47	128.86
2	A	800	NAD	C4A-C5A-N7A	-2.89	106.62	109.41
2	B	501	NAD	PN-O3-PA	-2.51	124.21	132.63
2	B	501	NAD	C4A-C5A-N7A	-2.42	107.07	109.41
2	A	800	NAD	O3B-C3B-C4B	-2.20	104.69	111.06
2	A	800	NAD	C5D-C4D-C3D	-2.13	107.28	115.29
2	A	800	NAD	C5N-C4N-C3N	-2.12	117.86	120.35
2	B	501	NAD	O2N-PN-O1N	2.09	122.78	112.14
2	B	501	NAD	C3N-C7N-N7N	2.72	120.92	117.76
2	A	800	NAD	C3N-C7N-N7N	3.40	121.71	117.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	801	GAI	1	0
4	A	802	SO4	1	0
2	B	501	NAD	1	0
3	B	502	GAI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	307/337 (91%)	-0.17	5 (1%) 72 74	18, 37, 62, 80	6 (1%)
1	B	306/337 (90%)	-0.16	12 (3%) 39 42	17, 36, 65, 94	6 (1%)
All	All	613/674 (90%)	-0.16	17 (2%) 53 56	17, 36, 64, 94	12 (1%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	397	TYR	4.0
1	B	165	PRO	3.4
1	B	396	GLU	3.4
1	B	398	GLN	3.3
1	B	136	GLU	3.3
1	B	88	ARG	3.2
1	B	127	ARG	3.0
1	B	150	TYR	2.7
1	B	248	GLN	2.6
1	B	343	GLY	2.6
1	A	181	ILE	2.5
1	A	247	LYS	2.4
1	A	136	GLU	2.4
1	B	126	LYS	2.2
1	B	110	MET	2.0
1	A	88	ARG	2.0
1	A	110	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GAI	A	801	4/4	0.92	0.19	67,68,69,70	0
3	GAI	B	502	4/4	0.95	0.30	51,51,55,56	0
4	SO4	A	802	5/5	0.96	0.27	77,77,80,82	0
2	NAD	A	800	44/44	0.98	0.11	26,36,40,41	0
2	NAD	B	501	44/44	0.99	0.10	29,38,41,45	0
4	SO4	B	503	5/5	0.99	0.12	53,53,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.