

# Full wwPDB X-ray Structure Validation Report (i)

#### Nov 29, 2023 – 12:20 AM EST

PDB ID : 7GLE

Title : Group deposition SARS-CoV-2 main protease in complex with inhibitors from

the COVID Moonshot – Crystal Structure of SARS-CoV-2 main protease in

complex with MAT-POS-5cd9ea36-13 (Mpro-P1889)

Authors: Fearon, D.; Aimon, A.; Aschenbrenner, J.C.; Balcomb, B.H.; Bertram, F.K.R.;

Brandao-Neto, J.; Dias, A.; Douangamath, A.; Dunnett, L.; Godoy, A.S.; Gorrie-Stone, T.J.; Koekemoer, L.; Krojer, T.; Lithgo, R.M.; Lukacik, P.; Marples, P.G.; Mikolajek, H.; Nelson, E.; Owen, C.D.; Powell, A.J.; Rangel, V.L.; Skyner, R.; Strain-Damerell, C.M.; Thompson, W.; Tomlinson, C.W.E.;

Wild, C.; Walsh, M.A.; von Delft, F.

Deposited on : 2023-08-11

Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)


Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{\rm A})}) \end{array}$
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	306	88%	11%	_			
1	В	306	89%	8%				

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DMS	В	401	-	-	-	X



# 2 Entry composition (i)


There are 5 unique types of molecules in this entry. The entry contains 5022 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 3C-like proteinase.

$\mathbf{Mol}$	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	305	Total 2363	C 1497	N 400	O 444	S 22	0	1	0
1	В	296	Total 2291	C 1455	N 390	O 426	S 20	0	0	0

• Molecule 2 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 4		O 1	S 1	0	0
2	A	1	Total 4		O 1		0	0
2	A	1	Total 4		O 1		0	0
2	A	1	Total 4	$\frac{\mathrm{C}}{2}$	O 1	S 1	0	0

Continued on next page...




n previous	paae
	n previous

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total C O S 4 2 1 1	0	0
2	В	1	Total C O S 4 2 1 1	0	0
2	В	1	Total C O S 4 2 1 1	0	0
2	В	1	Total C O S 4 2 1 1	0	0

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

• Molecule 4 is (4S)-6-chloro-2-[2-(cyclopropylamino)-2-oxoethyl]-N-(isoquinolin-4-yl)-1,2,3,4 -tetrahydroisoquinoline-4-carboxamide (three-letter code: R8O) (formula:  $C_{24}H_{23}ClN_4O_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
4	A	1	Total 31	C 24	Cl 1	N 4	O 2	0	0

• Molecule 5 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	196	Total O 196 196	0	0
5	В	108	Total O 108 108	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3C-like proteinase





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	68.04Å 101.09Å 104.52Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	72.66 - 1.98	Depositor
Resolution (A)	72.66 - 1.98	EDS
% Data completeness	64.4 (72.66-1.98)	Depositor
(in resolution range)	64.3 (72.66-1.98)	EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.61 (at 1.98Å)	Xtriage
Refinement program	BUSTER 2.10.4 (20-APR-2021)	Depositor
D.D.	0.217 , $0.259$	Depositor
$R, R_{free}$	0.211 , $0.251$	DCC
$R_{free}$ test set	1653 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.0	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.31 , 37.4	EDS
L-test for twinning <sup>2</sup>	$< L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	0.001 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5022	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, R8O, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.43	0/2419	0.61	0/3288	
1	В	0.41	0/2342	0.58	0/3181	
All	All	0.42	0/4761	0.59	0/6469	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2363	0	2310	21	0
1	В	2291	0	2245	17	0
2	A	16	0	24	0	0
2	В	16	0	24	1	0
3	A	1	0	0	0	0
4	A	31	0	0	1	0
5	A	196	0	0	0	0
5	В	108	0	0	2	0
All	All	5022	0	4603	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.



All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:298:ARG:HG3	1:B:303:VAL:HB	1.70	0.73
1:A:123:SER:HB2	1:B:303:VAL:O	1.95	0.67
1:A:233:VAL:HG21	1:A:269:LYS:HE3	1.81	0.62
1:A:213:ILE:HA	1:A:304:THR:HG23	1.84	0.60
1:A:44:CYS:SG	1:A:54:TYR:CE1	2.98	0.56
1:A:138:GLY:O	1:B:2:GLY:HA3	2.05	0.56
1:A:121:SER:HB3	1:B:304:THR:HB	1.89	0.54
1:B:109:GLY:HA2	1:B:200:ILE:HD13	1.89	0.53
1:A:163:HIS:NE2	4:A:406:R8O:N3	2.58	0.52
1:A:123:SER:HB3	1:B:304:THR:HG22	1.93	0.51
1:B:5:LYS:HD3	5:B:503:HOH:O	2.10	0.51
1:A:109:GLY:HA2	1:A:200:ILE:HD13	1.92	0.50
1:B:263:ASP:OD1	5:B:501:HOH:O	2.19	0.49
1:B:5:LYS:HG2	1:B:127:GLN:HB3	1.96	0.48
1:B:266:ALA:O	1:B:270:GLU:HG2	2.14	0.47
1:B:111:THR:HG22	1:B:129:ALA:HB2	1.97	0.47
1:A:3:PHE:HE1	1:A:300:CYS:HG	1.63	0.46
1:A:3:PHE:CE1	1:A:300:CYS:SG	3.09	0.46
1:A:5:LYS:HG2	1:A:127:GLN:HB3	1.97	0.45
1:A:95:ASN:HB3	1:A:98:THR:OG1	2.16	0.45
1:B:113:SER:O	1:B:149:GLY:HA2	2.17	0.45
1:B:260:ALA:O	1:B:263:ASP:HB3	2.16	0.45
1:A:163:HIS:CE1	1:A:172:HIS:HB3	2.51	0.45
1:B:95:ASN:HB3	1:B:98:THR:OG1	2.17	0.45
1:A:52:PRO:HD2	1:A:188:ARG:HG3	1.99	0.45
1:A:111:THR:HG22	1:A:129:ALA:HB2	1.99	0.44
1:B:298:ARG:CG	1:B:303:VAL:HB	2.45	0.43
1:A:117:CYS:O	1:A:144:SER:HA	2.20	0.42
1:B:126:TYR:HA	2:B:404:DMS:H22	2.02	0.42
1:B:140:PHE:HD2	1:B:172:HIS:CG	2.38	0.41
1:A:216:ASP:OD1	1:A:218:TRP:HD1	2.02	0.41
1:A:100:LYS:HE2	1:A:100:LYS:HB3	1.89	0.41
1:A:113:SER:O	1:A:149:GLY:HA2	2.21	0.41
1:A:66:PHE:CE1	1:A:87:LEU:HD21	2.56	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	A	304/306~(99%)	298 (98%)	6 (2%)	0	100	100
1	В	292/306~(95%)	284 (97%)	7 (2%)	1 (0%)	41	29
All	All	596/612 (97%)	582 (98%)	13 (2%)	1 (0%)	47	38

#### All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	154	TYR

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Rotameric Outliers	
1	A	263/263 (100%)	258 (98%)	5 (2%)	57 50
1	В	$253/263\ (96\%)$	251 (99%)	2 (1%)	81 80
All	All	516/526 (98%)	509 (99%)	7 (1%)	67 62

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	188	ARG
1	A	189	GLN
1	A	190	THR
1	A	222	ARG
1	A	235	MET

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	В	188	ARG
1	В	216	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	HIS
1	A	164	HIS
1	В	164	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Res	Link	Во	nd leng	ths	В	ond ang	les
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2										
2	DMS	A	401	-	3,3,3	0.33	0	3,3,3	0.75	0										
4	R8O	A	406	-	35,35,35	0.24	0	45,50,50	0.40	0										
2	DMS	В	402	-	3,3,3	0.10	0	3,3,3	0.07	0										
2	DMS	A	403	-	3,3,3	0.33	0	3,3,3	0.16	0										



Mol	Tuno	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DMS	A	402	-	3,3,3	0.30	0	3,3,3	0.54	0
2	DMS	A	404	-	3,3,3	0.32	0	3,3,3	0.18	0
2	DMS	В	401	-	3,3,3	0.33	0	3,3,3	0.29	0
2	DMS	В	403	-	3,3,3	0.28	0	3,3,3	0.30	0
2	DMS	В	404	-	3,3,3	0.23	0	3,3,3	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	$\operatorname{Res}$	Link	Chirals	Torsions	Rings
4	R8O	A	406	-	-	9/16/30/30	0/5/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	406	R8O	C5-C6-N1-C7
4	A	406	R8O	O-C6-N1-C7
4	A	406	R8O	C10-C11-C12-O1
4	A	406	R8O	C6-C5-N-C4
4	A	406	R8O	C14-C13-N2-C12
4	A	406	R8O	C6-C5-N-C10
4	A	406	R8O	C10-C11-C12-N2
4	A	406	R8O	C22-C11-C12-N2
4	A	406	R8O	C21-C13-N2-C12

There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	406	R8O	1	0
2	В	404	DMS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will



also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	# RSRZ > 2	$OWAB(Å^2)$	Q<0.9
1	A	305/306~(99%)	0.13	8 (2%) 56 58	21, 33, 52, 75	0
1	В	296/306~(96%)	0.36	18 (6%) 21 23	21, 40, 65, 88	0
All	All	601/612 (98%)	0.24	26 (4%) 35 37	21, 36, 59, 88	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	304	THR	6.6
1	В	60	ARG	6.1
1	В	305	PHE	5.5
1	A	154	TYR	4.7
1	В	57	LEU	4.7
1	A	301	SER	4.4
1	В	56	ASP	4.2
1	В	154	TYR	4.0
1	В	303	VAL	3.9
1	A	305	PHE	3.9
1	В	59	ILE	3.8
1	В	301	SER	3.5
1	A	302	GLY	3.4
1	A	303	VAL	3.4
1	В	119	ASN	3.0
1	В	189	GLN	3.0
1	В	232	LEU	2.9
1	A	298	ARG	2.9
1	В	294	PHE	2.7
1	В	54	TYR	2.6
1	В	304	THR	2.6
1	A	3	PHE	2.5
1	В	191	ALA	2.4
1	В	58	LEU	2.3

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Res	Type	RSRZ
1	В	64	HIS	2.2
1	В	42	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	DMS	В	401	4/4	0.32	0.57	124,124,124,124	0
2	DMS	В	404	4/4	0.79	0.31	74,74,74,74	0
2	DMS	A	404	4/4	0.82	0.26	66,67,67,67	0
2	DMS	В	403	4/4	0.85	0.21	85,85,85,85	4
4	R8O	A	406	31/31	0.91	0.14	33,37,40,41	31
2	DMS	A	402	4/4	0.92	0.18	65,65,65,66	0
2	DMS	A	401	4/4	0.93	0.23	72,72,72,72	0
2	DMS	A	403	4/4	0.94	0.19	69,69,69,69	0
2	DMS	В	402	4/4	0.98	0.11	47,47,47,47	0
3	CL	A	405	1/1	0.99	0.07	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers (i)

There are no such residues in this entry.

