



# Full wwPDB X-ray Structure Validation Report i

Mar 9, 2024 – 07:40 PM EST

PDB ID : 3GNE  
Title : Crystal structure of alginate lyase vAL-1 from Chlorella virus  
Authors : Ogura, K.; Yamasaki, M.; Hashidume, T.; Yamada, T.; Mikami, B.; Hashimoto, W.; Murata, K.  
Deposited on : 2009-03-17  
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 4787 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called VAL-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	243	Total	C	N	O	S	0	14	0
			1987	1280	338	361	8			
1	B	242	Total	C	N	O	S	0	10	0
			1955	1264	334	351	6			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	245	LEU	-	expression tag	UNP Q9DTZ2
A	246	GLU	-	expression tag	UNP Q9DTZ2
A	247	HIS	-	expression tag	UNP Q9DTZ2
A	248	HIS	-	expression tag	UNP Q9DTZ2
A	249	HIS	-	expression tag	UNP Q9DTZ2
A	250	HIS	-	expression tag	UNP Q9DTZ2
A	251	HIS	-	expression tag	UNP Q9DTZ2
A	252	HIS	-	expression tag	UNP Q9DTZ2
B	245	LEU	-	expression tag	UNP Q9DTZ2
B	246	GLU	-	expression tag	UNP Q9DTZ2
B	247	HIS	-	expression tag	UNP Q9DTZ2
B	248	HIS	-	expression tag	UNP Q9DTZ2
B	249	HIS	-	expression tag	UNP Q9DTZ2
B	250	HIS	-	expression tag	UNP Q9DTZ2
B	251	HIS	-	expression tag	UNP Q9DTZ2
B	252	HIS	-	expression tag	UNP Q9DTZ2

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 13 6 7	0	0
2	B	1	Total C O 13 6 7	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	384	Total O 384 384	0	0
4	B	399	Total O 399 399	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: VAL-1



- Molecule 1: VAL-1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.70Å 72.03Å 60.71Å 90.00° 112.35° 90.00°	Depositor
Resolution (Å)	10.00 – 1.20 20.21 – 1.20	Depositor EDS
% Data completeness (in resolution range)	89.7 (10.00-1.20) 94.3 (20.21-1.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	3.07 (at 1.20Å)	Xtriage
Refinement program	SHELXL-97	Depositor
$R$ , $R_{free}$	0.132 , 0.179 0.129 , 0.125	Depositor DCC
$R_{free}$ test set	7163 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	9.5	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 58.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.489 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	4787	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	0/2066	1.25	15/2783 (0.5%)
1	B	0.71	0/2029	1.22	15/2734 (0.5%)
All	All	0.72	0/4095	1.24	30/5517 (0.5%)

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	167	ARG	CD-NE-CZ	10.37	138.11	123.60
1	A	189	TYR	CB-CG-CD1	9.14	126.49	121.00
1	B	38	ARG	CD-NE-CZ	8.33	135.27	123.60
1	B	201	ARG	NE-CZ-NH2	-8.33	116.14	120.30
1	A	89	ARG	NE-CZ-NH1	-7.45	116.57	120.30
1	A	206[A]	ARG	NE-CZ-NH1	7.39	123.99	120.30
1	A	206[B]	ARG	NE-CZ-NH1	7.39	123.99	120.30
1	B	189	TYR	CB-CG-CD1	7.21	125.32	121.00
1	A	81	TYR	CB-CG-CD1	-6.88	116.87	121.00
1	B	201	ARG	NE-CZ-NH1	6.65	123.62	120.30
1	B	243	TYR	CA-C-O	-5.96	107.58	120.10
1	B	167	ARG	CD-NE-CZ	5.91	131.87	123.60
1	A	106	TYR	CB-CG-CD2	5.89	124.53	121.00
1	A	106	TYR	CG-CD1-CE1	5.69	125.85	121.30
1	B	89	ARG	NE-CZ-NH1	-5.58	117.51	120.30
1	B	80	PHE	CB-CG-CD2	5.37	124.56	120.80
1	A	206[A]	ARG	CD-NE-CZ	5.36	131.11	123.60
1	A	206[B]	ARG	CD-NE-CZ	5.36	131.11	123.60
1	B	45	ASP	CB-CG-OD2	-5.31	113.52	118.30
1	B	106	TYR	CG-CD1-CE1	5.25	125.50	121.30
1	B	206	ARG	NE-CZ-NH1	5.25	122.93	120.30
1	A	206[A]	ARG	NE-CZ-NH2	-5.23	117.68	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	206[B]	ARG	NE-CZ-NH2	-5.23	117.68	120.30
1	B	167	ARG	NE-CZ-NH2	-5.23	117.69	120.30
1	B	87	PHE	CB-CG-CD1	-5.22	117.14	120.80
1	A	80	PHE	CB-CG-CD2	5.22	124.45	120.80
1	A	178	LYS	O-C-N	5.20	131.01	122.70
1	A	44	TYR	CB-CG-CD2	-5.17	117.90	121.00
1	B	189	TYR	CA-CB-CG	5.16	123.20	113.40
1	B	201	ARG	CD-NE-CZ	5.01	130.62	123.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbit. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1987	0	1972	31	0
1	B	1955	0	1953	26	0
2	A	13	0	5	0	0
2	B	13	0	5	0	0
3	A	18	0	24	0	0
3	B	18	0	24	1	0
4	A	384	0	0	6	0
4	B	399	0	0	7	0
All	All	4787	0	3983	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:18:SER:H	3:B:605:GOL:H12	1.21	1.02
1:B:138:LYS:HE2	1:B:184:LEU:HB3	1.47	0.93
1:A:10:ASN:HD21	1:A:38:ARG:HE	1.25	0.79
1:A:138:LYS:HE2	1:A:184:LEU:HB3	1.62	0.79

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:LYS:HE2	1:A:134:ASP:OD2	1.83	0.78
1:B:191:ILE:HD11	1:B:196[B]:LYS:HE2	1.66	0.77
1:B:10:ASN:HD21	1:B:38:ARG:HE	1.33	0.76
1:A:68:LEU:HA	1:A:240[B]:MET:HE2	1.68	0.75
1:A:138:LYS:HE3	1:A:201:ARG:HB3	1.69	0.75
1:B:138:LYS:HE3	1:B:201:ARG:HB3	1.69	0.74
1:A:96[A]:THR:HG22	1:A:115:ASN:HB3	1.73	0.69
1:B:96[A]:THR:HG21	4:B:726:HOH:O	1.92	0.69
1:B:134:ASP:HB2	4:B:1250:HOH:O	1.93	0.68
1:A:96[A]:THR:HG21	1:A:172:THR:OG1	1.98	0.64
1:B:14[B]:LYS:HG3	4:B:942:HOH:O	1.99	0.62
1:B:41:LYS:HZ2	1:B:41:LYS:HB3	1.65	0.62
1:A:184:LEU:H	1:A:203:ASN:HD21	1.47	0.61
1:A:133[A]:SER:HB3	4:A:828:HOH:O	2.00	0.61
1:A:26[B]:MET:HG2	4:A:1200:HOH:O	2.02	0.60
1:B:157[B]:LYS:HE2	4:B:1206:HOH:O	2.02	0.59
1:A:184:LEU:H	1:A:203:ASN:ND2	2.00	0.58
1:A:120:GLN:HE21	1:A:153:GLN:HE21	1.50	0.57
1:A:62[B]:SER:OG	1:A:214:ARG:HD2	2.05	0.57
1:A:10:ASN:ND2	1:A:38:ARG:HE	2.00	0.57
1:A:200:LYS:O	1:A:201:ARG:HB2	2.06	0.55
1:B:120:GLN:HE21	1:B:153:GLN:HE21	1.53	0.55
1:B:184:LEU:H	1:B:203:ASN:HD21	1.56	0.54
1:B:200:LYS:O	1:B:201:ARG:HB2	2.07	0.54
1:A:138:LYS:NZ	1:A:201:ARG:HG2	2.23	0.54
1:B:184:LEU:H	1:B:203:ASN:ND2	2.06	0.54
1:B:26[A]:MET:HG2	4:B:1156:HOH:O	2.10	0.51
1:B:138:LYS:HE3	1:B:201:ARG:CB	2.37	0.51
1:B:28[B]:LYS:HZ3	1:B:43:VAL:HG11	1.76	0.50
1:A:10:ASN:HD21	1:A:38:ARG:NE	2.01	0.50
1:A:50:THR:H	1:A:53:ASN:HD22	1.61	0.49
1:A:73:ILE:HG12	1:A:240[B]:MET:CE	2.43	0.49
1:B:26[A]:MET:HG3	1:B:29:SER:O	2.12	0.49
1:B:14[B]:LYS:HE3	4:B:942:HOH:O	2.12	0.48
1:A:138:LYS:CE	1:A:184:LEU:HB3	2.38	0.48
1:B:10:ASN:ND2	1:B:38:ARG:HE	2.07	0.48
1:B:2:ASN:N	4:B:1157:HOH:O	2.47	0.48
1:A:73:ILE:HG12	1:A:240[B]:MET:HE3	1.95	0.47
1:A:120:GLN:NE2	1:A:153:GLN:HE21	2.13	0.47
1:A:157[A]:LYS:HE2	4:A:810:HOH:O	2.14	0.46
1:A:134:ASP:HB3	4:A:1177:HOH:O	2.14	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26[B]:MET:HG3	1:A:29:SER:O	2.16	0.46
1:A:201:ARG:HG3	4:A:1416:HOH:O	2.16	0.46
1:B:120:GLN:NE2	1:B:153:GLN:HE21	2.15	0.45
1:A:14:LYS:NZ	4:A:1203:HOH:O	2.50	0.45
1:B:200:LYS:HB2	1:B:201:ARG:HH11	1.81	0.45
1:A:184:LEU:O	1:A:201:ARG:HA	2.16	0.45
1:B:41:LYS:HB3	1:B:41:LYS:NZ	2.30	0.45
1:A:134:ASP:N	1:A:134:ASP:OD1	2.48	0.44
1:A:2:ASN:ND2	1:A:244[A]:GLU:OE1	2.50	0.43
1:B:50:THR:H	1:B:53:ASN:HD22	1.68	0.41
1:A:50:THR:N	1:A:53:ASN:HD22	2.19	0.41
1:B:136:LYS:HD3	1:B:136:LYS:HA	1.83	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	254/252 (101%)	245 (96%)	9 (4%)	0	100 100
1	B	250/252 (99%)	241 (96%)	8 (3%)	1 (0%)	34 11
All	All	504/504 (100%)	486 (96%)	17 (3%)	1 (0%)	47 19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	201	ARG

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	214/209 (102%)	211 (99%)	3 (1%)	67 32
1	B	209/209 (100%)	205 (98%)	4 (2%)	57 19
All	All	423/418 (101%)	416 (98%)	7 (2%)	59 24

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	178	LYS
1	A	181	ILE
1	A	201	ARG
1	B	136	LYS
1	B	178	LYS
1	B	201	ARG
1	B	203	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	20	ASN
1	A	53	ASN
1	A	120	GLN
1	A	203	ASN
1	B	10	ASN
1	B	20	ASN
1	B	53	ASN
1	B	120	GLN
1	B	158	ASN
1	B	203	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	GOL	B	605	-	5,5,5	0.35	0	5,5,5	0.29	0
3	GOL	A	602	-	5,5,5	0.39	0	5,5,5	0.89	0
3	GOL	B	601	-	5,5,5	0.64	0	5,5,5	0.57	0
3	GOL	B	606	-	5,5,5	0.56	0	5,5,5	0.67	0
2	FLC	B	502	-	12,12,12	1.08	1 (8%)	17,17,17	1.59	3 (17%)
2	FLC	A	501	-	12,12,12	1.00	1 (8%)	17,17,17	1.64	3 (17%)
3	GOL	A	604	-	5,5,5	0.58	0	5,5,5	1.04	0
3	GOL	A	603	-	5,5,5	0.39	0	5,5,5	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	605	-	-	4/4/4/4	-
3	GOL	A	602	-	-	0/4/4/4	-
3	GOL	B	601	-	-	0/4/4/4	-
3	GOL	B	606	-	-	0/4/4/4	-
2	FLC	B	502	-	-	0/16/16/16	-
2	FLC	A	501	-	-	0/16/16/16	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	604	-	-	0/4/4/4	-
3	GOL	A	603	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	502	FLC	OA2-CAC	-2.70	1.21	1.30
2	A	501	FLC	OA2-CAC	-2.53	1.22	1.30

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FLC	OB1-CBC-CB	-3.89	116.75	122.25
2	B	502	FLC	OB1-CBC-CB	-3.74	116.96	122.25
2	A	501	FLC	OB2-CBC-CB	3.21	118.62	113.05
2	A	501	FLC	OHB-CB-CBC	3.18	113.33	108.86
2	B	502	FLC	OB2-CBC-CB	2.99	118.25	113.05
2	B	502	FLC	OHB-CB-CBC	2.38	112.21	108.86

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	605	GOL	O1-C1-C2-C3
3	B	605	GOL	O1-C1-C2-O2
3	B	605	GOL	C1-C2-C3-O3
3	B	605	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	605	GOL	1	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	243/252 (96%)	-0.78	0 [100] [100]	7, 11, 28, 93	0
1	B	242/252 (96%)	-0.81	0 [100] [100]	7, 11, 24, 90	0
All	All	485/504 (96%)	-0.80	0 [100] [100]	7, 11, 26, 93	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	A	602	6/6	0.94	0.11	13,19,19,20	0
3	GOL	B	605	6/6	0.94	0.13	12,35,45,57	0
3	GOL	A	604	6/6	0.96	0.11	13,30,41,47	0
3	GOL	B	601	6/6	0.96	0.13	15,18,19,23	0
3	GOL	A	603	6/6	0.96	0.12	14,16,20,25	0
3	GOL	B	606	6/6	0.96	0.09	10,18,18,18	0
2	FLC	A	501	13/13	0.99	0.04	8,9,12,14	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FLC	B	502	13/13	0.99	0.04	8,9,12,14	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.