



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 08:22 PM EDT

PDB ID : 6GN8  
Title : Exoenzyme S from *Pseudomonas aeruginosa* in complex with human 14-3-3 protein beta, trimeric crystal form  
Authors : Karlberg, T.; Pinto, A.F.; Hornyak, P.; Thorsell, A.G.; Nareoja, K.; Schuler, H.  
Deposited on : 2018-05-30  
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	86% 9% 5%
1	B	243	5% 85% 7% 7%
2	C	244	8% 77% 8% 13%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5411 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 protein beta/alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	232	1871	1170	314	378	9	0	0	0
1	B	225	1814	1137	306	362	9	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	235	GLU	-	expression tag	UNP P31946
A	236	ASN	-	expression tag	UNP P31946
A	237	LEU	-	expression tag	UNP P31946
A	238	TYR	-	expression tag	UNP P31946
A	239	PHE	-	expression tag	UNP P31946
A	240	GLN	-	expression tag	UNP P31946
A	241	SER	-	expression tag	UNP P31946
A	242	LEU	-	expression tag	UNP P31946
A	243	GLU	-	expression tag	UNP P31946
B	235	GLU	-	expression tag	UNP P31946
B	236	ASN	-	expression tag	UNP P31946
B	237	LEU	-	expression tag	UNP P31946
B	238	TYR	-	expression tag	UNP P31946
B	239	PHE	-	expression tag	UNP P31946
B	240	GLN	-	expression tag	UNP P31946
B	241	SER	-	expression tag	UNP P31946
B	242	LEU	-	expression tag	UNP P31946
B	243	GLU	-	expression tag	UNP P31946

- Molecule 2 is a protein called Exoenzyme S.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	212	1587	978	292	313	4	0	1	0

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	210	MET	-	initiating methionine	UNP Q93SQ1
C	211	GLY	-	expression tag	UNP Q93SQ1
C	212	SER	-	expression tag	UNP Q93SQ1
C	213	SER	-	expression tag	UNP Q93SQ1
C	214	HIS	-	expression tag	UNP Q93SQ1
C	215	HIS	-	expression tag	UNP Q93SQ1
C	216	HIS	-	expression tag	UNP Q93SQ1
C	217	HIS	-	expression tag	UNP Q93SQ1
C	218	HIS	-	expression tag	UNP Q93SQ1
C	219	HIS	-	expression tag	UNP Q93SQ1
C	220	SER	-	expression tag	UNP Q93SQ1
C	221	GLN	-	expression tag	UNP Q93SQ1
C	222	ASP	-	expression tag	UNP Q93SQ1
C	223	PRO	-	expression tag	UNP Q93SQ1
C	224	ASN	-	expression tag	UNP Q93SQ1
C	225	SER	-	expression tag	UNP Q93SQ1
C	226	GLU	-	expression tag	UNP Q93SQ1
C	227	ASN	-	expression tag	UNP Q93SQ1
C	228	LEU	-	expression tag	UNP Q93SQ1
C	229	TYR	-	expression tag	UNP Q93SQ1
C	230	PHE	-	expression tag	UNP Q93SQ1
C	231	GLN	-	expression tag	UNP Q93SQ1
C	232	GLY	-	expression tag	UNP Q93SQ1
C	379	ALA	GLU	engineered mutation	UNP Q93SQ1
C	381	ALA	GLU	engineered mutation	UNP Q93SQ1

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total C O 8 6 2	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	59	Total O 59 59	0	0
4	B	18	Total O 18 18	0	0
4	C	54	Total O 54 54	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 14-3-3 protein beta/alpha

Chain A: 





- Molecule 1: 14-3-3 protein beta/alpha

Chain B: 



- Molecule 2: Exoenzyme S

Chain C: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	159.91Å 59.40Å 120.33Å 90.00° 125.79° 90.00°	Depositor
Resolution (Å)	97.61 – 2.34 97.61 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.7 (97.61-2.34) 99.7 (97.61-2.34)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.76 (at 2.34Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.200 , 0.248 0.202 , 0.253	Depositor DCC
$R_{free}$ test set	1957 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.3	Xtrriage
Anisotropy	0.401	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 46.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5411	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/1898	0.59	0/2555
1	B	0.47	0/1840	0.59	0/2474
2	C	0.53	0/1607	0.73	0/2155
All	All	0.50	0/5345	0.63	0/7184

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1871	0	1848	8	0
1	B	1814	0	1798	10	0
2	C	1587	0	1577	17	0
3	C	8	0	14	0	0
4	A	59	0	0	0	0
4	B	18	0	0	0	0
4	C	54	0	0	0	0
All	All	5411	0	5237	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:447:MET:HA	2:C:448:ARG:HB3	1.34	1.08
2:C:447:MET:HA	2:C:448:ARG:CB	2.15	0.77
2:C:447:MET:CA	2:C:448:ARG:HB3	2.16	0.73
2:C:447:MET:HB3	2:C:450:LEU:HB2	1.71	0.72
1:B:215:ASP:HA	2:C:446:ARG:HG3	1.71	0.70
1:A:115:GLU:HG2	1:A:168:ILE:HD12	1.90	0.53
2:C:448:ARG:HG2	2:C:449:GLY:H	1.75	0.51
2:C:447:MET:CB	2:C:450:LEU:HB2	2.39	0.50
1:B:221:GLN:HG2	1:B:224:ARG:HH12	1.78	0.48
2:C:347:ASN:HB3	2:C:350:VAL:HG22	1.96	0.48
2:C:356:GLN:HB3	2:C:359:ILE:HD11	1.95	0.48
1:A:114:PRO:HG3	1:A:161:GLU:HG2	1.96	0.46
1:B:215:ASP:CA	2:C:446:ARG:HG3	2.44	0.46
2:C:245:PHE:O	2:C:248:ASP:HB2	2.16	0.46
1:A:74:ASN:HB3	1:A:77:LYS:HB2	1.98	0.46
1:B:74:ASN:HB3	1:B:77:LYS:HB2	1.99	0.45
2:C:301:MET:HE3	2:C:341:TYR:CE1	2.52	0.45
1:A:103:LEU:HA	1:A:107:LEU:HB2	1.99	0.45
2:C:371:SER:HB3	2:C:380:LYS:HA	1.98	0.44
1:A:158:SER:HA	1:A:162:MET:HE3	1.98	0.44
2:C:398:ASP:HB3	2:C:400:GLN:H	1.83	0.44
2:C:327:PHE:CE1	2:C:390:MET:HB2	2.52	0.44
1:B:89:GLU:HG2	1:B:134:VAL:HB	1.99	0.44
1:B:103:LEU:HA	1:B:107:LEU:HB2	2.00	0.44
1:A:89:GLU:HG2	1:A:134:VAL:HB	2.01	0.42
2:C:448:ARG:HG2	2:C:449:GLY:N	2.35	0.42
1:A:64:ILE:HD11	1:B:18:ALA:HB2	2.02	0.41
1:B:68:GLU:OE1	1:B:85:ARG:HD3	2.19	0.41
1:B:48:VAL:HG21	2:C:448:ARG:HG3	2.03	0.41
1:B:114:PRO:HG3	1:B:161:GLU:HG2	2.02	0.40
1:A:68:GLU:OE1	1:A:85:ARG:HD3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/243 (95%)	227 (99%)	3 (1%)	0	100	100
1	B	221/243 (91%)	215 (97%)	6 (3%)	0	100	100
2	C	209/244 (86%)	193 (92%)	12 (6%)	4 (2%)	8	5
All	All	660/730 (90%)	635 (96%)	21 (3%)	4 (1%)	25	26

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	259	GLY
2	C	447	MET
2	C	448	ARG
2	C	275	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/217 (95%)	197 (96%)	9 (4%)	28	35
1	B	199/217 (92%)	194 (98%)	5 (2%)	47	58
2	C	160/189 (85%)	153 (96%)	7 (4%)	28	35
All	All	565/623 (91%)	544 (96%)	21 (4%)	34	43

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	SER
1	A	78	GLN
1	A	93	GLN
1	A	99	VAL
1	A	138	ASP
1	A	156	GLU
1	A	161	GLU
1	A	176	PHE
1	A	211	GLU
1	B	99	VAL
1	B	161	GLU
1	B	162	MET
1	B	176	PHE
1	B	218	LEU
2	C	345	SER
2	C	352	ARG
2	C	356	GLN
2	C	385	ASN
2	C	446	ARG
2	C	447	MET
2	C	450	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	GLN
1	B	79	GLN
2	C	385	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	MPD	C	501	-	7,7,7	0.57	0	9,10,10	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	C	501	-	-	0/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	232/243 (95%)	0.02	1 (0%) 92 96	35, 52, 98, 134	0
1	B	225/243 (92%)	0.40	13 (5%) 23 32	46, 85, 163, 211	0
2	C	212/244 (86%)	0.37	19 (8%) 9 15	37, 58, 119, 162	0
All	All	669/730 (91%)	0.26	33 (4%) 29 40	35, 62, 145, 211	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	205	LEU	7.3
2	C	354	PHE	7.1
2	C	446	ARG	6.1
2	C	353	SER	5.0
2	C	443	VAL	5.0
2	C	355	GLY	4.9
2	C	352	ARG	4.3
1	A	2	THR	3.9
2	C	350	VAL	3.5
1	B	165	THR	3.5
1	B	73	ARG	3.3
2	C	237	LEU	3.2
2	C	445	LEU	3.2
1	B	218	LEU	3.0
2	C	449	GLY	3.0
1	B	232	SER	2.9
1	B	201	ALA	2.8
2	C	448	ARG	2.8
2	C	450	LEU	2.7
2	C	351	ALA	2.5
2	C	322	ARG	2.5
1	B	77	LYS	2.4
2	C	432	PRO	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	108	ILE	2.3
1	B	121	LEU	2.3
2	C	444	ARG	2.3
1	B	219	ILE	2.2
2	C	236	ALA	2.1
1	B	76	LYS	2.1
1	B	74	ASN	2.1
1	B	203	ALA	2.1
2	C	453	ALA	2.1
2	C	258	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MPD	C	501	8/8	0.95	0.28	61,62,67,70	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.