



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 06:33 pm BST

PDB ID : 6GQZ
Title : Petrobactin-binding engineered lipocalin without ligand
Authors : Skerra, A.; Eichinger, A.
Deposited on : 2018-06-08
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 3288 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neutrophil gelatinase-associated lipocalin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	174	1475	953	245	272	5	0	11	0
1	B	174	1474	955	245	269	5	0	9	0

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	HIS	GLN	engineered mutation	UNP P80188
A	36	PHE	LEU	engineered mutation	UNP P80188
A	38	GLU	GLY	engineered mutation	UNP P80188
A	40	ILE	ALA	engineered mutation	UNP P80188
A	41	GLN	ILE	engineered mutation	UNP P80188
A	42	GLN	LEU	engineered mutation	UNP P80188
A	49	PRO	GLN	engineered mutation	UNP P80188
A	52	ILE	TYR	engineered mutation	UNP P80188
A	68	ASN	SER	engineered mutation	UNP P80188
A	70	ALA	LEU	engineered mutation	UNP P80188
A	71	SER	PHE	engineered mutation	UNP P80188
A	72	ASN	ARG	engineered mutation	UNP P80188
A	73	TRP	LYS	engineered mutation	UNP P80188
A	74	GLU	LYS	engineered mutation	UNP P80188
A	77	THR	ASP	engineered mutation	UNP P80188
A	79	ARG	TRP	engineered mutation	UNP P80188
A	81	LYS	ARG	engineered mutation	UNP P80188
A	87	SER	CYS	engineered mutation	UNP P80188
A	96	GLU	ASN	engineered mutation	UNP P80188
A	100	ARG	TYR	engineered mutation	UNP P80188
A	103	MET	LEU	engineered mutation	UNP P80188
A	125	THR	LYS	engineered mutation	UNP P80188
A	127	VAL	SER	engineered mutation	UNP P80188
A	132	LYS	TYR	engineered mutation	UNP P80188
A	134	TRP	LYS	engineered mutation	UNP P80188

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	28	HIS	GLN	engineered mutation	UNP P80188
B	36	PHE	LEU	engineered mutation	UNP P80188
B	38	GLU	GLY	engineered mutation	UNP P80188
B	40	ILE	ALA	engineered mutation	UNP P80188
B	41	GLN	ILE	engineered mutation	UNP P80188
B	42	GLN	LEU	engineered mutation	UNP P80188
B	49	PRO	GLN	engineered mutation	UNP P80188
B	52	ILE	TYR	engineered mutation	UNP P80188
B	68	ASN	SER	engineered mutation	UNP P80188
B	70	ALA	LEU	engineered mutation	UNP P80188
B	71	SER	PHE	engineered mutation	UNP P80188
B	72	ASN	ARG	engineered mutation	UNP P80188
B	73	TRP	LYS	engineered mutation	UNP P80188
B	74	GLU	LYS	engineered mutation	UNP P80188
B	77	THR	ASP	engineered mutation	UNP P80188
B	79	ARG	TRP	engineered mutation	UNP P80188
B	81	LYS	ARG	engineered mutation	UNP P80188
B	87	SER	CYS	engineered mutation	UNP P80188
B	96	GLU	ASN	engineered mutation	UNP P80188
B	100	ARG	TYR	engineered mutation	UNP P80188
B	103	MET	LEU	engineered mutation	UNP P80188
B	125	THR	LYS	engineered mutation	UNP P80188
B	127	VAL	SER	engineered mutation	UNP P80188
B	132	LYS	TYR	engineered mutation	UNP P80188
B	134	TRP	LYS	engineered mutation	UNP P80188


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	156	Total O 156 156	0	0
2	B	183	Total O 183 183	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Neutrophil gelatinase-associated lipocalin



- Molecule 1: Neutrophil gelatinase-associated lipocalin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	38.99Å 65.95Å 67.94Å 90.00° 96.77° 90.00°	Depositor
Resolution (Å)	33.41 – 1.40 33.39 – 1.40	Depositor EDS
% Data completeness (in resolution range)	96.0 (33.41-1.40) 96.0 (33.39-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.98 (at 1.40Å)	Xtrriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.137 , 0.181 0.138 , 0.182	Depositor DCC
R_{free} test set	3254 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtrriage
Anisotropy	0.612	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3288	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/1548	0.78	0/2098
1	B	0.59	0/1541	0.81	0/2088
All	All	0.59	0/3089	0.80	0/4186

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1475	0	1475	13	0
1	B	1474	0	1479	24	0
2	A	156	0	0	0	0
2	B	183	0	0	4	0
All	All	3288	0	2954	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:100[A]:ARG:HG2	1:B:100[A]:ARG:NH1	1.68	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:100[A]:ARG:HG2	1:B:100[A]:ARG:HH11	0.89	1.01
1:B:100[A]:ARG:HH11	1:B:100[A]:ARG:CG	1.79	0.94
1:A:73:TRP:CE2	1:A:74:GLU:HG3	2.32	0.65
1:A:25:ASN:O	1:A:28[B]:HIS:CE1	2.54	0.61
1:B:142:LYS:HE2	1:B:177:ASP:OD1	2.02	0.60
1:A:59:LYS:HE3	1:A:63[A]:SER:OG	2.01	0.59
1:A:106:TYR:CD2	1:A:125[A]:THR:HG22	2.38	0.58
1:A:42:GLN:O	1:A:43:ARG:HG3	2.04	0.57
1:A:43:ARG:O	1:A:44:GLU:HB2	2.03	0.57
1:B:8:ILE:CD1	1:B:165:HIS:CE1	2.88	0.57
1:B:8:ILE:HD12	1:B:165:HIS:CE1	2.40	0.56
1:A:59:LYS:HD3	1:A:65[A]:ASN:ND2	2.22	0.55
1:A:157:LYS:HE2	1:A:163:GLU:HG3	1.89	0.54
1:B:100[A]:ARG:NH1	1:B:100[A]:ARG:CG	2.50	0.52
1:A:25:ASN:O	1:A:28[B]:HIS:ND1	2.44	0.51
1:B:59:LYS:CE	1:B:63[A]:SER:OG	2.59	0.51
1:B:8:ILE:HG21	1:B:161:LEU:HD23	1.94	0.50
1:B:42:GLN:O	1:B:43:ARG:HB2	2.12	0.50
1:B:100[B]:ARG:NH2	2:B:201:HOH:O	2.24	0.50
1:B:73:TRP:CE2	1:B:74:GLU:HG3	2.47	0.49
1:B:50[B]:LYS:HD3	1:B:172:ILE:HG21	1.95	0.48
1:B:174:GLN:H	1:B:174:GLN:HE21	1.62	0.48
1:B:157:LYS:HE2	1:B:163:GLU:HG3	1.96	0.48
1:B:174:GLN:NE2	2:B:203:HOH:O	2.45	0.47
1:A:59:LYS:CE	1:A:63[A]:SER:OG	2.63	0.47
1:B:96:GLU:HB3	1:B:100[A]:ARG:NH1	2.30	0.47
1:A:33:VAL:HG13	1:A:136[B]:THR:HG23	1.98	0.46
1:A:136[B]:THR:HG21	1:A:138:TYR:CE1	2.52	0.44
1:B:135:ILE:HD11	1:B:159:LEU:HD12	1.98	0.44
1:B:96:GLU:O	1:B:100[A]:ARG:NH1	2.50	0.44
1:B:151[B]:ASN:ND2	2:B:206:HOH:O	2.50	0.43
1:B:14:SER:HB3	2:B:245:HOH:O	2.20	0.42
1:B:99:SER:OG	1:B:100[A]:ARG:NH1	2.48	0.42
1:B:157:LYS:CE	1:B:163:GLU:HG3	2.50	0.42
1:B:59:LYS:HE2	1:B:63[A]:SER:OG	2.20	0.41
1:A:43:ARG:O	1:A:44:GLU:CB	2.68	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	183/174 (105%)	176 (96%)	5 (3%)	2 (1%)	14	2
1	B	181/174 (104%)	175 (97%)	5 (3%)	1 (1%)	25	7
All	All	364/348 (105%)	351 (96%)	10 (3%)	3 (1%)	17	4

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	ARG
1	A	44	GLU
1	B	43	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/160 (107%)	170 (99%)	1 (1%)	86	70
1	B	169/160 (106%)	164 (97%)	5 (3%)	41	10
All	All	340/320 (106%)	334 (98%)	6 (2%)	62	28

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	B	13	LEU
1	B	42	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	100[A]	ARG
1	B	100[B]	ARG
1	B	174	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	174/174 (100%)	-0.09	9 (5%) 27 26	10, 15, 36, 84	0
1	B	174/174 (100%)	-0.01	11 (6%) 20 18	9, 14, 35, 73	0
All	All	348/348 (100%)	-0.05	20 (5%) 23 22	9, 15, 36, 84	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	41	GLN	6.4
1	B	73	TRP	6.0
1	B	40	ILE	4.5
1	A	5	SER	4.5
1	A	178	GLY	4.3
1	B	42	GLN	3.7
1	A	42	GLN	3.7
1	B	178	GLY	3.7
1	A	45	ASP	3.6
1	A	43	ARG	3.6
1	B	7	LEU	3.5
1	A	41	GLN	3.4
1	B	45	ASP	3.2
1	A	44	GLU	3.1
1	A	73	TRP	2.7
1	B	43	ARG	2.6
1	A	72	ASN	2.6
1	B	72	ASN	2.6
1	B	46	LYS	2.5
1	B	44	GLU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.