

# wwPDB NMR Structure Validation Summary Report (i)

#### Aug 21, 2022 – 08:14 AM EDT

PDB ID : 2GVB

Title : REFINED SOLUTION STRUCTURE OF THE TYR 41-> HIS MUTANT

OF THE M13 GENE V PROTEIN. A COMPARISON WITH THE CRYSTAL

STRUCTURE

Authors: Folkers, P.J.M.; Nilges, M.; Folmer, R.H.A.; Prompers, J.J.; Konings, R.N.H.;

Hilbers, C.W.

Deposited on : 1995-07-27

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

ShiftChecker : 2.29

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.29

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $SOLUTION\ NMR$ 

The overall completeness of chemical shifts assignment was not calculated.

There are no overall percentile quality scores available for this entry.

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	87	100%
1	В	87	100%



## 2 Ensemble composition and analysis (i)

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.



## 3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2760 atoms, of which 1400 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called GENE V PROTEIN.

$\mathbf{M}$	ol	Chain	Residues		Atoms							
1	1	Λ	87	Total	С	Н	N	О	S	0		
1	L	A	01	1380	435	700	117	125	3			
1		D	87	Total	С	Н	N	О	S	0		
1	1	Б	01	1380	435	700	117	125	3	U		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	41	HIS	TYR	conflict	UNP P69544
В	41	HIS	TYR	conflict	UNP P69544



## 4 Residue-property plots (i)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: GENE V PROTEIN

Chain	A:																		10	0%																					
M1 I2 K3 V4	E5 I6 K7	84 88 89 89	Q10	A11	F13	T14	T15 R16	S17	G18	6I >	R21	q22	G23	P25	Y26	S27	L28	E30	<b>Q31</b>	L32	Y34	V35	D36	638	N39	E40	H41	V43	L44	V45	147	T48	1.49	E51	G52	<b>0</b> 53	P54	V.56	A57	P58	G59 L60
Y61 T62 V63 H64	266 S66 S67	F68 K69	07V	G71 072	F73	G74	S/5 1.76	M7.7		670	L81	R82	L83	V84	A86	K87																									
• Mol	ecu	le i	1:	G	Ε	N	Ε	V	F	PΕ	RC	Γ(	ГΕ	ΣIJ	N																										
Chain	В:																		10	0%																					
M1 I2 K3 V4	E5 I6 K7	89 89 89	q10	A11	F13	T14	T15 R16	S17	G18	419	R21	q22	G23	P25	Y26	S27	L28	E30	<b>431</b>	L32	Y34	V35	D36	638	N39	E40	H41	V43	1.44	V45	147	T48	1.49	E51	G52	053	P54 AEE	VSG	A57	P58	G29 L60
Y61 T62 V63 H64	L65 S66 S67	F68 K69	070	G71 072	F73	G74	S75 1.76	M7.7	178	D/9	L81	R82	L83	V84 P85	A86	K87																									



#### Refinement protocol and experimental data overview (i) 5



Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DIANA	refinement	
X-PLOR	refinement	

No chemical shift data was provided.



## 6 Model quality (i)

### 6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	0	0	0	0
1	В	0	0	0	0
All	All	0	0	0	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

### 6.3 Torsion angles (i)

#### 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Analysed   Favoured   Allowed			Percentiles				
1	A	0	-	-	_	-				
1	В	0	-	=	-	-				
All	All	0	-	=	-	-				

There are no Ramachandran outliers.



#### 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	nalysed   Rotameric   Outliers				
1	A	0	-	-	-		
1	В	0	-	-	-		
All	All	0	-	-	-		

There are no protein residues with a non-rotameric sidechain to report.

#### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.6 Ligand geometry (i)

There are no ligands in this entry.

### 6.7 Other polymers (i)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 7 Chemical shift validation (i)

No chemical shift data were provided

