



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 12:23 pm BST

PDB ID : 6GVM
Title : Tubulin:F3II DARPin complex
Authors : Gigant, B.; Campanacci, V.; Cantos Fernandes, S.
Deposited on : 2018-06-21
Resolution : 3.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	 4% 86% 8% 5%
2	B	445	 3% 82% 13% ••
3	C	87	 10% 49% 24% • 25%
4	D	169	 9% 77% 16% • 6%

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 8397 atoms, of which 1 is hydrogen and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	428	3323	2105	566	631	21	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	SER	GLY	conflict	UNP D0VWZ0
A	340	SER	THR	conflict	UNP D0VWZ0

- Molecule 2 is a protein called Tubulin beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	B	425	3332	2092	1	568	645	26	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	203	CYS	SER	conflict	UNP D0VWY9
B	318	ILE	VAL	conflict	UNP D0VWY9


- Molecule 3 is a protein called STATHMIN-LIKE DOMAIN R1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	65	500	313	93	92	2	0	0	0

- Molecule 4 is a protein called F3II DARPIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	159	1181	742	204	234	1	0	0	0

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	A	1	32	10	5	14	3	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
6	A	1	1	1	0	0

- Molecule 7 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
7	B	1	28	10	5	11	2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha chain




- Molecule 2: Tubulin beta chain




- Molecule 3: STATHMIN-LIKE DOMAIN R1



- Molecule 4: F3II DARPIN





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	189.27Å 51.72Å 144.14Å 90.00° 121.03° 90.00°	Depositor
Resolution (Å)	49.28 – 3.50 49.28 – 3.50	Depositor EDS
% Data completeness (in resolution range)	50.0 (49.28-3.50) 50.1 (49.28-3.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 3.48Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.245 , 0.270 0.267 , 0.290	Depositor DCC
R_{free} test set	738 reflections (9.51%)	wwPDB-VP
Wilson B-factor (Å ²)	81.5	Xtrriage
Anisotropy	0.062	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 53.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	8397	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, GTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/3396	0.64	0/4613
2	B	0.42	0/3404	0.63	0/4612
3	C	0.46	0/506	0.66	0/678
4	D	0.41	0/1199	0.69	0/1631
All	All	0.43	0/8505	0.64	0/11534

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3323	0	3231	30	0
2	B	3331	1	3206	32	0
3	C	500	0	491	22	0
4	D	1181	0	1149	18	0
5	A	32	0	12	0	0
6	A	1	0	0	0	0
7	B	28	0	12	0	0
All	All	8396	1	8101	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:51:LEU:HD21	4:D:71:ALA:HB1	1.29	1.10
4:D:56:THR:O	4:D:90:VAL:HG11	1.76	0.85
2:B:108:TYR:HB3	3:C:82:VAL:HG21	1.65	0.79
1:A:336:LYS:HD3	3:C:24:LEU:HG	1.69	0.74
1:A:112:LYS:HE2	3:C:58:GLU:HG2	1.72	0.69
1:A:180:ALA:HB3	1:A:183:GLU:HG3	1.77	0.65
2:B:401:ARG:HB3	4:D:90:VAL:HG13	1.80	0.64
1:A:352:LYS:HG2	3:C:21:GLU:HG2	1.81	0.61
1:A:101:ASN:ND2	1:A:180:ALA:HB2	2.16	0.59
2:B:295:MET:HG2	2:B:377:PHE:HB2	1.86	0.58
2:B:106:GLY:O	2:B:111:GLY:HA3	2.04	0.57
3:C:79:ALA:O	3:C:82:VAL:HG22	2.06	0.56
1:A:412:GLY:O	3:C:57:ALA:HB1	2.06	0.56
1:A:112:LYS:HE2	3:C:58:GLU:CG	2.35	0.56
1:A:221:ARG:NE	2:B:325:MET:HB3	2.22	0.55
4:D:51:LEU:CD2	4:D:71:ALA:HB1	2.19	0.54
1:A:372:GLN:NE2	1:A:373:ARG:NH2	2.57	0.54
1:A:350:GLY:HA3	3:C:23:ILE:HD13	1.89	0.53
2:B:404:PHE:CZ	4:D:48:LEU:HD11	2.43	0.53
4:D:115:THR:HG22	4:D:118:HIS:ND1	2.25	0.52
3:C:8:VAL:HG13	3:C:20:TRP:CD1	2.45	0.52
2:B:214:PHE:O	2:B:218:LYS:HA	2.10	0.51
2:B:269:MET:HG3	2:B:303:ALA:HB3	1.92	0.51
1:A:101:ASN:HD22	2:B:258:ASN:HD21	1.58	0.51
1:A:349:THR:CG2	3:C:25:LYS:H	2.24	0.51
2:B:75:MET:HG3	2:B:94:PHE:CD1	2.47	0.50
2:B:162:PRO:HB2	3:C:65:PHE:CD1	2.48	0.49
4:D:118:HIS:HD2	4:D:152:ILE:HD11	1.77	0.49
1:A:307:PRO:HA	1:A:383:ALA:HB2	1.95	0.49
2:B:66:ILE:HG12	2:B:121:VAL:HG12	1.95	0.48
1:A:247:ALA:HB1	3:C:12:ASN:HB3	1.96	0.48
1:A:159:VAL:HG11	3:C:47:LEU:HB2	1.95	0.48
2:B:406:HIS:NE2	2:B:407:TRP:CD1	2.81	0.48
1:A:28:HIS:CE1	1:A:49:PHE:HB3	2.50	0.47
2:B:212:ILE:O	2:B:216:THR:HB	2.14	0.47
1:A:349:THR:HG22	3:C:25:LYS:H	1.80	0.47
1:A:320:ARG:HA	1:A:356:ASN:O	2.15	0.47
2:B:294:GLN:HE21	2:B:300:ASN:HD21	1.64	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:172:MET:HG3	2:B:387:LEU:HD21	1.96	0.46
1:A:159:VAL:HG22	3:C:46:SER:HB2	1.97	0.46
1:A:406:HIS:CG	2:B:263:PRO:HD3	2.51	0.46
4:D:30:VAL:O	4:D:34:MET:HG2	2.15	0.46
2:B:59:ASN:CG	2:B:59:ASN:O	2.52	0.46
2:B:118:VAL:HG11	2:B:153:LEU:HD11	1.99	0.45
4:D:129:VAL:HG21	4:D:161:LEU:HD11	1.99	0.45
1:A:109:THR:HG21	3:C:61:ARG:CD	2.47	0.45
4:D:90:VAL:O	4:D:90:VAL:HG12	2.17	0.45
2:B:404:PHE:HZ	4:D:48:LEU:HD11	1.80	0.44
2:B:162:PRO:HB2	3:C:65:PHE:HD1	1.82	0.44
4:D:114:TYR:HH	4:D:145:PHE:HE2	1.63	0.44
4:D:56:THR:HG23	4:D:86:LEU:HB3	1.99	0.44
1:A:156:ARG:HD3	3:C:50:ILE:HG21	1.99	0.44
2:B:83:PHE:O	2:B:86:ILE:HG22	2.18	0.44
1:A:412:GLY:O	3:C:57:ALA:CB	2.66	0.43
2:B:396:THR:O	2:B:400:ARG:HG3	2.19	0.43
2:B:422:GLU:HA	2:B:422:GLU:OE1	2.18	0.43
1:A:167:LEU:HG	1:A:200:CYS:HB3	2.01	0.43
2:B:217:LEU:HD22	2:B:277:SER:HB2	1.99	0.43
1:A:307:PRO:HA	1:A:383:ALA:CB	2.49	0.43
1:A:109:THR:HG21	3:C:61:ARG:HD2	2.01	0.43
1:A:221:ARG:HD2	2:B:326:LYS:N	2.34	0.42
2:B:406:HIS:NE2	4:D:81:ASP:OD2	2.53	0.42
3:C:47:LEU:HA	3:C:50:ILE:HD12	2.01	0.42
4:D:61:GLU:O	4:D:65:VAL:HG23	2.19	0.42
2:B:2:ARG:HB3	2:B:133:GLN:HE21	1.84	0.41
2:B:308:ARG:HG3	2:B:342:TYR:CZ	2.55	0.41
4:D:25:GLY:HA2	4:D:62:ILE:HD12	2.02	0.41
4:D:48:LEU:HD22	4:D:52:HIS:HB3	2.02	0.41
4:D:84:LEU:HD11	4:D:100:LEU:HD22	2.02	0.41
1:A:265:ILE:HG23	1:A:432:TYR:CE1	2.56	0.41
1:A:31:GLN:HE21	1:A:37:PRO:HD3	1.86	0.41
2:B:276:THR:HB	2:B:277:SER:H	1.71	0.41
3:C:13:LYS:HG2	3:C:18:GLN:HG3	2.01	0.41
2:B:171:VAL:HA	2:B:204:ILE:O	2.22	0.40
1:A:179:THR:HG22	2:B:248:LEU:HG	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/451 (94%)	411 (97%)	13 (3%)	0	100	100
2	B	421/445 (95%)	411 (98%)	10 (2%)	0	100	100
3	C	61/87 (70%)	57 (93%)	4 (7%)	0	100	100
4	D	157/169 (93%)	155 (99%)	2 (1%)	0	100	100
All	All	1063/1152 (92%)	1034 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	355/379 (94%)	345 (97%)	10 (3%)	43	72
2	B	365/383 (95%)	345 (94%)	20 (6%)	21	54
3	C	49/76 (64%)	44 (90%)	5 (10%)	7	32
4	D	120/133 (90%)	112 (93%)	8 (7%)	16	48
All	All	889/971 (92%)	846 (95%)	43 (5%)	25	60

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	84	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	179	THR
1	A	181	VAL
1	A	188	ILE
1	A	279	GLU
1	A	297	GLU
1	A	302	MET
1	A	335	ILE
1	A	349	THR
2	B	41	ASP
2	B	48	ARG
2	B	57	THR
2	B	75	MET
2	B	124	LYS
2	B	139	HIS
2	B	153	LEU
2	B	207	GLU
2	B	221	THR
2	B	229	HIS
2	B	241	CYS
2	B	247	GLN
2	B	248	LEU
2	B	275	LEU
2	B	278	ARG
2	B	293	GLN
2	B	295	MET
2	B	325	MET
2	B	338	LYS
2	B	371	LEU
3	C	11	LEU
3	C	24	LEU
3	C	49	GLU
3	C	84	LYS
3	C	85	ASN
4	D	28	ASP
4	D	40	VAL
4	D	56	THR
4	D	76	THR
4	D	100	LEU
4	D	106	VAL
4	D	122	LEU
4	D	164	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	101	ASN
1	A	107	HIS
1	A	285	GLN
1	A	301	GLN
2	B	300	ASN
4	D	92	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GTP	A	501	6	26,34,34	1.11	2 (7%)	33,54,54	2.09	6 (18%)
7	GDP	B	501	-	24,30,30	1.16	2 (8%)	31,47,47	2.10	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	A	501	6	-	4/18/38/38	0/3/3/3
7	GDP	B	501	-	-	5/12/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	501	GTP	C6-N1	3.42	1.39	1.33
7	B	501	GDP	C6-N1	3.27	1.38	1.33
7	B	501	GDP	C6-C5	3.12	1.46	1.41
5	A	501	GTP	PA-O5'	2.26	1.68	1.59

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	501	GDP	C5-C6-N1	-8.22	112.19	123.43
5	A	501	GTP	C5-C6-N1	-7.94	112.58	123.43
7	B	501	GDP	C6-N1-C2	6.03	125.50	115.93
5	A	501	GTP	C6-N1-C2	5.78	125.11	115.93
5	A	501	GTP	PB-O3B-PG	-3.85	119.60	132.83
7	B	501	GDP	C6-C5-C4	-3.40	117.56	120.80
5	A	501	GTP	C6-C5-C4	-3.02	117.92	120.80
5	A	501	GTP	N3-C2-N1	-2.87	123.39	127.22
7	B	501	GDP	N3-C2-N1	-2.68	123.64	127.22
5	A	501	GTP	C2-N3-C4	-2.16	112.89	115.36
7	B	501	GDP	C2-N3-C4	-2.01	113.06	115.36

There are no chirality outliers.


All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	501	GTP	C5'-O5'-PA-O1A
5	A	501	GTP	C5'-O5'-PA-O2A
7	B	501	GDP	C5'-O5'-PA-O1A
5	A	501	GTP	C5'-O5'-PA-O3A
7	B	501	GDP	C5'-O5'-PA-O3A
7	B	501	GDP	C5'-O5'-PA-O2A
5	A	501	GTP	PB-O3A-PA-O2A
7	B	501	GDP	PB-O3A-PA-O1A
7	B	501	GDP	PB-O3A-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/451 (94%)	0.24	20 (4%) 31 28	16, 56, 113, 155	0
2	B	425/445 (95%)	0.15	14 (3%) 46 41	20, 48, 89, 139	0
3	C	65/87 (74%)	0.67	9 (13%) 2 3	39, 72, 133, 209	0
4	D	159/169 (94%)	0.46	15 (9%) 8 9	7, 49, 126, 259	0
All	All	1077/1152 (93%)	0.26	58 (5%) 25 23	7, 53, 106, 259	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	169	ASN	6.2
1	A	336	LYS	5.9
4	D	167	LYS	5.3
3	C	6	MET	4.9
4	D	166	GLN	4.6
3	C	26	PRO	4.5
1	A	348	PRO	3.6
4	D	140	ASN	3.5
3	C	21	GLU	3.5
3	C	27	PRO	3.4
3	C	10	GLU	3.3
4	D	135	ASN	3.2
2	B	415	GLU	3.1
4	D	106	VAL	3.0
2	B	359	PRO	2.9
1	A	412	GLY	2.9
4	D	128	ILE	2.9
4	D	168	LEU	2.9
1	A	47	ASP	2.8
4	D	138	ASP	2.8
1	A	313	MET	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	293	ASN	2.7
4	D	153	SER	2.7
1	A	337	THR	2.7
2	B	32	PRO	2.6
4	D	161	LEU	2.6
2	B	435	TYR	2.6
3	C	28	SER	2.6
2	B	441	ASP	2.6
2	B	25	SER	2.6
2	B	83	PHE	2.5
2	B	81	GLY	2.5
1	A	351	PHE	2.5
4	D	163	GLU	2.4
2	B	369	ARG	2.3
2	B	408	TYR	2.3
2	B	82	PRO	2.3
2	B	321	GLY	2.3
2	B	38	GLY	2.3
3	C	45	PRO	2.2
1	A	245	ASP	2.2
1	A	30	ILE	2.2
4	D	165	LEU	2.2
1	A	346	TRP	2.1
1	A	434	GLU	2.1
1	A	357	TYR	2.1
1	A	199	ASP	2.1
4	D	137	ALA	2.1
1	A	433	GLU	2.1
4	D	139	VAL	2.1
3	C	24	LEU	2.1
1	A	295	CYS	2.1
1	A	329	ASN	2.1
1	A	372	GLN	2.0
1	A	322	ASP	2.0
1	A	349	THR	2.0
2	B	414	ASP	2.0
3	C	75	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GTP	A	501	32/32	0.92	0.23	17,22,28,35	0
7	GDP	B	501	28/28	0.94	0.22	27,33,37,42	0
6	MG	A	502	1/1	0.95	0.45	59,59,59,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.