



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 10, 2023 – 09:10 am GMT

PDB ID : 8H26  
Title : Crystal structure of MnmM from *S. aureus* complexed with SAH (1.50 Å)  
Authors : Kim, J.; Cho, G.; Lee, J.  
Deposited on : 2022-10-05  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.3  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.3


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	195	 3% 77% 13% 10%
1	B	195	 3% 84% 6% 10%
1	C	195	 3% 78% 15% 6%
1	D	195	 4% 79% 10% 10%
1	E	195	 5% 77% 10% 13%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	195	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment at the beginning labeled '3%', a large green segment labeled '78%', a small yellow segment labeled '6%', and a grey segment at the end labeled '14%'.</p>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9201 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 16S rRNA (Cytosine(1402)-N(4))-methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	175	1406	896	241	263	6	0	2	0
1	B	175	1402	893	241	262	6	0	2	0
1	C	184	1495	951	259	278	7	0	5	0
1	D	175	1413	897	244	266	6	0	4	0
1	E	169	1353	860	235	253	5	0	2	0
1	F	168	1350	860	234	251	5	0	2	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	188	GLY	-	expression tag	UNP Q2FXG9
A	189	HIS	-	expression tag	UNP Q2FXG9
A	190	HIS	-	expression tag	UNP Q2FXG9
A	191	HIS	-	expression tag	UNP Q2FXG9
A	192	HIS	-	expression tag	UNP Q2FXG9
A	193	HIS	-	expression tag	UNP Q2FXG9
A	194	HIS	-	expression tag	UNP Q2FXG9
A	195	GLY	-	expression tag	UNP Q2FXG9
B	188	GLY	-	expression tag	UNP Q2FXG9
B	189	HIS	-	expression tag	UNP Q2FXG9
B	190	HIS	-	expression tag	UNP Q2FXG9
B	191	HIS	-	expression tag	UNP Q2FXG9
B	192	HIS	-	expression tag	UNP Q2FXG9
B	193	HIS	-	expression tag	UNP Q2FXG9
B	194	HIS	-	expression tag	UNP Q2FXG9
B	195	GLY	-	expression tag	UNP Q2FXG9
C	188	GLY	-	expression tag	UNP Q2FXG9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	189	HIS	-	expression tag	UNP Q2FXG9
C	190	HIS	-	expression tag	UNP Q2FXG9
C	191	HIS	-	expression tag	UNP Q2FXG9
C	192	HIS	-	expression tag	UNP Q2FXG9
C	193	HIS	-	expression tag	UNP Q2FXG9
C	194	HIS	-	expression tag	UNP Q2FXG9
C	195	GLY	-	expression tag	UNP Q2FXG9
D	188	GLY	-	expression tag	UNP Q2FXG9
D	189	HIS	-	expression tag	UNP Q2FXG9
D	190	HIS	-	expression tag	UNP Q2FXG9
D	191	HIS	-	expression tag	UNP Q2FXG9
D	192	HIS	-	expression tag	UNP Q2FXG9
D	193	HIS	-	expression tag	UNP Q2FXG9
D	194	HIS	-	expression tag	UNP Q2FXG9
D	195	GLY	-	expression tag	UNP Q2FXG9
E	188	GLY	-	expression tag	UNP Q2FXG9
E	189	HIS	-	expression tag	UNP Q2FXG9
E	190	HIS	-	expression tag	UNP Q2FXG9
E	191	HIS	-	expression tag	UNP Q2FXG9
E	192	HIS	-	expression tag	UNP Q2FXG9
E	193	HIS	-	expression tag	UNP Q2FXG9
E	194	HIS	-	expression tag	UNP Q2FXG9
E	195	GLY	-	expression tag	UNP Q2FXG9
F	188	GLY	-	expression tag	UNP Q2FXG9
F	189	HIS	-	expression tag	UNP Q2FXG9
F	190	HIS	-	expression tag	UNP Q2FXG9
F	191	HIS	-	expression tag	UNP Q2FXG9
F	192	HIS	-	expression tag	UNP Q2FXG9
F	193	HIS	-	expression tag	UNP Q2FXG9
F	194	HIS	-	expression tag	UNP Q2FXG9
F	195	GLY	-	expression tag	UNP Q2FXG9

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C<sub>14</sub>H<sub>20</sub>N<sub>6</sub>O<sub>5</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	C	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	D	1	Total	C	N	O	S	0	0
			26	14	6	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	119	Total	O	0	0
			119	119		
3	B	134	Total	O	0	0
			134	134		
3	C	130	Total	O	0	0
			130	130		
3	D	90	Total	O	0	0
			90	90		
3	E	111	Total	O	0	0
			111	111		
3	F	94	Total	O	0	0
			94	94		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 16S rRNA (Cytosine(1402)-N(4))-methyltransferase




- Molecule 1: 16S rRNA (Cytosine(1402)-N(4))-methyltransferase




- Molecule 1: 16S rRNA (Cytosine(1402)-N(4))-methyltransferase






- Molecule 1: 16S rRNA (Cytosine(1402)-N(4))-methyltransferase







- Molecule 1: 16S rRNA (Cytosine(1402)-N(4))-methyltransferase



- Molecule 1: 16S rRNA (Cytosine(1402)-N(4))-methyltransferase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.33Å 69.36Å 74.61Å 91.25° 112.97° 114.05°	Depositor
Resolution (Å)	67.09 – 1.50 67.08 – 1.50	Depositor EDS
% Data completeness (in resolution range)	63.4 (67.09-1.50) 63.4 (67.08-1.50)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.82 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.183 , 0.219 0.190 , 0.225	Depositor DCC
$R_{free}$ test set	5112 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.6	Xtrriage
Anisotropy	0.015	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 39.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9201	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	1/1432 (0.1%)	0.86	3/1937 (0.2%)
1	B	0.68	0/1427	0.81	1/1930 (0.1%)
1	C	0.73	1/1525 (0.1%)	0.85	3/2063 (0.1%)
1	D	0.72	2/1438 (0.1%)	0.81	0/1944
1	E	0.70	2/1379 (0.1%)	0.82	1/1865 (0.1%)
1	F	0.71	1/1376 (0.1%)	0.89	1/1861 (0.1%)
All	All	0.71	7/8577 (0.1%)	0.84	9/11600 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	58	GLU	CG-CD	8.71	1.65	1.51
1	A	21	GLU	CD-OE2	-6.62	1.18	1.25
1	C	122	SER	CB-OG	-6.49	1.33	1.42
1	D	44	GLU	CD-OE2	5.47	1.31	1.25
1	F	4	GLU	CD-OE1	-5.10	1.20	1.25
1	E	29[A]	CYS	CB-SG	-5.01	1.73	1.81
1	E	29[B]	CYS	CB-SG	-5.01	1.73	1.81

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	86	ASP	CB-CG-OD2	6.37	124.03	118.30
1	B	153	ASP	CB-CG-OD1	6.15	123.83	118.30
1	A	114	ASP	CB-CG-OD2	6.14	123.83	118.30
1	F	153	ASP	CB-CG-OD1	5.72	123.45	118.30
1	C	55	LEU	CB-CG-CD1	-5.72	101.28	111.00
1	C	62	ASP	CB-CG-OD2	-5.65	113.22	118.30
1	A	86	ASP	CB-CG-OD1	-5.34	113.49	118.30
1	C	62	ASP	CB-CG-OD1	5.27	123.05	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	E	86	ASP	CB-CG-OD1	5.00	122.80	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1406	0	1401	16	0
1	B	1402	0	1403	9	0
1	C	1495	0	1488	26	0
1	D	1413	0	1407	13	0
1	E	1353	0	1342	10	0
1	F	1350	0	1339	8	0
2	A	26	0	19	0	0
2	B	26	0	19	1	0
2	C	26	0	19	2	0
2	D	26	0	19	2	0
3	A	119	0	0	1	0
3	B	134	0	0	1	0
3	C	130	0	0	6	1
3	D	90	0	0	3	0
3	E	111	0	0	3	1
3	F	94	0	0	2	0
All	All	9201	0	8456	72	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:155:LEU:HD13	1:C:181[B]:CYS:SG	1.94	1.06
1:A:155:LEU:HD13	1:A:181[B]:CYS:SG	2.05	0.97
1:F:139:HIS:ND1	3:F:201:HOH:O	2.04	0.80

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:21:GLU:OE1	3:D:301:HOH:O	2.01	0.78
1:C:118:GLN:OE1	3:C:301:HOH:O	2.02	0.77
1:C:155:LEU:CD1	1:C:181[B]:CYS:SG	2.75	0.73
1:D:144:GLY:N	3:D:303:HOH:O	2.26	0.69
1:C:123:LEU:O	1:C:127[A]:MET:HG2	1.94	0.68
1:B:123:LEU:O	1:B:127[B]:MET:HG2	1.93	0.67
1:B:153:ASP:HB3	1:D:145:GLN:NE2	2.10	0.67
1:A:123:LEU:O	1:A:127[A]:MET:HG2	1.94	0.67
1:C:147:GLU:OE1	3:C:302:HOH:O	2.12	0.66
1:C:145:GLN:OE1	3:C:303:HOH:O	2.13	0.65
1:C:7:LEU:HD23	3:C:393:HOH:O	1.97	0.64
1:D:157:THR:OG1	3:D:302:HOH:O	2.02	0.64
1:E:84:ILE:O	1:E:89:LYS:NZ	2.33	0.62
1:D:123:LEU:O	1:D:127[A]:MET:HG2	2.00	0.61
1:A:7:LEU:HD12	1:C:21:GLU:HG2	1.81	0.61
1:A:155:LEU:CD1	1:A:181[B]:CYS:SG	2.86	0.61
1:C:55:LEU:C	1:C:55:LEU:HD12	2.22	0.60
1:D:14:ILE:HD12	1:D:94:ALA:HB1	1.82	0.60
1:C:17:HIS:HE1	1:C:184:GLU:OE1	1.85	0.60
1:A:7:LEU:HD12	1:C:21:GLU:CG	2.34	0.58
1:F:174:ARG:HG3	1:F:177:ALA:HB3	1.86	0.57
1:A:17:HIS:HE1	1:A:184:GLU:OE1	1.89	0.55
1:F:152:LEU:HD21	1:F:179:PHE:HZ	1.73	0.53
1:B:171:LEU:HD21	1:C:167:GLN:OE1	2.10	0.52
1:A:5:ARG:HD3	3:A:402:HOH:O	2.10	0.52
1:C:145:GLN:NE2	1:F:157:THR:OG1	2.44	0.50
1:C:7:LEU:HB2	1:C:8:PRO:HD3	1.93	0.50
1:C:98:ASN:HD22	2:C:201:SAH:HB2	1.77	0.50
1:F:139:HIS:CE1	1:F:175:ASN:O	2.65	0.49
1:D:98:ASN:HB2	2:D:201:SAH:HG1	1.95	0.48
1:C:65:LYS:HG2	3:C:312:HOH:O	2.14	0.48
1:C:13:LEU:O	1:C:17:HIS:HD2	1.97	0.47
1:B:159:ASP:OD1	1:B:161:LYS:HD3	2.14	0.47
1:E:148:LYS:N	3:E:209:HOH:O	2.47	0.47
1:D:98:ASN:HD22	2:D:201:SAH:HB2	1.80	0.47
1:C:101:TYR:CE1	1:C:110:VAL:HG13	2.50	0.46
1:A:13:LEU:O	1:A:17:HIS:HD2	1.98	0.46
1:C:98:ASN:ND2	2:C:201:SAH:HN1	2.14	0.46
1:A:61:ARG:HD2	1:A:72:LEU:HD12	1.97	0.45
1:A:7:LEU:CD1	1:C:21:GLU:CG	2.95	0.45
1:A:173:GLN:HG2	3:E:252:HOH:O	2.17	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:LYS:HE2	1:A:179:PHE:CE1	2.52	0.44
1:C:55:LEU:HD12	1:C:56:ALA:N	2.32	0.44
1:F:130:GLU:HG2	3:F:269:HOH:O	2.16	0.44
1:B:98:ASN:ND2	2:B:201:SAH:HN1	2.17	0.43
1:D:1:MET:SD	1:F:164:GLN:HG3	2.59	0.43
1:A:164:GLN:HG3	1:E:1:MET:SD	2.58	0.43
1:C:29[A]:CYS:SG	1:C:57:LEU:HD23	2.58	0.43
1:D:127[B]:MET:CE	1:D:133:ILE:HG13	2.49	0.43
1:E:41:GLN:NE2	3:E:212:HOH:O	2.52	0.43
1:E:162:HIS:HD2	1:E:187:SER:O	2.02	0.43
1:B:15:LYS:NZ	3:B:309:HOH:O	2.52	0.42
1:A:135:LEU:HD13	1:A:151:LEU:HD21	2.01	0.42
1:D:25:VAL:O	1:D:95:ALA:HA	2.19	0.42
1:D:112:LYS:HD3	1:D:114:ASP:HB2	2.02	0.42
1:C:25:VAL:O	1:C:95:ALA:HA	2.19	0.42
1:A:25:VAL:O	1:A:95:ALA:HA	2.20	0.41
1:A:139:HIS:HE1	1:A:177:ALA:O	2.04	0.41
1:B:25:VAL:O	1:B:95:ALA:HA	2.20	0.41
1:B:153:ASP:HB3	1:D:145:GLN:CD	2.41	0.41
1:E:139:HIS:HA	1:E:148:LYS:NZ	2.36	0.41
1:E:175:ASN:O	1:E:175:ASN:CG	2.59	0.41
1:E:162:HIS:HB2	1:E:189:HIS:O	2.21	0.41
1:E:37:PHE:O	1:E:41:GLN:HG2	2.21	0.41
1:E:25:VAL:O	1:E:95:ALA:HA	2.22	0.40
1:B:164:GLN:HG3	1:C:1:MET:SD	2.62	0.40
1:C:130:GLU:HG2	3:C:319:HOH:O	2.20	0.40
1:F:25:VAL:O	1:F:95:ALA:HA	2.21	0.40
1:C:135:LEU:HD13	1:C:151:LEU:HD21	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:406:HOH:O	3:E:280:HOH:O[1_665]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/195 (88%)	168 (98%)	3 (2%)	0	100	100
1	B	171/195 (88%)	170 (99%)	1 (1%)	0	100	100
1	C	185/195 (95%)	182 (98%)	3 (2%)	0	100	100
1	D	173/195 (89%)	171 (99%)	2 (1%)	0	100	100
1	E	165/195 (85%)	163 (99%)	2 (1%)	0	100	100
1	F	164/195 (84%)	161 (98%)	1 (1%)	2 (1%)	13	2
All	All	1029/1170 (88%)	1015 (99%)	12 (1%)	2 (0%)	47	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	140	GLY
1	F	175	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	157/171 (92%)	156 (99%)	1 (1%)	86	74
1	B	157/171 (92%)	156 (99%)	1 (1%)	86	74
1	C	167/171 (98%)	166 (99%)	1 (1%)	86	74
1	D	158/171 (92%)	155 (98%)	3 (2%)	57	27
1	E	151/171 (88%)	147 (97%)	4 (3%)	46	16

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	150/171 (88%)	147 (98%)	3 (2%)	55	25
All	All	940/1026 (92%)	927 (99%)	13 (1%)	67	42

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	HIS
1	B	68	ASN
1	C	139	HIS
1	D	57	LEU
1	D	68	ASN
1	D	77	HIS
1	E	77	HIS
1	E	115	THR
1	E	118	GLN
1	E	132	ILE
1	F	77	HIS
1	F	139	HIS
1	F	152	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	HIS
1	A	139	HIS
1	B	68	ASN
1	B	79	ASN
1	B	98	ASN
1	B	145	GLN
1	B	176	HIS
1	C	17	HIS
1	C	98	ASN
1	C	118	GLN
1	C	145	GLN
1	C	160	GLN
1	D	33	ASN
1	D	98	ASN
1	D	176	HIS
1	E	41	GLN
1	E	82	HIS
1	E	91	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	E	162	HIS
1	F	53	GLN
1	F	82	HIS
1	F	141	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SAH	C	201	-	24,28,28	0.68	0	25,40,40	0.98	2 (8%)
2	SAH	B	201	-	24,28,28	0.79	2 (8%)	25,40,40	0.97	2 (8%)
2	SAH	D	201	-	24,28,28	0.74	0	25,40,40	0.98	1 (4%)
2	SAH	A	201	-	24,28,28	0.74	1 (4%)	25,40,40	1.18	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	C	201	-	-	0/11/31/31	0/3/3/3
2	SAH	B	201	-	-	0/11/31/31	0/3/3/3
2	SAH	D	201	-	-	0/11/31/31	0/3/3/3
2	SAH	A	201	-	-	0/11/31/31	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	SAH	OXT-C	-2.21	1.23	1.30
2	B	201	SAH	C8-N7	-2.17	1.30	1.34
2	B	201	SAH	OXT-C	-2.09	1.23	1.30

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	SAH	OXT-C-O	-3.38	116.41	124.09
2	C	201	SAH	O4'-C1'-C2'	-2.78	102.86	106.93
2	A	201	SAH	OXT-C-CA	2.67	122.47	113.38
2	C	201	SAH	C5-C6-N6	2.52	124.19	120.35
2	B	201	SAH	C5-C6-N6	2.47	124.11	120.35
2	A	201	SAH	O4'-C1'-C2'	-2.45	103.34	106.93
2	D	201	SAH	C5-C6-N6	2.08	123.51	120.35
2	B	201	SAH	O4'-C1'-C2'	-2.04	103.94	106.93
2	A	201	SAH	C5-C6-N6	2.02	123.42	120.35

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	201	SAH	2	0
2	B	201	SAH	1	0
2	D	201	SAH	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	175/195 (89%)	0.19	6 (3%) 45 49	14, 28, 58, 71	0
1	B	175/195 (89%)	0.08	5 (2%) 51 56	14, 25, 51, 89	0
1	C	184/195 (94%)	0.13	5 (2%) 54 59	12, 24, 43, 53	0
1	D	175/195 (89%)	0.17	7 (4%) 38 42	14, 27, 52, 78	0
1	E	169/195 (86%)	0.24	10 (5%) 22 24	14, 27, 58, 88	0
1	F	168/195 (86%)	0.21	5 (2%) 50 55	15, 28, 52, 82	0
All	All	1046/1170 (89%)	0.17	38 (3%) 42 47	12, 27, 53, 89	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	110	VAL	5.7
1	F	148	LYS	5.6
1	E	189	HIS	5.2
1	F	101	TYR	4.8
1	F	141	HIS	4.5
1	C	187	SER	4.4
1	B	175	ASN	4.3
1	E	139	HIS	4.2
1	A	110	VAL	4.2
1	E	176	HIS	4.0
1	C	188	GLY	3.9
1	D	144	GLY	3.9
1	B	139	HIS	3.8
1	A	176	HIS	3.7
1	D	175	ASN	3.6
1	B	176	HIS	3.4
1	C	162[A]	HIS	3.4
1	D	55	LEU	3.3
1	B	174	ARG	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	145	GLN	3.0
1	D	176	HIS	3.0
1	B	144	GLY	2.9
1	D	174	ARG	2.8
1	E	173	GLN	2.8
1	E	113	PRO	2.8
1	C	101	TYR	2.7
1	E	175	ASN	2.6
1	E	148	LYS	2.6
1	E	149	HIS	2.5
1	F	139	HIS	2.5
1	A	57	LEU	2.5
1	A	101	TYR	2.4
1	E	80	ILE	2.4
1	F	173	GLN	2.4
1	E	186	ILE	2.3
1	A	54	ASP	2.2
1	D	173	GLN	2.1
1	A	53	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SAH	A	201	26/26	0.93	0.09	25,29,43,52	0
2	SAH	D	201	26/26	0.94	0.08	22,27,32,39	0
2	SAH	C	201	26/26	0.95	0.09	16,18,22,25	0
2	SAH	B	201	26/26	0.96	0.08	16,22,37,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around SAH D 201:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around SAH C 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.