



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 4, 2024 – 03:45 am GMT

PDB ID : 9H40
Title : Pinorexinol hydroxylase from Pseudomonas sp.
Authors : Guerriere, T.B.; Mattevi, A.
Deposited on : 2024-10-17
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	 5% 86% 10% ..
1	C	557	 5% 83% 12% ..

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8996 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called p-cresol methylhydroxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	538	Total 4273	C 2759	N 715	O 777	S 22	0	0	0
1	C	533	Total 4237	C 2738	N 707	O 770	S 22	0	0	0

- Molecule 2 is ALANINE (three-letter code: ALA) (formula: $C_3H_7NO_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 6	C 3	N 1	O 2	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	53	27	9	15	2	0	0
3	C	1	53	27	9	15	2	0	0

- Molecule 4 is GLYCINE (three-letter code: GLY) (formula: $C_2H_5NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	C	1	5	2	1	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	170	Total 170	O 170	0	0
5	C	199	Total 199	O 199	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: p-cresol methylhydroxylase



- Molecule 1: p-cresol methylhydroxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.21Å 131.81Å 137.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.76 – 1.80 51.76 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.1 (51.76-1.80) 98.1 (51.76-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 1.0	Depositor
R, R_{free}	0.186 , 0.228 0.197 , 0.234	Depositor DCC
R_{free} test set	4826 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtrriage
Anisotropy	0.057	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 34.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.000 for -h,l,k	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8996	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.25 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.6888e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/4400	0.88	8/5963 (0.1%)
1	C	0.70	0/4361	0.92	8/5909 (0.1%)
All	All	0.71	0/8761	0.90	16/11872 (0.1%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	162	ARG	NE-CZ-NH2	-19.18	110.71	120.30
1	A	162	ARG	NE-CZ-NH2	-16.79	111.91	120.30
1	C	162	ARG	NE-CZ-NH1	15.01	127.80	120.30
1	C	496	ARG	NE-CZ-NH1	12.28	126.44	120.30
1	C	496	ARG	NE-CZ-NH2	-10.66	114.97	120.30
1	A	162	ARG	NE-CZ-NH1	9.45	125.02	120.30
1	A	496	ARG	NE-CZ-NH1	8.49	124.55	120.30
1	C	496	ARG	CB-CG-CD	8.15	132.79	111.60
1	A	518	ARG	NE-CZ-NH2	-8.00	116.30	120.30
1	A	162	ARG	CG-CD-NE	-7.97	95.06	111.80
1	A	518	ARG	NE-CZ-NH1	6.77	123.69	120.30
1	C	162	ARG	CG-CD-NE	-6.11	98.97	111.80
1	A	496	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	A	496	ARG	CB-CG-CD	5.57	126.08	111.60
1	C	286	LEU	CA-CB-CG	5.51	127.98	115.30
1	C	162	ARG	CD-NE-CZ	5.42	131.19	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4273	0	4176	46	0
1	C	4237	0	4145	50	0
2	A	6	0	4	0	0
3	A	53	0	31	4	0
3	C	53	0	31	4	0
4	C	5	0	2	0	0
5	A	170	0	0	2	0
5	C	199	0	0	3	0
All	All	8996	0	8389	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:411:HIS:NE2	3:A:702:FAD:HM82	1.78	0.96
1:A:464:LEU:HD11	1:A:476:SER:HB3	1.62	0.80
1:A:206:MET:SD	1:C:419:PRO:HB3	2.26	0.76
1:A:411:HIS:CE1	3:A:702:FAD:HM82	2.24	0.72
1:A:156:VAL:HG13	1:A:230:MET:CE	2.22	0.70
1:C:227:VAL:HG11	1:C:230:MET:CE	2.23	0.69
1:A:227:VAL:HG11	1:A:230:MET:HE2	1.76	0.67
1:A:518:ARG:NH2	1:C:189:GLU:OE2	2.27	0.66
1:C:439:LEU:HD11	1:C:482:HIS:CD2	2.32	0.65
1:C:161:ASP:OD2	1:C:496:ARG:HD3	1.95	0.65
1:A:460:PHE:CD2	1:A:462:ILE:HD11	2.33	0.64
1:A:460:PHE:CE2	1:A:462:ILE:HD11	2.34	0.62
1:A:156:VAL:HG13	1:A:230:MET:HE3	1.80	0.61
1:A:242:LEU:HD23	1:A:243:LYS:N	2.16	0.60
1:A:462:ILE:HD12	1:A:462:ILE:N	2.16	0.60
1:A:89:GLY:HA2	1:A:411:HIS:O	2.03	0.59
3:C:602:FAD:H8A	3:C:602:FAD:O5B	2.02	0.59
1:A:518:ARG:HH22	1:C:189:GLU:CD	2.07	0.58
1:C:448:ILE:HB	1:C:460:PHE:CE1	2.39	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:90:TYR:CD1	1:C:496:ARG:HD2	2.38	0.57
1:C:227:VAL:HG11	1:C:230:MET:HE2	1.85	0.57
1:C:436:ALA:HB1	1:C:441:ILE:HG23	1.88	0.56
1:A:519:PHE:CE2	1:A:523:ILE:HD11	2.41	0.55
1:C:289:LEU:HD22	1:C:292:PHE:CD1	2.40	0.55
1:A:280:ALA:HA	1:A:326:ILE:HD11	1.89	0.55
3:A:702:FAD:O5B	3:A:702:FAD:H8A	2.07	0.55
1:C:464:LEU:HB2	1:C:480:PHE:CZ	2.41	0.55
1:A:419:PRO:HB3	1:C:206:MET:SD	2.47	0.55
1:C:411:HIS:NE2	3:C:602:FAD:HM82	2.22	0.55
1:A:464:LEU:HD11	1:A:476:SER:CB	2.35	0.54
1:A:473:ASN:O	1:A:477:ARG:HG3	2.08	0.54
1:C:359:ASP:N	1:C:359:ASP:OD1	2.41	0.54
1:C:439:LEU:CD1	1:C:482:HIS:CD2	2.91	0.54
1:A:413:TRP:CZ3	1:A:463:PRO:HB3	2.44	0.53
1:A:227:VAL:HG11	1:A:230:MET:CE	2.39	0.52
1:A:67:GLU:OE2	1:A:70:ARG:NH1	2.42	0.52
1:A:416:PRO:HD2	1:A:460:PHE:O	2.10	0.52
1:A:156:VAL:HG13	1:A:230:MET:HE1	1.91	0.52
1:A:161:ASP:OD2	1:A:496:ARG:HD3	2.10	0.51
1:C:300:GLN:HG2	1:C:305:MET:CG	2.43	0.49
1:C:170:GLN:HG2	1:C:171:PHE:CD2	2.48	0.48
1:C:215:ILE:O	1:C:218:ILE:HG12	2.13	0.48
1:A:448:ILE:HB	1:A:460:PHE:CE1	2.48	0.48
1:C:274:PRO:O	1:C:452:PRO:HG2	2.12	0.48
1:A:242:LEU:HD23	1:A:242:LEU:C	2.33	0.48
1:C:460:PHE:HE2	1:C:462:ILE:HD11	1.79	0.48
1:A:373:THR:OG1	1:A:376:GLN:HG3	2.14	0.47
1:C:146:ASP:OD1	1:C:165:GLY:HA3	2.14	0.47
1:C:531:GLY:HA3	1:C:542:PRO:CB	2.44	0.47
1:A:449:VAL:CG2	5:A:895:HOH:O	2.63	0.47
1:A:162:ARG:HD3	5:C:806:HOH:O	2.14	0.47
1:C:81:THR:O	3:C:602:FAD:H2B	2.15	0.46
1:A:526:ALA:HB2	1:C:526:ALA:HB2	1.98	0.46
1:C:286:LEU:CD2	5:C:878:HOH:O	2.64	0.46
1:C:289:LEU:HD13	1:C:292:PHE:HE1	1.80	0.46
1:C:156:VAL:HG13	1:C:230:MET:CE	2.46	0.46
1:C:243:LYS:HZ3	1:C:286:LEU:HD21	1.80	0.46
1:A:146:ASP:OD1	1:A:165:GLY:HA3	2.15	0.46
1:C:288:GLY:HA2	5:C:834:HOH:O	2.16	0.46
1:A:468:GLU:O	1:A:470:PRO:HD3	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:TYR:CD1	1:A:410:GLY:HA3	2.51	0.46
1:A:255:VAL:HA	1:A:422:GLY:HA3	1.98	0.45
1:A:438:ARG:HB2	5:A:944:HOH:O	2.15	0.45
1:A:90:TYR:O	1:A:499:PRO:HD3	2.16	0.44
1:A:286:LEU:HD22	1:A:382:TYR:CG	2.52	0.44
1:A:274:PRO:O	1:A:452:PRO:HG2	2.17	0.44
1:C:293:LEU:HA	1:C:293:LEU:HD23	1.76	0.44
1:C:448:ILE:HD12	1:C:460:PHE:CZ	2.52	0.44
1:C:275:ASP:OD2	1:C:286:LEU:HD22	2.17	0.43
1:C:277:ASN:HB3	1:C:285:SER:HB3	2.01	0.43
1:A:504:ASP:OD1	1:A:551:TYR:HE1	2.00	0.43
1:C:44:TYR:CD1	1:C:49:GLU:HB2	2.53	0.43
1:A:88:LEU:O	3:A:702:FAD:H9	2.18	0.43
1:C:251:TYR:OH	1:C:426:LEU:HD22	2.19	0.43
1:C:300:GLN:HG2	1:C:305:MET:HG3	2.01	0.43
1:A:435:GLU:HG3	1:A:486:VAL:HG21	2.01	0.42
1:C:368:TYR:HB3	1:C:372:LEU:HD21	2.01	0.42
1:C:198:MET:HB3	1:C:201:SER:HB2	2.01	0.42
1:C:547:ASP:OD1	1:C:550:ARG:NH1	2.52	0.41
1:C:460:PHE:CE2	1:C:462:ILE:HD11	2.55	0.41
1:A:170:GLN:HG2	1:A:171:PHE:CE2	2.56	0.41
1:A:242:LEU:C	1:A:242:LEU:CD2	2.89	0.41
1:C:230:MET:HE3	1:C:230:MET:HB2	1.97	0.41
1:C:289:LEU:HD13	1:C:292:PHE:CE1	2.56	0.41
1:A:39:ASP:HB3	1:A:42:SER:HB2	2.02	0.41
1:C:285:SER:HB2	1:C:286:LEU:H	1.65	0.41
1:A:251:TYR:OH	1:A:426:LEU:HD22	2.21	0.41
1:C:164:ALA:HA	1:C:172:ARG:O	2.20	0.40
1:C:255:VAL:HA	1:C:422:GLY:HA3	2.04	0.40
1:C:73:ASN:O	1:C:530:LYS:HD3	2.22	0.40
1:C:89:GLY:HA3	1:C:413:TRP:CZ3	2.57	0.40
1:C:87:ASN:HB2	3:C:602:FAD:O1A	2.22	0.40
1:C:370:LEU:HD23	1:C:372:LEU:HG	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	534/557 (96%)	526 (98%)	8 (2%)	0	100	100
1	C	527/557 (95%)	515 (98%)	12 (2%)	0	100	100
All	All	1061/1114 (95%)	1041 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	444/458 (97%)	438 (99%)	6 (1%)	62	56
1	C	441/458 (96%)	428 (97%)	13 (3%)	37	26
All	All	885/916 (97%)	866 (98%)	19 (2%)	48	38

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	161	ASP
1	A	242	LEU
1	A	438	ARG
1	A	441	ILE
1	A	449	VAL
1	A	552	LYS
1	C	20	ASN
1	C	154	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	161	ASP
1	C	178	HIS
1	C	286	LEU
1	C	289	LEU
1	C	292	PHE
1	C	359	ASP
1	C	394	PHE
1	C	444	ILE
1	C	449	VAL
1	C	453	SER
1	C	489	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	250	GLN
1	A	350	GLN
1	A	411	HIS
1	C	20	ASN
1	C	73	ASN
1	C	137	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ALA	A	701	-	5,5,5	1.02	0	6,6,6	2.89	2 (33%)
3	FAD	C	602	-	53,58,58	0.66	0	68,89,89	0.81	1 (1%)
4	GLY	C	601	-	4,4,4	1.54	1 (25%)	3,4,4	2.33	2 (66%)
3	FAD	A	702	-	53,58,58	0.70	1 (1%)	68,89,89	0.78	3 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALA	A	701	-	-	0/4/4/4	-
3	FAD	C	602	-	-	6/30/50/50	0/6/6/6
4	GLY	C	601	-	-	0/2/2/2	-
3	FAD	A	702	-	-	7/30/50/50	0/6/6/6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	601	GLY	OXT-C	-2.96	1.20	1.30
3	A	702	FAD	C8A-N7A	-2.02	1.31	1.34

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	ALA	OXT-C-O	-6.07	110.30	124.09
4	C	601	GLY	OXT-C-O	-3.26	115.17	123.30
2	A	701	ALA	OXT-C-CA	2.72	123.87	114.06
4	C	601	GLY	OXT-C-CA	2.38	122.93	113.45
3	C	602	FAD	C5A-C6A-N6A	2.25	123.77	120.35
3	A	702	FAD	C5A-C6A-N6A	2.18	123.66	120.35
3	A	702	FAD	C4-N3-C2	-2.13	121.71	125.64
3	A	702	FAD	O2P-P-O1P	2.01	122.19	112.24

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	702	FAD	O4'-C4'-C5'-O5'
3	C	602	FAD	O4'-C4'-C5'-O5'
3	C	602	FAD	C2'-C3'-C4'-O4'
3	A	702	FAD	C2'-C3'-C4'-O4'
3	A	702	FAD	C3'-C4'-C5'-O5'
3	C	602	FAD	C3'-C4'-C5'-O5'
3	A	702	FAD	C2'-C3'-C4'-C5'
3	C	602	FAD	C2'-C3'-C4'-C5'
3	C	602	FAD	C4'-C5'-O5'-P
3	A	702	FAD	C4'-C5'-O5'-P
3	A	702	FAD	O3'-C3'-C4'-O4'
3	C	602	FAD	O3'-C3'-C4'-O4'
3	A	702	FAD	O3'-C3'-C4'-C5'

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	602	FAD	4	0
3	A	702	FAD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	538/557 (96%)	0.28	28 (5%) 34 31	21, 34, 68, 110	0
1	C	533/557 (95%)	0.10	26 (4%) 36 33	21, 32, 60, 100	0
All	All	1071/1114 (96%)	0.19	54 (5%) 35 32	21, 32, 64, 110	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	286	LEU	6.2
1	C	394	PHE	5.7
1	A	441	ILE	5.5
1	C	293	LEU	5.1
1	C	292	PHE	5.0
1	C	287	ALA	4.6
1	A	460	PHE	4.3
1	C	289	LEU	4.2
1	A	439	LEU	4.1
1	C	439	LEU	4.0
1	A	436	ALA	4.0
1	A	443	LEU	3.9
1	C	436	ALA	3.9
1	C	460	PHE	3.9
1	A	413	TRP	3.7
1	C	410	GLY	3.6
1	A	294	ALA	3.5
1	C	291	GLU	3.5
1	C	288	GLY	3.5
1	C	396	ILE	3.5
1	C	441	ILE	3.4
1	A	438	ARG	3.1
1	C	444	ILE	3.1
1	A	395	ALA	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	440	GLY	2.9
1	C	393	THR	2.8
1	C	461	ILE	2.8
1	A	461	ILE	2.7
1	C	300	GLN	2.7
1	C	440	GLY	2.7
1	A	293	LEU	2.7
1	C	290	HIS	2.7
1	A	296	GLY	2.7
1	A	480	PHE	2.6
1	C	454	TRP	2.5
1	A	287	ALA	2.5
1	C	285	SER	2.5
1	A	437	ARG	2.4
1	C	438	ARG	2.4
1	C	411	HIS	2.4
1	A	394	PHE	2.4
1	C	443	LEU	2.3
1	A	326	ILE	2.3
1	A	297	PRO	2.3
1	C	480	PHE	2.3
1	A	302	PRO	2.2
1	A	292	PHE	2.2
1	A	435	GLU	2.1
1	A	454	TRP	2.1
1	A	414	PHE	2.1
1	A	152	TRP	2.1
1	A	288	GLY	2.1
1	A	286	LEU	2.1
1	A	412	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ALA	A	701	6/6	0.70	0.21	52,52,54,55	0
4	GLY	C	601	5/5	0.85	0.13	42,43,48,51	0
3	FAD	A	702	53/53	0.95	0.08	27,32,41,45	0
3	FAD	C	602	53/53	0.98	0.04	21,24,28,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.