

# Full wwPDB X-ray Structure Validation Report (i)

Jan 28, 2025 - 12:06 pm GMT

PDB ID	:	9H45
Title	:	Crystal Structure of Hfq V22A
Authors	:	McQuail, J.; Krepl, M.; Katsuya-Gaviria, K.; Tabib-Salazar, A.; Burchell, L.;
		Bischler, T.; Grafenhan, T.; Brear, P.; Sponer, J.; Luisi, B.
Deposited on	:	2024-10-17
Resolution	:	2.08  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R <sub>free</sub>	164625	7574(2.10-2.06)
Clashscore	180529	8325 (2.10-2.06)
Ramachandran outliers	177936	8271 (2.10-2.06)
Sidechain outliers	177891	8272 (2.10-2.06)
RSRZ outliers	164620	7574 (2.10-2.06)
RNA backbone	3690	1067 (2.46-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
			37%				
1	А	102	49%	14% •	36%		
			25%				
1	В	102	46%	15% •	38%		
			37%				
1	С	102	48%	13% •	38%		
			39%				
1	D	102	42%	17% •	40%		



Conti	nued fron	<i>i</i> previous	page				
Mol	Chain	Length	Quality of chain				
			35%				
1	Ε	102	43%	15% •	38%		
			28%				
1	F	102	47%	15% •	37%		
2	Q	18	28%	61%	11%		



## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Ato	$\mathbf{ms}$			ZeroOcc	AltConf	Trace
1	Λ	65	Total	С	Ν	Ο	$\mathbf{S}$	0	0	0
	05	517	332	91	93	1	0	0	0	
1	В	63	Total	С	Ν	Ο	S	0	0	0
1	D	05	502	324	88	89	1	0	0	0
1	С	C 63	Total	С	Ν	Ο	S	0	0	0
	U		502	324	88	89	1			
1	Л	61	Total	С	Ν	Ο	S	0	0	0
1	D		491	318	86	86	1	0		
1	F	63	Total	С	Ν	Ο	S	0	0	0
	05	502	324	88	89	1	0	0	0	
1 E	6.4	Total	С	Ν	Ο	S	0	2	0	
	Г	64	525	339	94	91	1		2	U

• Molecule 1 is a protein called RNA-binding protein Hfq.

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	22	ALA	VAL	engineered mutation	UNP P0A6X3
В	22	ALA	VAL	engineered mutation	UNP P0A6X3
С	22	ALA	VAL	engineered mutation	UNP P0A6X3
D	22	ALA	VAL	engineered mutation	UNP P0A6X3
Е	22	ALA	VAL	engineered mutation	UNP P0A6X3
F	22	ALA	VAL	engineered mutation	UNP P0A6X3

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Q	18	Total 396	C 180	N 90	0 108	Р 18	0	0	0

• Molecule 3 is water.



9H45	)

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	36	Total O 36 36	0	0
3	В	39	Total         O           39         39	0	0
3	С	31	Total         O           31         31	0	0
3	D	38	Total         O           38         38	0	0
3	Е	41	Total         O           41         41	0	0
3	F	43	Total         O           43         43	0	0
3	Q	54	Total O 54 54	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: RNA-binding protein Hfq





Chain Q:	28%	61%	11%
다 <b>이 아 아 아</b>	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	116.25Å 67.15Å 84.58Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $104.64^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{Posolution} \left( \overset{\circ}{\mathbf{A}} \right)$	56.24 - 2.08	Depositor
Resolution (A)	56.24 - 2.08	EDS
% Data completeness	99.0 (56.24-2.08)	Depositor
(in resolution range)	99.1 (56.24-2.08)	EDS
$R_{merge}$	0.51	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.62 (at 2.08 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
P. P.	0.274 , $0.323$	Depositor
$n, n_{free}$	0.290 , $0.330$	DCC
$R_{free}$ test set	1879 reflections $(4.98\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.1	Xtriage
Anisotropy	0.326	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34 , $44.4$	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	3717	wwPDB-VP
Average B, all atoms $(Å^2)$	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 12.40% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bo	ond angles
	Ullaili	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.29	0/526	0.54	0/713
1	В	0.33	0/511	0.57	0/693
1	С	0.47	0/511	0.73	0/693
1	D	0.55	0/499	0.70	0/675
1	Е	0.44	0/511	0.65	0/693
1	F	0.47	0/540	0.71	0/730
2	Q	0.40	0/449	1.10	5/698~(0.7%)
All	All	0.43	0/3547	0.74	5/4895~(0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1
1	Ε	0	2
All	All	0	3

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	Q	4	A	P-O3'-C3'	-7.62	110.55	119.70
2	Q	18	A	OP1-P-OP2	-7.25	108.72	119.60
2	Q	17	А	OP1-P-O3'	6.47	119.43	105.20
2	Q	13	А	P-O3'-C3'	-6.43	111.98	119.70
2	Q	10	А	P-O3'-C3'	-6.04	112.45	119.70

There are no chirality outliers.

All (3) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	D	19	ARG	Sidechain
1	Ε	16	ARG	Sidechain
1	Ε	66	ARG	Sidechain

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	517	0	539	14	0
1	В	502	0	526	7	0
1	С	502	0	526	10	0
1	D	491	0	515	13	0
1	Е	502	0	526	9	0
1	F	525	0	560	13	0
2	Q	396	0	199	2	0
3	А	36	0	0	3	0
3	В	39	0	0	1	1
3	С	31	0	0	1	0
3	D	38	0	0	4	0
3	Е	41	0	0	1	1
3	F	43	0	0	0	0
3	Q	54	0	0	1	0
All	All	3717	0	3391	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:15:LEU:HD13	1:F:20:VAL:HG13	1.47	0.94
1:F:68:VAL:HG23	1:F:68:VAL:O	1.93	0.69
1:A:7:LEU:HD13	1:C:45:LEU:HD21	1.77	0.67
1:F:15:LEU:HD13	1:F:20:VAL:CG1	2.25	0.64
1:A:7:LEU:HD23	1:C:40:ASP:HB3	1.85	0.59
1:E:56:LYS:HA	1:E:59:ILE:HD12	1.86	0.58
1:A:7:LEU:CD1	1:C:45:LEU:HD21	2.34	0.57



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:58:ALA:HB3	3:B:207:HOH:O	2.03	0.57
1:D:53:MET:HB3	1:F:62:VAL:HG13	1.88	0.56
1:C:38:SER:OG	1:C:45:LEU:HD23	2.07	0.54
1:B:16:ARG:O	1:B:19:ARG:HD3	2.08	0.53
1:A:40:ASP:HB2	3:A:214:HOH:O	2.08	0.52
3:C:202:HOH:O	1:E:43:VAL:HG13	2.09	0.51
2:Q:1:A:O5'	2:Q:18:A:O3'	2.29	0.50
1:A:24:ILE:HG12	1:A:62:VAL:HG22	1.94	0.49
1:D:53:MET:HB3	1:F:62:VAL:CG1	2.43	0.49
1:D:38:SER:HB2	1:F:7:LEU:HD21	1.95	0.49
1:E:40:ASP:OD1	1:E:43:VAL:HG22	2.12	0.49
1:D:15:LEU:HD12	1:D:36:ILE:CD1	2.43	0.49
1:B:37:GLU:HB3	1:B:47:LYS:HG3	1.94	0.49
1:D:43:VAL:HG22	3:D:208:HOH:O	2.12	0.49
1:D:53:MET:HE1	1:F:8:GLN:HA	1.94	0.48
1:B:61:THR:HG22	1:B:63:VAL:HG23	1.94	0.48
3:D:208:HOH:O	1:F:8:GLN:HB2	2.14	0.47
1:A:15:LEU:HD23	1:A:20:VAL:CG1	2.44	0.47
1:A:58:ALA:HB3	3:A:206:HOH:O	2.15	0.46
1:E:7:LEU:HD21	1:F:45:LEU:HD12	1.96	0.46
1:A:7:LEU:HD11	1:C:43:VAL:CG2	2.46	0.46
1:A:8:GLN:CD	3:A:209:HOH:O	2.55	0.45
1:A:9:ASP:HB2	1:A:10:PRO:HD3	1.99	0.45
1:F:68:VAL:O	1:F:68:VAL:CG2	2.64	0.45
1:C:45:LEU:N	1:C:45:LEU:HD22	2.32	0.45
1:A:15:LEU:HD21	1:A:64:PRO:HG3	2.00	0.44
1:C:32:LEU:HB3	1:C:46:LEU:CD2	2.48	0.43
1:D:48:ASN:C	3:D:201:HOH:O	2.57	0.43
1:A:7:LEU:CD2	1:C:40:ASP:HB3	2.47	0.43
1:B:31:LYS:HD3	3:Q:117:HOH:O	2.18	0.43
1:D:25:TYR:CD2	2:Q:12:A:C4	3.07	0.42
1:E:47:LYS:CE	1:E:51:SER:HB2	2.49	0.42
1:F:40:ASP:OD1	1:F:43:VAL:HG22	2.20	0.42
1:F:56:LYS:HA	1:F:59:ILE:HD12	2.01	0.42
1:D:15:LEU:CD2	1:D:68:VAL:HG21	2.50	0.42
1:D:56:LYS:HA	1:D:59:ILE:HD12	2.01	0.42
1:A:5:GLN:CG	1:A:6:SER:H	2.33	0.41
1:E:49:THR:HG22	1:E:50:VAL:HG13	2.01	0.41
1:F:28:ASN:OD1	1:F:30:ILE:HD13	2.20	0.41
1:D:20:VAL:HG13	1:D:66:ARG:CZ	2.50	0.41
1:B:52:GLN:OE1	1:D:61:THR:HG21	2.20	0.41



9H45
------

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:23:SER:HB2	1:E:63:VAL:HG13	2.03	0.41
1:D:19:ARG:NH1	3:D:203:HOH:O	2.54	0.41
1:E:11:PHE:O	1:E:15:LEU:HD12	2.21	0.41
1:A:8:GLN:NE2	1:C:55:TYR:OH	2.54	0.41
1:B:7:LEU:C	1:B:7:LEU:HD12	2.41	0.41
1:E:8:GLN:N	3:E:201:HOH:O	2.54	0.41
1:C:47:LYS:HE3	1:C:50:VAL:C	2.41	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:211:HOH:O	3:E:229:HOH:O[1_545]	1.96	0.24

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	63/102~(62%)	60~(95%)	2(3%)	1 (2%)	8 4
1	В	61/102~(60%)	58~(95%)	2(3%)	1 (2%)	8 4
1	С	61/102~(60%)	59 (97%)	1 (2%)	1 (2%)	8 4
1	D	57/102~(56%)	54 (95%)	3 (5%)	0	100 100
1	Е	61/102~(60%)	58 (95%)	2(3%)	1 (2%)	8 4
1	F	64/102~(63%)	61~(95%)	3~(5%)	0	100 100
All	All	367/612~(60%)	350 (95%)	13 (4%)	4 (1%)	12 7

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	48	ASN
	<i>a</i>	-	-



 $Continued \ from \ previous \ page...$ 

Mol	Chain	Res	Type
1	А	48	ASN
1	В	48	ASN
1	Е	48	ASN

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	60/88~(68%)	58~(97%)	2(3%)	33 34
1	В	58/88~(66%)	52~(90%)	6 (10%)	6 3
1	С	58/88~(66%)	53 (91%)	5 (9%)	8 5
1	D	57/88~(65%)	52 (91%)	5 (9%)	8 5
1	Е	58/88~(66%)	52 (90%)	6 (10%)	6 3
1	F	61/88~(69%)	56~(92%)	5 (8%)	9 6
All	All	352/528~(67%)	323~(92%)	29 (8%)	10 6

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	6	SER
1	А	66	ARG
1	В	9	ASP
1	В	17	ARG
1	В	18	GLU
1	В	20	VAL
1	В	48	ASN
1	В	66	ARG
1	С	9	ASP
1	С	16	ARG
1	С	20	VAL
1	С	23	SER
1	С	47	LYS
1	D	7	LEU
1	D	18	GLU



Mol	Chain	Res	Type
1	D	19	ARG
1	D	40	ASP
1	D	60	SER
1	Е	23	SER
1	Е	30	ILE
1	Ε	37	GLU
1	Е	40	ASP
1	Ε	47	LYS
1	Ε	66	ARG
1	F	17[A]	ARG
1	F	17[B]	ARG
1	F	30	ILE
1	F	37	GLU
1	F	49	THR

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	В	48	ASN

#### 5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	Q	17/18~(94%)	7 (41%)	0

All (7) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	Q	2	А
2	Q	5	А
2	Q	8	А
2	Q	9	А
2	Q	11	А
2	Q	14	А
2	Q	17	А

There are no RNA pucker outliers to report.



### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry (i)

There are no ligands in this entry.

#### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

**Warning**: The R factor obtained from EDS is 0.3321, which does not match the depositor's R factor of 0.274. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2		$OWAB(Å^2)$	Q<0.9
1	А	65/102~(63%)	2.25	38 (58%) 0 (	0	10, 26, 47, 79	0
1	В	63/102~(61%)	2.11	26 (41%) 1	1	8, 21, 43, 64	0
1	С	63/102~(61%)	2.45	38 (60%) 0	0	6, 24, 41, 54	0
1	D	61/102~(59%)	2.43	40 (65%) 0	0	4,25,41,55	0
1	Ε	63/102~(61%)	2.27	36 (57%) 0	0	10, 23, 44, 65	0
1	F	64/102~(62%)	2.12	29~(45%) 1	1	9,24,44,71	2 (3%)
2	Q	18/18~(100%)	1.09	0 100 100		15, 24, 29, 30	0
All	All	397/630~(63%)	2.22	207 (52%) 0	0	4, 24, 47, 79	2(0%)

All (207) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	68	VAL	6.7
1	С	68	VAL	6.7
1	А	68	VAL	6.0
1	D	49	THR	5.9
1	Ε	6	SER	5.9
1	D	15	LEU	5.8
1	Е	68	VAL	5.8
1	F	17[A]	ARG	5.7
1	F	68	VAL	5.6
1	В	67	PRO	5.4
1	D	68	VAL	5.3
1	А	69	SER	5.2
1	С	63	VAL	5.1
1	С	7	LEU	4.9
1	F	39	PHE	4.9



Mol	Chain	Res	Type	RSRZ
1	С	67	PRO	4.7
1	А	7	LEU	4.6
1	С	59	ILE	4.6
1	С	49	THR	4.6
1	Е	7	LEU	4.5
1	А	14	ALA	4.2
1	С	37	GLU	4.1
1	С	14	ALA	4.1
1	Е	49	THR	4.1
1	С	47	LYS	4.1
1	D	30	ILE	4.0
1	А	67	PRO	4.0
1	Е	67	PRO	4.0
1	D	67	PRO	3.8
1	D	48	ASN	3.8
1	F	7	LEU	3.8
1	С	22	ALA	3.8
1	С	20	VAL	3.7
1	F	62	VAL	3.7
1	В	46	LEU	3.7
1	С	45	LEU	3.7
1	А	36	ILE	3.7
1	D	7	LEU	3.6
1	Е	60	SER	3.6
1	F	49	THR	3.6
1	Е	38	SER	3.6
1	А	5	GLN	3.6
1	А	30	ILE	3.5
1	В	7	LEU	3.5
1	Е	14	ALA	3.5
1	F	30	ILE	3.5
1	F	36	ILE	3.5
1	В	62	VAL	3.5
1	D	36	ILE	3.5
1	В	47	LYS	3.4
1	D	51	SER	3.4
1	А	55	TYR	3.4
1	D	47	LYS	3.4
1	D	54	VAL	3.4
1	В	49	THR	3.4
1	С	34	GLY	3.4
1	D	17	ARG	3.4



Mol	Chain	$\operatorname{Res}$	Type	RSRZ
1	F	63	VAL	3.3
1	В	51	SER	3.3
1	В	41	GLN	3.3
1	Е	47	LYS	3.3
1	Е	63	VAL	3.2
1	С	15	LEU	3.2
1	Е	10	PRO	3.2
1	С	43	VAL	3.1
1	С	51	SER	3.1
1	F	51	SER	3.1
1	F	5	GLN	3.1
1	С	18	GLU	3.1
1	С	62	VAL	3.1
1	D	35	GLN	3.1
1	Е	27	VAL	3.1
1	А	10	PRO	3.0
1	А	62	VAL	3.0
1	С	54	VAL	3.0
1	В	15	LEU	3.0
1	Е	30	ILE	3.0
1	С	17	ARG	3.0
1	С	6	SER	3.0
1	А	51	SER	3.0
1	D	63	VAL	2.9
1	D	53	MET	2.9
1	А	20	VAL	2.9
1	В	20	VAL	2.9
1	D	62	VAL	2.9
1	Е	46	LEU	2.9
1	Е	48	ASN	2.9
1	D	22	ALA	2.9
1	Е	51	SER	2.9
1	D	66	ARG	2.9
1	D	25	TYR	2.9
1	A	37	GLU	2.9
1	A	46	LEU	2.8
1	В	39	PHE	2.8
1	В	38	SER	2.8
1	С	38	SER	2.8
1	Е	32	LEU	2.8
1	В	22	ALA	2.8
1	А	33	GLN	2.8



Mol	Chain	Res	Type	RSRZ
1	F	15	LEU	2.8
1	D	65	SER	2.8
1	В	48	ASN	2.8
1	Е	37	GLU	2.8
1	F	48	ASN	2.8
1	А	24	ILE	2.8
1	А	66	ARG	2.8
1	F	45	LEU	2.8
1	А	63	VAL	2.8
1	D	13	ASN	2.7
1	А	18	GLU	2.7
1	D	45	LEU	2.7
1	Е	22	ALA	2.7
1	С	36	ILE	2.7
1	А	65	SER	2.7
1	Е	59	ILE	2.7
1	А	32	LEU	2.7
1	С	16	ARG	2.7
1	F	35	GLN	2.7
1	В	24	ILE	2.7
1	В	55	TYR	2.6
1	С	50	VAL	2.6
1	D	20	VAL	2.6
1	F	27	VAL	2.6
1	D	33	GLN	2.6
1	А	22	ALA	2.6
1	А	49	THR	2.6
1	С	39	PHE	2.6
1	В	66	ARG	2.6
1	Е	64	PRO	2.5
1	А	48	ASN	2.5
1	D	55	TYR	2.5
1	D	12	LEU	2.5
1	A	17	ARG	2.5
1	В	13	ASN	2.5
1	С	42	PHE	2.4
1	F	22	ALA	2.4
1	С	61	THR	2.4
1	F	67	PRO	2.4
1	F	12	LEU	2.4
1	F	32	LEU	2.4
1	В	14	ALA	2.4



Mol	Chain	Res	Type	RSRZ
1	С	48	ASN	2.4
1	А	47	LYS	2.4
1	F	47	LYS	2.4
1	В	64	PRO	2.4
1	F	44	ILE	2.4
1	С	8	GLN	2.4
1	D	8	GLN	2.4
1	Е	58	ALA	2.4
1	D	64	PRO	2.4
1	Е	65	SER	2.4
1	В	30	ILE	2.4
1	D	50	VAL	2.3
1	Е	54	VAL	2.3
1	Е	9	ASP	2.3
1	А	39	PHE	2.3
1	Е	55	TYR	2.3
1	С	65	SER	2.3
1	Е	20	VAL	2.3
1	D	42	PHE	2.3
1	С	55	TYR	2.3
1	D	44	ILE	2.3
1	С	33	GLN	2.3
1	А	28	ASN	2.3
1	Е	42	PHE	2.3
1	F	13	ASN	2.3
1	Е	45	LEU	2.3
1	Е	13	ASN	2.2
1	Е	66	ARG	2.2
1	А	25	TYR	2.2
1	Е	26	LEU	2.2
1	F	18	GLU	2.2
1	F	64	PRO	2.2
1	А	34	GLY	2.2
1	D	43	VAL	2.2
1	A	41	GLN	2.2
1	С	35	GLN	2.2
1	D	59	ILE	2.2
1	С	9	ASP	2.2
1	D	38	SER	2.2
1	С	13	ASN	2.2
1	Е	62	VAL	2.2
1	А	15	LEU	2.1



Mol	Chain	Res	Type	RSRZ
1	D	46	LEU	2.1
1	Е	25	TYR	2.1
1	С	27	VAL	2.1
1	F	20	VAL	2.1
1	А	59	ILE	2.1
1	С	30	ILE	2.1
1	Е	44	ILE	2.1
1	В	35	GLN	2.1
1	В	43	VAL	2.1
1	В	19	ARG	2.1
1	А	44	ILE	2.1
1	F	43	VAL	2.1
1	А	38	SER	2.1
1	А	35	GLN	2.1
1	А	52	GLN	2.1
1	F	66	ARG	2.0
1	В	23	SER	2.0
1	D	60	SER	2.0
1	D	26	LEU	2.0
1	D	32	LEU	2.0
1	D	61	THR	2.0
1	Е	39	PHE	2.0
1	D	28	ASN	2.0
1	F	14	ALA	2.0

Continued from previous page...

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

There are no ligands in this entry.

#### 6.5 Other polymers (i)

There are no such residues in this entry.

