

wwPDB X-ray Structure Validation Summary Report (i)

Dec 12, 2024 – 09:32 pm GMT

PDB ID : 9H8Q

Title: Eugenol Oxidase (EUGO) from Rhodococcus jostii RHA1, mutant DTT-

T425G

Authors: Rozeboom, H.J.; Fraaije, M.W.

Deposited on : 2024-10-29

Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.40

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	164625	4274 (1.60-1.60)
Clashscore	180529	4682 (1.60-1.60)
Ramachandran outliers	177936	4583 (1.60-1.60)
Sidechain outliers	177891	4582 (1.60-1.60)
RSRZ outliers	164620	4272 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	526	92%	7%
1	В	526	94%	6%
1	С	526	94%	5% •
1	D	526	94%	5% •
1	Е	526	92%	7%



Continued from previous page...

	J	1	<i>5</i>	
Mol	Chain	Length	Quality of chain	
1	F	526	94% 5% •	•
1	G	526	93% 6% •	
1	Н	526	94% 5%	



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 36661 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Probable vanillyl-alcohol oxidase.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	A	525	Total	С	N	О	S	0	2	0
1	A	323	4146	2642	702	776	26	U	2	
1	В	525	Total	С	N	О	S	0	1	0
1	Б	525	4138	2637	701	775	25	U	1	
1	С	525	Total	С	N	О	S	0	3	0
1		323	4154	2646	703	779	26	Ü	3	
1	D	525	Total	С	N	О	S	0	0	0
1	D		4130	2632	700	774	24	O	0	
1	Е	525	Total	С	N	О	S	0	2	0
1	l L	323	4146	2642	702	776	26	U		
1	F	525	Total	С	N	O	S	0	0	0
1	I.	323	4130	2632	700	774	24	U	0	
1	G	525	Total	С	N	О	S	0	1	0
1	G	323	4138	2638	701	775	24	U	1	
1	Н	525	Total	С	N	О	S	0	0	0
1	П	525	4130	2632	700	774	24	U	0	

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	81	HIS	SER	engineered mutation	UNP Q0SBK1
A	394	VAL	SER	engineered mutation	UNP Q0SBK1
A	423	MET	ALA	engineered mutation	UNP Q0SBK1
A	425	GLY	GLN	engineered mutation	UNP Q0SBK1
A	427	THR	ILE	engineered mutation	UNP Q0SBK1
A	434	TYR	HIS	engineered mutation	UNP Q0SBK1
A	445	ASP	ILE	engineered mutation	UNP Q0SBK1
A	518	PRO	SER	engineered mutation	UNP Q0SBK1
В	81	HIS	SER	engineered mutation	UNP Q0SBK1
В	394	VAL	SER	engineered mutation	UNP Q0SBK1
В	423	MET	ALA	engineered mutation	UNP Q0SBK1
В	425	GLY	GLN	engineered mutation	UNP Q0SBK1
В	427	THR	ILE	engineered mutation	UNP Q0SBK1



Continued from previous page...


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	434	TYR	HIS	engineered mutation	UNP Q0SBK1
В	445	ASP	ILE	engineered mutation	UNP Q0SBK1
В	518	PRO	SER	engineered mutation	UNP Q0SBK1
С	81	HIS	SER	engineered mutation	UNP Q0SBK1
С	394	VAL	SER	engineered mutation	UNP Q0SBK1
С	423	MET	ALA	engineered mutation	UNP Q0SBK1
С	425	GLY	GLN	engineered mutation	UNP Q0SBK1
С	427	THR	ILE	engineered mutation	UNP Q0SBK1
С	434	TYR	HIS	engineered mutation	UNP Q0SBK1
С	445	ASP	ILE	engineered mutation	UNP Q0SBK1
С	518	PRO	SER	engineered mutation	UNP Q0SBK1
D	81	HIS	SER	engineered mutation	UNP Q0SBK1
D	394	VAL	SER	engineered mutation	UNP Q0SBK1
D	423	MET	ALA	engineered mutation	UNP Q0SBK1
D	425	GLY	GLN	engineered mutation	UNP Q0SBK1
D	427	THR	ILE	engineered mutation	UNP Q0SBK1
D	434	TYR	HIS	engineered mutation	UNP Q0SBK1
D	445	ASP	ILE	engineered mutation	UNP Q0SBK1
D	518	PRO	SER	engineered mutation	UNP Q0SBK1
Е	81	HIS	SER	engineered mutation	UNP Q0SBK1
Е	394	VAL	SER	engineered mutation	UNP Q0SBK1
Е	423	MET	ALA	engineered mutation	UNP Q0SBK1
Е	425	GLY	GLN	engineered mutation	UNP Q0SBK1
Е	427	THR	ILE	engineered mutation	UNP Q0SBK1
Е	434	TYR	HIS	engineered mutation	UNP Q0SBK1
Е	445	ASP	ILE	engineered mutation	UNP Q0SBK1
Е	518	PRO	SER	engineered mutation	UNP Q0SBK1
F	81	HIS	SER	engineered mutation	UNP Q0SBK1
F	394	VAL	SER	engineered mutation	UNP Q0SBK1
F	423	MET	ALA	engineered mutation	UNP Q0SBK1
F	425	GLY	GLN	engineered mutation	UNP Q0SBK1
F	427	THR	ILE	engineered mutation	UNP Q0SBK1
F	434	TYR	HIS	engineered mutation	UNP Q0SBK1
F	445	ASP	ILE	engineered mutation	UNP Q0SBK1
F	518	PRO	SER	engineered mutation	UNP Q0SBK1
G	81	HIS	SER	engineered mutation	UNP Q0SBK1
G	394	VAL	SER	engineered mutation	UNP Q0SBK1
G	423	MET	ALA	engineered mutation	UNP Q0SBK1
G	425	GLY	GLN	engineered mutation	UNP Q0SBK1
G	427	THR	ILE	engineered mutation	UNP Q0SBK1
G	434	TYR	HIS	engineered mutation	UNP Q0SBK1
G	445	ASP	ILE	engineered mutation	UNP Q0SBK1



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
G	518	PRO	SER	engineered mutation	UNP Q0SBK1
Н	81	HIS	SER	engineered mutation	UNP Q0SBK1
Н	394	VAL	SER	engineered mutation	UNP Q0SBK1
Н	423	MET	ALA	engineered mutation	UNP Q0SBK1
Н	425	GLY	GLN	engineered mutation	UNP Q0SBK1
Н	427	THR	ILE	engineered mutation	UNP Q0SBK1
Н	434	TYR	HIS	engineered mutation	UNP Q0SBK1
Н	445	ASP	ILE	engineered mutation	UNP Q0SBK1
Н	518	PRO	SER	engineered mutation	UNP Q0SBK1

 \bullet Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2).$




Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	A	1	Total	С	N	О	Р	0	0
	Λ	1	53	27	9	15	2	O	0
2	В	1	Total	С	N	О	Р	0	0
	Ъ	1	53	27	9	15	2		0
2	С	1	Total	С	N	О	Р	0	0
2		1	53	27	9	15	2		
2	D	1	Total	С	N	О	Р	0	0
	ט	1	53	27	9	15	2	U	0
2	E	1	Total	С	N	О	Р	0	0
		1	53	27	9	15	2	0	0
2	F	1	Total	С	N	О	Р	0	0
	I'	1	53	27	9	15	2	U	U



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	G	1	Total	С	N	О	Р	0	0
	G	1	53	27	9	15	2	U	
2	Н	1	Total	С	N	О	Р	0	0
	Π	1	53	27	9	15	2		


• Molecule 3 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYL AMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: B3P) (formula: $C_{11}H_{26}N_2O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	Λ	1	Total	С	N	О	0	0	
)	3 A	1	19	11	2	6			
3	Λ	1	Total	С	N	О	0	0	
3	3 A	1	19	11	2	6	U		
3	F	1	Total	С	N	О	0	0	
3	I'	1	19	11	2	6	U		
3	3 F	1	Total	С	N	О	0	0	
3	Г	1	19	11	2	6	U	U	

• Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 8 6 2	0	0
4	В	1	Total C O 8 6 2	0	0
4	С	1	Total C O 8 6 2	0	0
4	D	1	Total C O 8 6 2	0	0
4	F	1	Total C O 8 6 2	0	0
4	Н	1	Total C O 8 6 2	0	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
5	В	1	Total Cl 1 1	0	0
5	F	1	Total Cl 1 1	0	0

• Molecule 6 is water.

\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
6	A	392	Total O 392 392	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	371	Total O 371 371	0	0
6	С	374	Total O 374 374	0	0
6	D	389	Total O 389 389	0	0
6	Е	420	Total O 420 420	0	0
6	F	385	Total O 385 385	0	0
6	G	329	Total O 329 329	0	0
6	Н	339	Total O 339 339	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Probable vanillyl-alcohol oxidase




• Molecule 1: Probable vanillyl-alcohol oxidase



• Molecule 1: Probable vanillyl-alcohol oxidase




• Molecule 1: Probable vanillyl-alcohol oxidase



• Molecule 1: Probable vanillyl-alcohol oxidase







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	90.44Å 110.20Å 117.54Å	Donositon
a, b, c, α , β , γ	89.87° 89.34° 68.51°	Depositor
Resolution (Å)	102.75 - 1.60	Depositor
rtesolution (A)	102.75 - 1.60	EDS
% Data completeness	93.2 (102.75-1.60)	Depositor
(in resolution range)	93.2 (102.75-1.60)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.58 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.177 , 0.198	Depositor
	0.188 , 0.207	DCC
R_{free} test set	25315 reflections $(4.88%)$	wwPDB-VP
Wilson B-factor (Å ²)	17.0	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 24.3	EDS
L-test for twinning ²	$< L > = 0.48, < L^2> = 0.31$	Xtriage
	0.020 for h,h-k,-l	
Estimated twinning fraction	0.137 for -h,-k,l	Xtriage
	$0.019 ext{ for -h,-h+k,-l}$	
F_o, F_c correlation	0.96	EDS
Total number of atoms	36661	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, FAD, CL, B3P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	В	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.48	0/4255	0.85	12/5772~(0.2%)
1	В	0.47	0/4247	0.80	5/5762 (0.1%)
1	С	0.46	0/4263	0.82	8/5783 (0.1%)
1	D	0.51	0/4239	0.85	$12/5752 \ (0.2\%)$
1	Е	0.51	0/4255	0.86	$13/5772 \ (0.2\%)$
1	F	0.48	0/4239	0.84	9/5752~(0.2%)
1	G	0.43	0/4247	0.79	$6/5763 \; (0.1\%)$
1	Н	0.46	0/4239	0.81	$10/5752 \ (0.2\%)$
All	All	0.48	0/33984	0.83	75/46108 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	В	0	2
1	С	0	2
1	D	0	2
1	Е	0	2
1	F	0	2
1	G	0	3
1	Н	0	1
All	All	0	16

There are no bond length outliers.

The worst 5 of 75 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	С	433	MET	CG-SD-CE	-12.34	80.46	100.20
1	G	176	MET	CG-SD-CE	10.52	117.03	100.20
1	A	423	MET	CG-SD-CE	-10.51	83.39	100.20
1	Н	109	ARG	NE-CZ-NH2	-9.70	115.45	120.30
1	G	109	ARG	NE-CZ-NH2	-9.39	115.61	120.30

There are no chirality outliers.

5 of 16 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	112	ARG	Sidechain
1	A	348	ARG	Sidechain
1	В	20	ARG	Sidechain
1	В	348	ARG	Sidechain
1	С	13	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4146	0	4012	22	0
1	В	4138	0	4004	20	0
1	С	4154	0	4015	22	0
1	D	4130	0	3996	12	0
1	Е	4146	0	4012	24	0
1	F	4130	0	3996	16	0
1	G	4138	0	4006	28	0
1	Н	4130	0	3996	13	0
2	A	53	0	30	1	0
2	В	53	0	30	6	0
2	С	53	0	31	7	0
2	D	53	0	30	1	0
2	Е	53	0	30	6	0
2	F	53	0	30	1	0
2	G	53	0	31	7	0
2	Н	53	0	30	1	0
3	A	38	0	52	0	0
3	F	38	0	52	1	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	8	0	14	1	0
4	В	8	0	14	2	0
4	С	8	0	14	0	0
4	D	8	0	14	0	0
4	F	8	0	14	0	0
4	Н	8	0	14	0	0
5	В	1	0	0	0	0
5	F	1	0	0	0	0
6	A	392	0	0	5	0
6	В	371	0	0	6	0
6	С	374	0	0	2	0
6	D	389	0	0	6	0
6	Ε	420	0	0	6	0
6	F	385	0	0	9	0
6	G	329	0	0	4	0
6	Н	339	0	0	4	0
All	All	36661	0	32467	164	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 164 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:E:390:HIS:NE2	2:E:600:FAD:C8M	1.68	1.56
1:B:390:HIS:NE2	2:B:601:FAD:C8M	1.69	1.56
1:C:390:HIS:NE2	2:C:600:FAD:C8M	1.74	1.47
1:G:390:HIS:NE2	2:G:600:FAD:C8M	1.79	1.41
1:G:390:HIS:NE2	2:G:600:FAD:HM82	1.02	1.34

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	525/526~(100%)	515 (98%)	10 (2%)	0	100	100
1	В	$524/526 \; (100\%)$	512 (98%)	12 (2%)	0	100	100
1	С	$526/526 \; (100\%)$	514 (98%)	12 (2%)	0	100	100
1	D	523/526~(99%)	512 (98%)	11 (2%)	0	100	100
1	E	$525/526 \; (100\%)$	514 (98%)	11 (2%)	0	100	100
1	F	523/526~(99%)	511 (98%)	12 (2%)	0	100	100
1	G	524/526 (100%)	512 (98%)	12 (2%)	0	100	100
1	Н	523/526 (99%)	513 (98%)	10 (2%)	0	100	100
All	All	4193/4208 (100%)	4103 (98%)	90 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	435/434 (100%)	429 (99%)	6 (1%)	62 43
1	В	434/434 (100%)	427 (98%)	7 (2%)	58 37
1	С	436/434 (100%)	428 (98%)	8 (2%)	54 31
1	D	433/434 (100%)	426 (98%)	7 (2%)	58 37
1	E	$435/434 \ (100\%)$	430 (99%)	5 (1%)	70 53
1	F	433/434 (100%)	426 (98%)	7 (2%)	58 37
1	G	434/434 (100%)	427 (98%)	7 (2%)	58 37
1	Н	433/434 (100%)	427 (99%)	6 (1%)	62 43
All	All	3473/3472 (100%)	3420 (98%)	53 (2%)	60 39

5 of 53 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	Е	154	TRP
1	F	273	ASN
1	Н	216	ASP
1	Е	198	LEU
1	Е	471	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	256	GLN
1	Н	205	GLN
1	F	67	GLN
1	Н	419	ASN
1	G	419	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 20 ligands modelled in this entry, 2 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Во	ond leng	ths	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	F	601	-	53,58,58	0.83	1 (1%)	68,89,89	0.98	2 (2%)
3	B3P	A	601	-	18,18,18	0.36	0	21,23,23	0.80	1 (4%)
2	FAD	D	600	-	53,58,58	0.77	1 (1%)	68,89,89	1.07	4 (5%)
2	FAD	С	600	-	53,58,58	0.74	1 (1%)	68,89,89	0.75	0
4	MPD	С	601	-	7,7,7	0.21	0	9,10,10	0.56	0
2	FAD	Н	602	-	53,58,58	0.69	0	68,89,89	0.95	4 (5%)
3	ВЗР	A	602	-	18,18,18	0.21	0	21,23,23	0.82	1 (4%)
3	ВЗР	F	604	-	18,18,18	0.23	0	21,23,23	1.04	1 (4%)
2	FAD	G	600	-	53,58,58	0.67	0	68,89,89	0.82	1 (1%)
3	B3P	F	602	-	18,18,18	0.20	0	21,23,23	0.80	0
4	MPD	D	601	-	7,7,7	0.20	0	9,10,10	0.40	0
2	FAD	A	600	_	53,58,58	0.81	1 (1%)	68,89,89	0.85	3 (4%)
4	MPD	Н	601	-	7,7,7	0.25	0	9,10,10	0.34	0
4	MPD	A	603	-	7,7,7	0.30	0	9,10,10	0.42	0
2	FAD	Е	600	-	53,58,58	0.65	0	68,89,89	0.83	2 (2%)
4	MPD	F	603	-	7,7,7	0.21	0	9,10,10	0.62	0
2	FAD	В	601	-	53,58,58	0.70	0	68,89,89	1.00	5 (7%)
4	MPD	В	602	-	7,7,7	0.35	0	9,10,10	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	F	601	-	-	4/30/50/50	0/6/6/6
3	ВЗР	A	601	-	-	3/28/28/28	-
2	FAD	D	600	-	-	3/30/50/50	0/6/6/6
2	FAD	С	600	-	-	6/30/50/50	0/6/6/6
4	MPD	С	601	-	-	1/5/5/5	-
2	FAD	Н	602	-	-	7/30/50/50	0/6/6/6
3	ВЗР	A	602	-	-	0/28/28/28	-
3	ВЗР	F	604	-	-	9/28/28/28	-
2	FAD	G	600	-	-	5/30/50/50	0/6/6/6
3	ВЗР	F	602	-	-	3/28/28/28	-
4	MPD	D	601	-	-	0/5/5/5	-
2	FAD	A	600	-	-	4/30/50/50	0/6/6/6
4	MPD	Н	601	-	-	0/5/5/5	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	A	603	-	-	0/5/5/5	-
2	FAD	${ m E}$	600	-	-	5/30/50/50	0/6/6/6
4	MPD	F	603	-	-	0/5/5/5	-
2	FAD	В	601	-	-	5/30/50/50	0/6/6/6
4	MPD	В	602	-	-	1/5/5/5	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
2	D	600	FAD	C1'-C2'	2.49	1.56	1.52
2	С	600	FAD	C8A-N7A	-2.30	1.30	1.34
2	A	600	FAD	C8A-N7A	-2.22	1.30	1.34
2	F	601	FAD	O4B-C1B	2.13	1.44	1.41

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
2	В	601	FAD	O2'-C2'-C3'	-4.04	99.27	109.10
2	D	600	FAD	P-O3P-PA	-3.45	120.99	132.83
2	D	600	FAD	O2'-C2'-C3'	-3.42	100.79	109.10
2	F	601	FAD	P-O3P-PA	-3.15	122.02	132.83
2	Е	600	FAD	P-O3P-PA	-3.00	122.52	132.83

There are no chirality outliers.

5 of 56 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	602	ВЗР	N1-C4-C7-O6
3	F	602	ВЗР	C5-C4-C7-O6
3	F	602	ВЗР	C6-C4-C7-O6
3	F	604	ВЗР	C6-C4-C5-O4
3	F	604	ВЗР	N1-C4-C7-O6

There are no ring outliers.

11 monomers are involved in 34 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	601	FAD	1	0
2	D	600	FAD	1	0
2	С	600	FAD	7	0





Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Н	602	FAD	1	0
3	F	604	ВЗР	1	0
2	G	600	FAD	7	0
2	A	600	FAD	1	0
4	A	603	MPD	1	0
2	Е	600	FAD	6	0
2	В	601	FAD	6	0
4	В	602	MPD	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#	∤RSR	Z>2	$OWAB(A^2)$	Q < 0.9
1	A	525/526~(99%)	-1.30	0	100	100	10, 18, 31, 56	2 (0%)
1	В	525/526~(99%)	-1.28	0	100	100	11, 18, 33, 63	1 (0%)
1	С	525/526~(99%)	-1.25	0	100	100	10, 19, 34, 59	3 (0%)
1	D	525/526~(99%)	-1.31	0	100	100	10, 16, 32, 60	0
1	E	525/526~(99%)	-1.32	0	100	100	10, 17, 29, 66	2 (0%)
1	F	525/526~(99%)	-1.28	0	100	100	10, 17, 31, 71	0
1	G	525/526~(99%)	-1.20	0	100	100	11, 22, 40, 61	1 (0%)
1	Н	525/526 (99%)	-1.26	0	100	100	12, 18, 34, 73	0
All	All	4200/4208~(99%)	-1.27	0	100	100	10, 18, 34, 73	9 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ m \AA}^2)$	Q<0.9
3	ВЗР	A	602	19/19	0.97	0.05	37,47,52,53	0
3	ВЗР	F	602	19/19	0.97	0.06	30,35,43,51	0
3	ВЗР	A	601	19/19	0.98	0.04	26,31,37,38	0
3	ВЗР	F	604	19/19	0.98	0.05	22,27,39,42	0
4	MPD	A	603	8/8	0.98	0.06	31,33,35,40	0
4	MPD	С	601	8/8	0.98	0.05	31,32,34,35	0
4	MPD	Н	601	8/8	0.98	0.05	27,29,32,33	0
4	MPD	В	602	8/8	0.99	0.04	29,30,32,33	0
2	FAD	E	600	53/53	0.99	0.02	11,13,16,20	0
4	MPD	D	601	8/8	0.99	0.04	25,28,31,32	0
4	MPD	F	603	8/8	0.99	0.05	27,29,32,33	0
2	FAD	С	600	53/53	0.99	0.02	12,14,17,18	0
2	FAD	G	600	53/53	1.00	0.02	12,16,19,22	0
2	FAD	Н	602	53/53	1.00	0.02	9,11,14,18	0
2	FAD	A	600	53/53	1.00	0.02	11,13,16,20	0
2	FAD	D	600	53/53	1.00	0.02	11,13,15,17	0
2	FAD	В	601	53/53	1.00	0.02	12,14,16,21	0
2	FAD	F	601	53/53	1.00	0.02	11,13,15,20	0
5	CL	В	603	1/1	1.00	0.04	25,25,25,25	0
5	CL	F	605	1/1	1.00	0.04	22,22,22,22	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around B3P A 602:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around B3P F 602:












Electron density around B3P A 601:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around B3P F 604:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around FAD E 600:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around FAD C 600:












Electron density around FAD G 600:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around FAD H 602:












Electron density around FAD A 600:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around FAD D 600:












Electron density around FAD B 601:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around FAD F 601:









6.5 Other polymers (i)

There are no such residues in this entry.

