



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 03:23 AM EDT

PDB ID : 3HJ4  
Title : Minor Editosome-Associated TUTase 1  
Authors : Stagno, J.; Luecke, H.  
Deposited on : 2009-05-20  
Resolution : 1.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	384	
1	B	384	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7116 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Minor Editosome-Associated TUTase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	381	3110	1964	552	576	6	12	0	6	0
1	B	383	3127	1976	553	580	6	12	0	7	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ASN	DELETION	UNP Q4GZ86
B	?	-	ASN	DELETION	UNP Q4GZ86


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	439	Total	O	0	0
			439	439		
2	B	440	Total	O	0	0
			440	440		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Minor Editosome-Associated TUTase



- Molecule 1: Minor Editosome-Associated TUTase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.21Å 102.03Å 66.07Å 90.00° 111.99° 90.00°	Depositor
Resolution (Å)	31.41 – 1.56 31.41 – 1.56	Depositor EDS
% Data completeness (in resolution range)	98.1 (31.41-1.56) 97.9 (31.41-1.56)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.43 (at 1.56Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.162 , 0.203 0.176 , 0.214	Depositor DCC
$R_{free}$ test set	5431 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.0	Xtrriage
Anisotropy	0.560	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 43.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.074 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7116	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.59	0/3182	0.70	2/4274 (0.0%)
1	B	0.61	0/3202	0.73	3/4301 (0.1%)
All	All	0.60	0/6384	0.72	5/8575 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	293	ARG	NE-CZ-NH1	6.92	123.76	120.30
1	A	93	MSE	CG-SE-CE	-6.36	84.90	98.90
1	B	93	MSE	CG-SE-CE	-5.84	86.05	98.90
1	A	92	ARG	NE-CZ-NH1	5.33	122.97	120.30
1	B	362	ARG	NE-CZ-NH2	-5.30	117.65	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3110	0	3053	12	0
1	B	3127	0	3072	16	0
2	A	439	0	0	1	0
2	B	440	0	0	5	0
All	All	7116	0	6125	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:LEU:HD21	1:A:298:LYS:HE2	1.54	0.88
1:A:78:TRP:HZ3	1:A:288:LEU:HD12	1.38	0.86
1:A:78:TRP:CZ3	1:A:288:LEU:HD12	2.21	0.74
1:B:111:ILE:HD12	1:B:117:VAL:CG2	2.22	0.70
1:B:294:LEU:HD21	1:B:298:LYS:HE3	1.76	0.68
1:B:362:ARG:NH2	1:B:367:ASP:OD1	2.29	0.66
1:A:383:THR:HG22	1:A:383:THR:O	1.99	0.63
1:A:230:TYR:HA	1:A:233:LEU:HD22	1.82	0.60
1:B:115:ILE:HG22	2:B:448:HOH:O	2.04	0.57
1:B:230:TYR:HA	1:B:233:LEU:HD22	1.88	0.55
1:B:111:ILE:HB	1:B:117:VAL:HG22	1.90	0.54
1:A:171:GLU:OE1	1:A:383:THR:HB	2.09	0.53
1:B:111:ILE:HD12	1:B:117:VAL:HG21	1.92	0.50
1:B:115:ILE:HG23	2:B:729:HOH:O	2.12	0.47
1:A:294:LEU:CD2	1:A:298:LYS:HE2	2.36	0.47
1:B:294:LEU:C	1:B:294:LEU:HD23	2.35	0.46
1:B:149:ARG:NH2	2:B:721:HOH:O	2.41	0.46
1:B:294:LEU:HD21	1:B:298:LYS:CE	2.45	0.45
1:A:38:ASP:HB3	1:A:103:MSE:HE1	2.00	0.44
1:A:78:TRP:CZ3	1:A:288:LEU:CD1	2.97	0.43
1:B:329:SER:HB3	2:B:794:HOH:O	2.18	0.43
1:B:56:VAL:HG11	2:B:670:HOH:O	2.18	0.42
1:A:362:ARG:NH2	1:A:367:ASP:OD1	2.51	0.42
1:B:362:ARG:HH22	1:B:367:ASP:CG	2.21	0.42
1:A:56:VAL:HG11	2:A:808:HOH:O	2.18	0.42
1:B:75:PHE:CZ	1:B:134:GLY:HA3	2.55	0.41
1:A:78:TRP:HZ3	1:A:288:LEU:CD1	2.20	0.41
1:B:111:ILE:HD12	1:B:117:VAL:HG22	1.99	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/384 (100%)	378 (98%)	7 (2%)	0	100	100
1	B	388/384 (101%)	382 (98%)	6 (2%)	0	100	100
All	All	773/768 (101%)	760 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/325 (105%)	332 (97%)	9 (3%)	46	17
1	B	344/325 (106%)	335 (97%)	9 (3%)	46	17
All	All	685/650 (105%)	667 (97%)	18 (3%)	47	17

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	VAL
1	A	103	MSE
1	A	115	ILE
1	A	130	ASP
1	A	152	PHE
1	A	219	MSE
1	A	233	LEU
1	A	350	ARG
1	A	362	ARG
1	B	112	ARG
1	B	130	ASP
1	B	152	PHE
1	B	177	ARG
1	B	233	LEU

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	B	350	ARG
1	B	362	ARG
1	B	370[A]	SER
1	B	370[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	369/384 (96%)	0.29	32 (8%) 10 11	8, 14, 26, 42	0
1	B	371/384 (96%)	0.23	22 (5%) 22 26	8, 15, 25, 41	0
All	All	740/768 (96%)	0.26	54 (7%) 15 17	8, 14, 26, 42	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	113	ALA	5.9
1	B	115	ILE	5.6
1	A	112	ARG	5.0
1	A	115	ILE	4.8
1	A	383	THR	4.6
1	B	113	ALA	4.6
1	B	3	VAL	4.1
1	B	111	ILE	4.0
1	B	384	VAL	3.9
1	A	3	VAL	3.8
1	A	176	GLU	3.8
1	A	235	ASP	3.8
1	B	112	ARG	3.6
1	B	114	ARG	3.6
1	B	192	VAL	3.1
1	A	111	ILE	3.0
1	A	222	GLN	3.0
1	A	223	GLU	3.0
1	B	23	PRO	3.0
1	B	265	VAL	2.9
1	A	265	VAL	2.9
1	A	110	TYR	2.9
1	A	267	LEU	2.8
1	A	23	PRO	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	228	PRO	2.8
1	B	235	ASP	2.8
1	B	110	TYR	2.8
1	A	114	ARG	2.7
1	A	382	ASP	2.7
1	B	24	GLU	2.6
1	B	109	ARG	2.6
1	A	192	VAL	2.6
1	A	229	ILE	2.6
1	B	370[A]	SER	2.6
1	A	375	ASP	2.6
1	A	123	GLY	2.5
1	B	254	TYR	2.4
1	A	24	GLU	2.4
1	B	373	LYS	2.3
1	A	109	ARG	2.3
1	A	231	ASP	2.3
1	A	254	TYR	2.2
1	A	145	LEU	2.2
1	B	190	LEU	2.2
1	A	255	TYR	2.2
1	B	333	VAL	2.2
1	B	163	VAL	2.1
1	A	374	PHE	2.1
1	A	186	THR	2.1
1	B	222	GLN	2.1
1	A	177	ARG	2.1
1	B	294	LEU	2.1
1	A	376	GLU	2.0
1	A	232	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.