



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 20, 2023 – 02:25 AM EDT

PDB ID : 5HK5  
Title : Structure of the Grem2-GDF5 Inhibitory Complex  
Authors : Nolan, K.; Thompson, T.B.; Read, R.J.  
Deposited on : 2016-01-13  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	147	
1	F	147	
1	G	147	
1	H	147	
2	A	120	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	B	120	 <p>2% 74% 12% 12%</p>
2	C	120	 <p>% 77% 11% 12%</p>
2	D	120	 <p>4% 79% 8% 12%</p>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7436 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gremlin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	E	128	1033	652	187	184	10	0	0	0
1	F	129	1042	658	189	185	10	0	0	0
1	G	129	1042	658	189	185	10	0	0	0
1	H	119	972	611	178	173	10	0	0	0


- Molecule 2 is a protein called Growth/differentiation factor 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	105	834	525	141	157	11	0	0	0
2	B	105	834	525	141	157	11	0	0	0
2	C	105	845	535	142	157	11	0	1	0
2	D	105	834	525	141	157	11	0	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gremlin-2




- Molecule 1: Gremlin-2




- Molecule 1: Gremlin-2




- Molecule 1: Gremlin-2








● Molecule 2: Growth/differentiation factor 5




● Molecule 2: Growth/differentiation factor 5



● Molecule 2: Growth/differentiation factor 5



● Molecule 2: Growth/differentiation factor 5



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	114.72Å 125.79Å 125.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.55 – 2.90 50.52 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.55-2.90) 99.2 (50.52-2.90)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.10 (at 2.91Å)	Xtrriage
Refinement program	REFMAC 5.8.0124	Depositor
R, $R_{free}$	0.261 , 0.282 0.262 , 0.282	Depositor DCC
$R_{free}$ test set	2113 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.9	Xtrriage
Anisotropy	0.173	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 1.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.467 for -h,l,k	Xtrriage
Reported twinning fraction	0.517 for H, K, L 0.483 for -H, L, K	Depositor
Outliers	6 of 40921 reflections (0.015%)	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	7436	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.46 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7298e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	E	0.68	0/1057	0.95	5/1424 (0.4%)
1	F	0.65	0/1066	0.88	1/1435 (0.1%)
1	G	0.61	0/1066	0.86	2/1435 (0.1%)
1	H	0.68	0/993	0.97	2/1336 (0.1%)
2	A	0.50	0/856	0.78	1/1163 (0.1%)
2	B	0.51	0/856	0.78	1/1163 (0.1%)
2	C	0.54	0/872	0.77	1/1186 (0.1%)
2	D	0.54	0/856	0.75	1/1163 (0.1%)
All	All	0.60	0/7622	0.85	14/10305 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	2

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	60	LEU	CA-CB-CG	7.06	131.54	115.30
1	E	68	LEU	CB-CG-CD2	6.88	122.69	111.00
1	G	68	LEU	CB-CG-CD2	6.85	122.64	111.00
1	E	140	LEU	CA-CB-CG	6.46	130.16	115.30
2	B	120	ARG	NE-CZ-NH2	-6.24	117.18	120.30
1	E	108	ARG	NE-CZ-NH1	6.22	123.41	120.30
1	H	51	GLU	N-CA-C	-6.06	94.64	111.00
2	C	120	ARG	NE-CZ-NH2	-5.88	117.36	120.30
1	E	95	ARG	NE-CZ-NH1	5.35	122.98	120.30
1	H	87	CYS	N-CA-C	5.34	125.43	111.00

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	95	ARG	NE-CZ-NH1	5.17	122.89	120.30
1	F	65	ARG	NE-CZ-NH1	5.10	122.85	120.30
2	D	99	ASP	CB-CG-OD1	5.04	122.83	118.30
1	E	46	HIS	N-CA-C	5.00	124.50	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	134	GLU	Peptide
1	H	135	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1033	0	1032	23	0
1	F	1042	0	1045	19	1
1	G	1042	0	1045	23	0
1	H	972	0	973	21	0
2	A	834	0	796	8	1
2	B	834	0	796	29	0
2	C	845	0	806	7	0
2	D	834	0	796	6	0
All	All	7436	0	7289	110	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:60:LEU:HD11	2:B:118:GLY:CA	1.31	1.61
2:B:60:LEU:CD1	2:B:118:GLY:CA	2.09	1.31
2:B:60:LEU:CD1	2:B:118:GLY:HA2	1.61	1.29

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:56:LEU:HD21	2:B:62:PRO:HG3	1.27	1.15
2:B:60:LEU:HD21	2:B:119:CYS:N	1.69	1.07
1:E:84:GLU:OE2	1:E:147:LYS:NZ	1.88	1.07
1:H:135:LEU:HG	1:H:136:GLU:HB3	1.38	1.05
2:B:60:LEU:CD1	2:B:118:GLY:HA3	1.80	1.02
2:B:60:LEU:HD21	2:B:119:CYS:H	0.85	1.02
2:B:60:LEU:HD13	2:B:118:GLY:HA2	1.42	0.98
2:B:60:LEU:HD11	2:B:118:GLY:HA2	1.23	0.96
2:B:60:LEU:CD2	2:B:119:CYS:H	1.79	0.94
2:B:56:LEU:CD2	2:B:62:PRO:HG3	1.99	0.92
2:B:58:SER:O	2:B:60:LEU:N	2.00	0.92
1:H:135:LEU:HG	1:H:136:GLU:CB	2.00	0.92
2:B:60:LEU:HD11	2:B:118:GLY:HA3	0.93	0.91
2:B:56:LEU:HD21	2:B:62:PRO:CG	2.05	0.87
1:E:95:ARG:NH2	2:B:30:ASP:O	2.08	0.86
1:H:83:SER:OG	1:H:89:SER:HB3	1.77	0.84
1:G:95:ARG:NH2	2:A:30:ASP:O	2.12	0.83
1:H:75:THR:OG1	1:H:95:ARG:HD3	1.81	0.80
1:G:33:PRO:HG3	2:D:67:VAL:HG13	1.68	0.75
1:F:83:SER:OG	1:F:89:SER:HB3	1.88	0.74
1:F:75:THR:OG1	1:F:95:ARG:HD3	1.89	0.73
1:E:75:THR:OG1	1:E:95:ARG:HD3	1.89	0.72
1:G:75:THR:OG1	1:G:95:ARG:HD3	1.89	0.72
1:H:140:LEU:HD21	1:H:145:ARG:HH21	1.53	0.71
1:G:127:ARG:NH1	1:G:129:THR:HG21	2.05	0.70
1:E:33:PRO:HG3	2:C:67:VAL:HG13	1.72	0.70
2:B:58:SER:OG	2:B:59:HIS:N	2.23	0.69
2:A:57:ARG:O	2:A:60:LEU:HG	1.93	0.68
1:E:127:ARG:HD2	1:G:127:ARG:HD2	1.78	0.65
2:C:89:ARG:HG2	2:C:116:SER:OG	2.00	0.61
1:H:135:LEU:CG	1:H:136:GLU:HB3	2.23	0.61
1:G:127:ARG:NH1	1:G:129:THR:CG2	2.64	0.59
1:H:60:LEU:HD22	1:H:110:VAL:HG21	1.84	0.59
1:H:135:LEU:HD23	1:H:136:GLU:CD	2.22	0.59
1:E:60:LEU:HD22	1:E:110:VAL:HG21	1.84	0.59
1:F:60:LEU:HD22	1:F:110:VAL:HG21	1.85	0.59
1:G:60:LEU:HD22	1:G:110:VAL:HG21	1.85	0.59
1:G:59:ALA:HB1	2:B:105:VAL:HG13	1.85	0.58
1:F:112:LYS:HG2	1:F:113:GLU:H	1.69	0.58
1:F:130:SER:OG	1:F:148:LYS:HD2	2.02	0.57
2:D:29:LYS:HE3	2:D:41:LEU:HD13	1.87	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:29:LYS:HE3	2:A:41:LEU:HD13	1.87	0.57
2:B:56:LEU:HD11	2:B:62:PRO:HG3	1.86	0.56
2:B:60:LEU:O	2:B:61:GLU:HB2	2.05	0.56
1:G:127:ARG:CZ	1:G:129:THR:CG2	2.84	0.56
2:B:29:LYS:HE3	2:B:41:LEU:HD13	1.86	0.56
1:F:132:ILE:HG23	1:F:146:ILE:HG23	1.88	0.56
2:C:29:LYS:HE3	2:C:41:LEU:HD13	1.88	0.55
1:G:127:ARG:CZ	1:G:129:THR:HG21	2.36	0.55
1:G:75:THR:OG1	1:G:95:ARG:CD	2.55	0.54
1:E:75:THR:OG1	1:E:95:ARG:CD	2.54	0.54
1:E:92:ILE:HG23	2:D:74:SER:O	2.09	0.52
2:B:57:ARG:CG	2:B:59:HIS:CD2	2.93	0.52
1:E:59:ALA:HB1	2:A:105:VAL:HG13	1.92	0.52
1:E:107:PRO:HG3	1:F:152:VAL:HG23	1.92	0.51
2:A:18:ARG:NH1	2:A:53:GLU:O	2.42	0.51
2:B:56:LEU:CD1	2:B:62:PRO:HG3	2.42	0.50
1:G:59:ALA:HB1	2:B:105:VAL:CG1	2.41	0.49
2:D:18:ARG:NH1	2:D:53:GLU:O	2.43	0.49
1:E:108:ARG:HE	1:F:150:GLN:CD	2.15	0.49
1:H:135:LEU:HG	1:H:136:GLU:N	2.26	0.49
1:H:80:GLN:CG	1:H:92:ILE:HD11	2.42	0.49
2:B:57:ARG:HG2	2:B:59:HIS:HD2	1.78	0.49
1:F:75:THR:OG1	1:F:95:ARG:CD	2.61	0.49
1:G:57:GLN:NE2	2:B:108:GLN:OE1	2.39	0.49
1:E:96:PHE:HB2	1:F:105:TYR:O	2.13	0.49
1:G:107:PRO:HG3	1:H:152:VAL:HG23	1.95	0.48
1:G:140:LEU:HD13	1:G:145:ARG:NH2	2.28	0.48
1:G:92:ILE:HG23	2:C:74:SER:O	2.14	0.48
1:H:135:LEU:C	1:H:136:GLU:HG2	2.34	0.48
1:G:74:LYS:HG3	1:G:76:GLN:HE21	1.78	0.47
1:H:87:CYS:O	1:H:136:GLU:HG3	2.13	0.47
2:C:18:ARG:NH1	2:C:53:GLU:O	2.43	0.47
1:F:83:SER:OG	1:F:89:SER:CB	2.62	0.47
1:F:112:LYS:HG2	1:F:113:GLU:N	2.29	0.47
2:B:57:ARG:HG3	2:B:59:HIS:CD2	2.51	0.46
1:E:26:PRO:HA	1:G:74:LYS:NZ	2.31	0.46
1:E:93:LEU:CD2	2:D:71:LEU:HD11	2.47	0.45
1:E:114:GLU:HG2	1:E:115:ASP:N	2.32	0.44
1:G:97:CYS:HB2	1:H:105:TYR:HB3	1.98	0.44
1:E:127:ARG:NH1	1:E:153:LYS:HG3	2.33	0.44
2:D:56:LEU:HG	2:D:70:THR:HG21	2.00	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:56:LEU:HG	2:A:70:THR:HG21	2.00	0.44
1:G:127:ARG:HH12	1:G:129:THR:HG21	1.81	0.43
1:E:105:TYR:CG	1:F:125:PRO:HD3	2.53	0.43
1:E:114:GLU:HG2	1:E:115:ASP:OD1	2.17	0.43
1:E:97:CYS:HB2	1:F:105:TYR:HB3	2.01	0.43
2:B:57:ARG:NH2	2:B:59:HIS:CE1	2.87	0.43
1:E:45:TRP:N	1:E:51:GLU:OE1	2.52	0.43
1:H:80:GLN:HG2	1:H:92:ILE:HD11	2.00	0.43
2:C:56:LEU:HG	2:C:70:THR:HG21	1.99	0.43
1:G:93:LEU:CD2	2:C:71:LEU:HD11	2.49	0.43
1:F:33:PRO:HD2	1:F:52:VAL:HG13	2.01	0.42
1:H:84:GLU:O	1:H:86:GLY:O	2.37	0.42
1:H:136:GLU:O	1:H:143:PRO:O	2.38	0.42
1:F:30:ILE:HD12	1:F:52:VAL:HG21	2.01	0.42
1:H:135:LEU:HG	1:H:136:GLU:CA	2.50	0.42
2:B:57:ARG:HG2	2:B:59:HIS:CD2	2.54	0.42
1:E:78:LEU:HD21	1:F:110:VAL:HG11	2.01	0.42
1:G:105:TYR:CG	1:H:125:PRO:HD3	2.55	0.42
1:E:59:ALA:HB1	2:A:105:VAL:CG1	2.49	0.42
1:F:80:GLN:CD	1:F:92:ILE:HD11	2.41	0.41
2:A:60:LEU:HD12	2:A:62:PRO:HD3	2.01	0.41
1:G:78:LEU:HD21	1:H:110:VAL:HG11	2.03	0.41
2:B:56:LEU:HD21	2:B:62:PRO:CD	2.49	0.41
1:E:105:TYR:CE2	1:F:125:PRO:HG3	2.56	0.41
1:H:135:LEU:HG	1:H:136:GLU:CG	2.50	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:90:ARG:NH1	2:A:53:GLU:OE1[4_785]	1.96	0.24

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	124/147 (84%)	111 (90%)	9 (7%)	4 (3%)	4	16
1	F	125/147 (85%)	115 (92%)	10 (8%)	0	100	100
1	G	125/147 (85%)	115 (92%)	9 (7%)	1 (1%)	19	51
1	H	117/147 (80%)	102 (87%)	12 (10%)	3 (3%)	5	20
2	A	103/120 (86%)	102 (99%)	1 (1%)	0	100	100
2	B	103/120 (86%)	99 (96%)	2 (2%)	2 (2%)	8	28
2	C	104/120 (87%)	103 (99%)	1 (1%)	0	100	100
2	D	103/120 (86%)	102 (99%)	1 (1%)	0	100	100
All	All	904/1068 (85%)	849 (94%)	45 (5%)	10 (1%)	14	42

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	113	GLU
2	B	58	SER
1	H	88	ARG
2	B	59	HIS
1	G	85	GLU
1	H	135	LEU
1	E	141	ASP
1	E	115	ASP
1	H	46	HIS
1	E	140	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	119/137 (87%)	117 (98%)	2 (2%)	60	86
1	F	120/137 (88%)	117 (98%)	3 (2%)	47	78

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	120/137 (88%)	118 (98%)	2 (2%)	60	86
1	H	113/137 (82%)	108 (96%)	5 (4%)	28	61
2	A	96/108 (89%)	96 (100%)	0	100	100
2	B	96/108 (89%)	95 (99%)	1 (1%)	76	92
2	C	97/108 (90%)	96 (99%)	1 (1%)	76	92
2	D	96/108 (89%)	96 (100%)	0	100	100
All	All	857/980 (87%)	843 (98%)	14 (2%)	62	86

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	45	TRP
1	E	137	CYS
1	F	30	ILE
1	F	137	CYS
1	F	146	ILE
1	G	137	CYS
1	G	140	LEU
1	H	47	HIS
1	H	106	ILE
1	H	113	GLU
1	H	137	CYS
1	H	148	LYS
2	B	68	ILE
2	C	68	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	126	GLN
1	E	154	HIS
1	G	76	GLN
2	A	108	GLN
2	B	59	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	E	128/147 (87%)	0.75	9 (7%) 16 12	12, 34, 63, 103	0
1	F	129/147 (87%)	1.21	13 (10%) 7 5	13, 38, 80, 99	0
1	G	129/147 (87%)	0.76	8 (6%) 20 16	16, 32, 71, 97	0
1	H	119/147 (80%)	1.02	17 (14%) 2 2	17, 39, 71, 84	0
2	A	105/120 (87%)	0.85	1 (0%) 82 82	12, 26, 54, 75	0
2	B	105/120 (87%)	0.67	3 (2%) 51 47	15, 32, 60, 90	0
2	C	105/120 (87%)	0.77	1 (0%) 82 82	13, 28, 48, 84	0
2	D	105/120 (87%)	0.70	5 (4%) 30 27	19, 30, 52, 77	0
All	All	925/1068 (86%)	0.85	57 (6%) 20 16	12, 32, 66, 103	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	33	PRO	8.3
1	F	32	SER	7.4
1	F	28	GLY	7.2
1	F	26	PRO	7.2
1	G	110	VAL	6.8
1	F	34	TYR	5.7
1	F	146	ILE	5.2
1	F	31	PRO	4.8
1	F	27	ALA	4.7
1	F	135	LEU	4.6
2	D	16	LYS	4.5
1	F	30	ILE	4.3
1	E	110	VAL	4.2
1	G	111	LYS	4.0
1	H	146	ILE	3.9
1	H	139	GLY	3.8

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	H	163	SER	3.8
1	E	112	LYS	3.7
1	H	54	ALA	3.6
2	B	60	LEU	3.4
1	H	52	VAL	3.3
1	H	135	LEU	3.2
1	H	143	PRO	3.0
1	G	112	LYS	3.0
1	E	140	LEU	2.9
1	H	136	GLU	2.9
2	A	36	TRP	2.8
1	H	72	TRP	2.7
1	H	128	VAL	2.7
2	B	36	TRP	2.6
1	G	26	PRO	2.6
1	H	55	SER	2.6
1	H	138	PRO	2.6
1	H	148	LYS	2.5
1	E	126	GLN	2.5
1	E	111	LYS	2.5
2	D	120	ARG	2.5
1	F	72	TRP	2.5
1	G	126	GLN	2.4
2	C	16	LYS	2.3
1	F	47	HIS	2.3
1	E	120	CYS	2.2
2	D	109	TYR	2.2
1	H	88	ARG	2.2
2	D	35	ASP	2.2
1	E	68	LEU	2.1
2	D	95	ILE	2.1
2	B	95	ILE	2.1
1	G	127	ARG	2.1
1	E	85	GLU	2.1
1	E	93	LEU	2.1
1	G	138	PRO	2.1
1	F	46	HIS	2.1
1	H	132	ILE	2.0
1	G	135	LEU	2.0
1	H	133	VAL	2.0
1	H	140	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.