



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 20, 2023 – 03:10 AM EDT

PDB ID : 2HOI  
Title : Crystal structure of the tetrameric pre-cleavage synaptic complex in the cre-loxp site-specific recombination  
Authors : Ghosh, K.; Van Duyne, G.D.  
Deposited on : 2006-07-14  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	35	
1	E	35	
2	D	35	
2	F	35	
3	A	343	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	B	343	 83% 11% 6%
3	G	343	%  80% 12% • 6%
3	H	343	%  82% 10% • 6%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13811 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called LoxP DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	C	35	Total	C	N	O	P	0	0	0
			710	344	124	208	34			
1	E	34	Total	C	N	O	P	0	0	0
			693	334	122	203	34			

- Molecule 2 is a DNA chain called LoxP DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	D	34	Total	C	N	O	P	0	0	0
			701	337	125	205	34			
2	F	34	Total	C	N	O	P	0	0	0
			701	337	125	205	34			

- Molecule 3 is a protein called Recombinase Cre.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	322	Total	C	N	O	S	0	0	0
			2546	1581	485	465	15			
3	B	322	Total	C	N	O	S	0	0	0
			2546	1581	485	465	15			
3	G	322	Total	C	N	O	S	0	0	0
			2546	1581	485	465	15			
3	H	322	Total	C	N	O	S	0	0	0
			2546	1581	485	465	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	201	ALA	LYS	engineered mutation	UNP P06956
B	201	ALA	LYS	engineered mutation	UNP P06956
G	201	ALA	LYS	engineered mutation	UNP P06956
H	201	ALA	LYS	engineered mutation	UNP P06956


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	50	Total O 50 50	0	0
4	D	42	Total O 42 42	0	0
4	E	36	Total O 36 36	0	0
4	F	50	Total O 50 50	0	0
4	A	168	Total O 168 168	0	0
4	B	177	Total O 177 177	0	0
4	G	144	Total O 144 144	0	0
4	H	155	Total O 155 155	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: LoxP DNA




- Molecule 1: LoxP DNA




- Molecule 2: LoxP DNA



- Molecule 2: LoxP DNA





- Molecule 3: Recombinase Cre





- Molecule 3: Recombinase Cre







• Molecule 3: Recombinase Cre



• Molecule 3: Recombinase Cre



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	136.78Å 136.78Å 218.28Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.92 – 2.60 19.92 – 2.60	Depositor EDS
% Data completeness (in resolution range)	81.5 (19.92-2.60) 81.5 (19.92-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	9.02 (at 2.59Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.184 , 0.246 0.187 , 0.247	Depositor DCC
$R_{free}$ test set	3007 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.1	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.018 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	13811	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	C	1.07	3/795 (0.4%)	1.93	60/1224 (4.9%)
1	E	1.09	4/776 (0.5%)	1.83	55/1194 (4.6%)
2	D	1.04	2/786 (0.3%)	1.90	51/1212 (4.2%)
2	F	1.01	1/786 (0.1%)	1.91	57/1212 (4.7%)
3	A	0.38	0/2587	0.58	1/3489 (0.0%)
3	B	0.36	0/2587	0.58	0/3489
3	G	0.35	0/2587	0.57	0/3489
3	H	0.37	0/2587	0.61	0/3489
All	All	0.60	10/13491 (0.1%)	1.09	224/18798 (1.2%)

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	1	DT	C5-C7	8.11	1.54	1.50
2	D	22	DC	N1-C2	-7.00	1.33	1.40
1	E	27	DC	N1-C2	-6.54	1.33	1.40
2	D	27	DC	N1-C2	-5.88	1.34	1.40
2	F	22	DC	N1-C2	-5.78	1.34	1.40

The worst 5 of 224 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	20	DC	O4'-C1'-N1	10.33	115.23	108.00
2	F	17	DG	O4'-C4'-C3'	-9.91	100.05	106.00
2	D	17	DG	O4'-C4'-C3'	-8.54	100.88	106.00
2	D	26	DA	O4'-C4'-C3'	-8.44	100.94	106.00
2	F	6	DG	O4'-C4'-C3'	-7.72	101.37	106.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	710	0	400	11	0
1	E	693	0	387	11	0
2	D	701	0	388	6	0
2	F	701	0	388	8	0
3	A	2546	0	2563	37	0
3	B	2546	0	2563	22	0
3	G	2546	0	2563	29	0
3	H	2546	0	2563	25	0
4	A	168	0	0	4	0
4	B	177	0	0	4	0
4	C	50	0	0	3	0
4	D	42	0	0	0	0
4	E	36	0	0	2	0
4	F	50	0	0	4	0
4	G	144	0	0	2	0
4	H	155	0	0	3	0
All	All	13811	0	11815	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 124 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:F:428:HOH:O	3:H:121:ARG:HD2	1.69	0.92
1:C:22:DT:H72	4:A:430:HOH:O	1.70	0.91
3:A:72:ARG:HG3	3:A:116:VAL:HG11	1.56	0.86
3:A:72:ARG:HG3	3:A:116:VAL:CG1	2.05	0.85
4:C:60:HOH:O	3:A:97:MET:SD	2.36	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	320/343 (93%)	318 (99%)	2 (1%)	0	100	100
3	B	320/343 (93%)	319 (100%)	1 (0%)	0	100	100
3	G	320/343 (93%)	314 (98%)	6 (2%)	0	100	100
3	H	320/343 (93%)	317 (99%)	3 (1%)	0	100	100
All	All	1280/1372 (93%)	1268 (99%)	12 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	268/286 (94%)	253 (94%)	15 (6%)	21	42
3	B	268/286 (94%)	258 (96%)	10 (4%)	34	60
3	G	268/286 (94%)	253 (94%)	15 (6%)	21	42
3	H	268/286 (94%)	252 (94%)	16 (6%)	19	39
All	All	1072/1144 (94%)	1016 (95%)	56 (5%)	23	46

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	116	VAL
3	H	323	ASN
3	G	227	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	H	293	VAL
3	H	149	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	281	GLN
3	H	311	GLN
3	H	323	ASN
3	H	255	GLN
3	A	323	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	C	35/35 (100%)	-0.23	1 (2%) 51 45	16, 30, 60, 77	0
1	E	34/35 (97%)	-0.44	0 100 100	15, 31, 57, 62	0
2	D	34/35 (97%)	-0.55	1 (2%) 51 45	17, 31, 48, 74	0
2	F	34/35 (97%)	-0.56	1 (2%) 51 45	16, 33, 55, 79	0
3	A	322/343 (93%)	-0.49	4 (1%) 79 76	18, 28, 39, 50	0
3	B	322/343 (93%)	-0.54	1 (0%) 94 93	19, 28, 38, 42	0
3	G	322/343 (93%)	-0.42	4 (1%) 79 76	17, 32, 49, 54	0
3	H	322/343 (93%)	-0.55	3 (0%) 84 82	16, 26, 38, 46	0
All	All	1425/1512 (94%)	-0.49	15 (1%) 80 78	15, 29, 43, 79	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	DT	5.5
3	H	341	ASP	5.0
3	B	279	SER	3.0
3	A	21	ASP	2.7
2	D	2	DA	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.