



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 12:21 pm GMT

PDB ID : 2HOR
Title : Crystal structure of alliinase from garlic- apo form
Authors : Shimon, L.J.W.; Rabinkov, A.; Wilcheck, M.; Mirelman, D.; Frolow, F.
Deposited on : 2006-07-16
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtrriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2957 (1.60-1.60)
Clashscore	122126	3202 (1.60-1.60)
Ramachandran outliers	120053	3117 (1.60-1.60)
Sidechain outliers	120020	3116 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	427	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BMA	A	504	X	-	-	-

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 4122 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alliin lyase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	425	3506	2242	581	657	26	6	15	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	176	ASP	ASN	SEE REMARK 999	UNP Q01594

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0

- Molecule 3 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula: C₆H₁₂O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	10	6	4	0	0

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	10	6	4	0	0

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	2	Total	Cl	0	0
			2	2		

- Molecule 9 is NITRATE ION (three-letter code: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	N	O	0	0
			4	1	3		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	496	Total	O	0	0
			496	496		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alliin lyase 1



4 Data and refinement statistics i

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	81.12Å 81.12Å 164.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.60 – 1.60 39.57 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.60-1.60) 99.9 (39.57-1.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.47 (at 1.60Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.163 , 0.179 0.234 , 0.243	Depositor DCC
R_{free} test set	3663 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtrriage
Anisotropy	0.361	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 55.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4122	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CL, NO3, FUC, ACT, MAN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.60	2/3640 (0.1%)	0.69	5/4922 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	55[A]	GLN	N-CA	-13.55	1.19	1.46
1	A	55[B]	GLN	N-CA	-13.55	1.19	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	55[A]	GLN	CA-C-N	7.45	131.10	116.20
1	A	55[B]	GLN	CA-C-N	7.45	131.10	116.20
1	A	55[A]	GLN	CA-C-O	-7.26	104.85	120.10
1	A	55[B]	GLN	CA-C-O	-7.26	104.85	120.10
1	A	54	ILE	C-N-CA	5.53	135.52	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3506	0	3438	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	42	0	36	0	0
3	A	10	0	10	0	0
4	A	10	0	7	0	0
5	A	23	0	21	0	0
6	A	25	0	0	0	0
7	A	4	0	3	0	0
8	A	2	0	0	0	0
9	A	4	0	0	0	0
10	A	496	0	0	5	0
All	All	4122	0	3515	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:354[A]:ARG:NH2	10:A:1172:HOH:O	2.26	0.66
1:A:2[B]:MET:HA	1:A:2[B]:MET:HE3	1.78	0.63
1:A:294:LEU:HG	1:A:298:LYS:HE2	1.82	0.61
1:A:232:HIS:ND1	1:A:318:GLY:HA3	2.18	0.59
1:A:19:ASN:HD22	1:A:21[B]:SER:H	1.51	0.57
1:A:298:LYS:HE3	10:A:899:HOH:O	2.05	0.55
1:A:19:ASN:HD22	1:A:21[A]:SER:H	1.51	0.55
1:A:19:ASN:ND2	1:A:21[B]:SER:H	2.06	0.54
1:A:19:ASN:ND2	1:A:21[A]:SER:H	2.06	0.53
1:A:2[B]:MET:HA	1:A:2[B]:MET:CE	2.39	0.49
1:A:143:LEU:HD22	1:A:221:LYS:HD3	1.97	0.47
1:A:19:ASN:C	1:A:19:ASN:HD22	2.20	0.45
1:A:68:PHE:HB2	1:A:321[A]:LYS:HE2	2.00	0.43
1:A:74:LYS:NZ	10:A:1138:HOH:O	2.33	0.43
1:A:83:LEU:HD13	1:A:83:LEU:C	2.38	0.43
1:A:124:ARG:NH2	1:A:266:LYS:HE2	2.34	0.43
1:A:333:LEU:HD22	1:A:339:PHE:HB2	1.99	0.43
1:A:226[B]:MET:HE1	1:A:240:ALA:HB2	2.02	0.41
1:A:354[B]:ARG:NH1	10:A:946:HOH:O	2.34	0.41
1:A:11[B]:GLU:HG3	10:A:970:HOH:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/427 (103%)	426 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/375 (104%)	383 (99%)	5 (1%)	71	53

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	LYS
1	A	19	ASN
1	A	25	ARG
1	A	374	LYS
1	A	423	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	500	1,3,2	14,14,15	0.90	1 (7%)	17,19,21	0.79	0
3	FUC	A	501	2	9,10,11	0.37	0	13,14,16	1.04	0
2	NAG	A	503	2,4	14,14,15	0.59	0	17,19,21	0.96	1 (5%)
4	BMA	A	504	2,5	10,10,12	0.85	0	11,13,17	0.86	0
5	MAN	A	505	4	11,11,12	0.42	0	15,15,17	0.92	1 (6%)
5	MAN	A	506	4	12,12,12	0.53	0	17,17,17	1.41	4 (23%)
2	NAG	A	555	1	14,14,15	0.52	0	17,19,21	1.27	1 (5%)
6	SO4	A	600	-	4,4,4	0.39	0	6,6,6	0.53	0
6	SO4	A	601	-	4,4,4	0.44	0	6,6,6	0.45	0
7	ACT	A	602	-	1,3,3	2.08	1 (100%)	0,3,3	0.00	-
6	SO4	A	701	-	4,4,4	0.17	0	6,6,6	0.15	0
6	SO4	A	702	-	4,4,4	0.34	0	6,6,6	0.20	0
6	SO4	A	703	-	4,4,4	0.23	0	6,6,6	0.48	0
9	NO3	A	707	-	1,3,3	3.57	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	500	1,3,2	-	0/6/23/26	0/1/1/1
3	FUC	A	501	2	-	0/0/17/20	0/1/1/1
2	NAG	A	503	2,4	-	0/6/23/26	0/1/1/1
4	BMA	A	504	2,5	1/1/3/5	0/2/15/22	0/1/1/1
5	MAN	A	505	4	-	0/2/19/22	0/1/1/1
5	MAN	A	506	4	-	0/2/22/22	0/1/1/1
2	NAG	A	555	1	-	0/6/23/26	0/1/1/1
6	SO4	A	600	-	-	0/0/0/0	0/0/0/0
6	SO4	A	601	-	-	0/0/0/0	0/0/0/0
7	ACT	A	602	-	-	0/0/0/0	0/0/0/0
6	SO4	A	701	-	-	0/0/0/0	0/0/0/0
6	SO4	A	702	-	-	0/0/0/0	0/0/0/0
6	SO4	A	703	-	-	0/0/0/0	0/0/0/0
9	NO3	A	707	-	-	0/0/0/0	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	NAG	O5-C1	-2.31	1.40	1.43
7	A	602	ACT	CH3-C	2.08	1.51	1.48
9	A	707	NO3	O1-N	3.57	1.41	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	503	NAG	O4-C4-C3	-2.73	103.98	110.34
5	A	506	MAN	C3-C4-C5	2.36	114.47	110.24
5	A	506	MAN	O5-C5-C6	2.37	112.39	106.43
5	A	505	MAN	C1-O5-C5	2.69	115.89	112.19
5	A	506	MAN	C4-C3-C2	2.75	115.64	110.83
5	A	506	MAN	C1-C2-C3	2.83	116.02	110.36
2	A	555	NAG	C1-O5-C5	3.54	117.05	112.19

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	504	BMA	C3

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.