



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 02:17 AM EDT

PDB ID : 5HOG
Title : Crystal structure of the carboxy-terminal domain of yeast Ctf4 bound to Dna2.
Authors : Simon, A.C.; Pellegrini, L.
Deposited on : 2016-01-19
Resolution : 3.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	
1	B	478	
1	C	478	
2	D	17	
2	E	17	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9598 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase alpha-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	425	Total 3425	C 2199	N 568	O 643	S 15	0	1	0
1	B	432	Total 3481	C 2233	N 578	O 654	S 16	0	1	0
1	C	296	Total 2405	C 1562	N 392	O 440	S 11	0	1	0

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	450	MET	HIS	conflict	UNP Q01454
A	451	GLY	ASN	conflict	UNP Q01454
A	452	SER	GLU	conflict	UNP Q01454
A	453	SER	HIS	conflict	UNP Q01454
A	454	HIS	SER	conflict	UNP Q01454
A	455	HIS	TYR	conflict	UNP Q01454
A	456	HIS	SER	conflict	UNP Q01454
A	457	HIS	ARG	conflict	UNP Q01454
A	458	HIS	VAL	conflict	UNP Q01454
A	460	SER	LYS	conflict	UNP Q01454
A	461	GLN	THR	conflict	UNP Q01454
A	462	ASP	HIS	conflict	UNP Q01454
A	463	PRO	SER	conflict	UNP Q01454
A	464	GLU	PHE	conflict	UNP Q01454
A	465	ASN	PRO	conflict	UNP Q01454
A	466	LEU	ILE	conflict	UNP Q01454
A	467	TYR	SER	conflict	UNP Q01454
A	468	PHE	LEU	conflict	UNP Q01454
A	469	GLN	ALA	conflict	UNP Q01454
A	470	GLY	ASN	conflict	UNP Q01454
B	450	MET	HIS	conflict	UNP Q01454
B	451	GLY	ASN	conflict	UNP Q01454
B	452	SER	GLU	conflict	UNP Q01454

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	453	SER	HIS	conflict	UNP Q01454
B	454	HIS	SER	conflict	UNP Q01454
B	455	HIS	TYR	conflict	UNP Q01454
B	456	HIS	SER	conflict	UNP Q01454
B	457	HIS	ARG	conflict	UNP Q01454
B	458	HIS	VAL	conflict	UNP Q01454
B	460	SER	LYS	conflict	UNP Q01454
B	461	GLN	THR	conflict	UNP Q01454
B	462	ASP	HIS	conflict	UNP Q01454
B	463	PRO	SER	conflict	UNP Q01454
B	464	GLU	PHE	conflict	UNP Q01454
B	465	ASN	PRO	conflict	UNP Q01454
B	466	LEU	ILE	conflict	UNP Q01454
B	467	TYR	SER	conflict	UNP Q01454
B	468	PHE	LEU	conflict	UNP Q01454
B	469	GLN	ALA	conflict	UNP Q01454
B	470	GLY	ASN	conflict	UNP Q01454
C	450	MET	HIS	conflict	UNP Q01454
C	451	GLY	ASN	conflict	UNP Q01454
C	452	SER	GLU	conflict	UNP Q01454
C	453	SER	HIS	conflict	UNP Q01454
C	454	HIS	SER	conflict	UNP Q01454
C	455	HIS	TYR	conflict	UNP Q01454
C	456	HIS	SER	conflict	UNP Q01454
C	457	HIS	ARG	conflict	UNP Q01454
C	458	HIS	VAL	conflict	UNP Q01454
C	460	SER	LYS	conflict	UNP Q01454
C	461	GLN	THR	conflict	UNP Q01454
C	462	ASP	HIS	conflict	UNP Q01454
C	463	PRO	SER	conflict	UNP Q01454
C	464	GLU	PHE	conflict	UNP Q01454
C	465	ASN	PRO	conflict	UNP Q01454
C	466	LEU	ILE	conflict	UNP Q01454
C	467	TYR	SER	conflict	UNP Q01454
C	468	PHE	LEU	conflict	UNP Q01454
C	469	GLN	ALA	conflict	UNP Q01454
C	470	GLY	ASN	conflict	UNP Q01454

- Molecule 2 is a protein called Dna2p.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
			Total	C	N				O
2	D	13	106	64	17	25	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	14	Total	C	N	O	0	0	0
			110	66	18	26			


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	29	Total	O	0	0
			29	29		
3	B	29	Total	O	0	0
			29	29		
3	C	13	Total	O	0	0
			13	13		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase alpha-binding protein

Chain A: 




- Molecule 1: DNA polymerase alpha-binding protein

Chain B: 



- Molecule 1: DNA polymerase alpha-binding protein

Chain C: 



ILE
LEU
ASN
LYS
LYS
ASN
GLU
ILE
GLY
ALA
ASP
THR
GLU
ALA
LEU
GLU
GLY
PHE
GLU
ASP
LYS
GLU
ILE
ILE
GLN
PRO
VAL
VAL
SER
MET
LYS
ALA
ALA
GLU
GLU
TYR
LEU
LEU
ARG
SER
LYS
GLN
VAL
VAL
SER
ALA
ALA
GLU
HIS
GLU
LEU
LEU
SER
GLN
ASP
VAL
SER
SER
GLU
LEU
LEU
THR
ALA
VAL
VAL
LYS
LYS
ILE
SER
GLU
ASN
ASP
GLY
ALA
MET
TYR
LEU
GLY
ASN



GLU
VAL
LEU
ALA
ALA
LEU
ASN
GLY
ALA
TYR
ASP
LYS
ALA
LEU
LEU
ARG
LEU
PHE
ALA
SER
ALA
CYS
SER
SER
ASP
GLN
ASN
ASN
VAL
VAL
GLU
SER
LYS
ALA
ALA
SER
LEU
LEU
GLU
HIS
GLU
LEU
LEU
LYS
GLN
ASP
VAL
SER
SER
GLU
LEU
LEU
THR
ALA
VAL
VAL
LYS
LYS
ILE
SER
GLU
ASN
ASP
GLY
ALA
MET
TYR
LEU
GLY
ASN

LYS
ILE
ASN
ASN
ILE
ARG
GLU
ALA
ARG
TYR
ASP
GLU
GLN
GLN
LEU
LYS

● Molecule 2: Dna2p



● Molecule 2: Dna2p



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	88.68Å 99.55Å 218.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.98 – 3.09 48.98 – 3.09	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.98-3.09) 99.7 (48.98-3.09)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.14 (at 3.07Å)	Xtrriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.180 , 0.226 0.176 , 0.223	Depositor DCC
R_{free} test set	1818 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	70.7	Xtrriage
Anisotropy	0.292	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9598	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/3510	0.42	0/4752
1	B	0.25	0/3567	0.41	0/4828
1	C	0.27	0/2481	0.44	0/3370
2	D	0.20	0/105	0.35	0/141
2	E	0.20	0/109	0.33	0/146
All	All	0.25	0/9772	0.42	0/13237

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3425	0	3369	27	0
1	B	3481	0	3431	22	0
1	C	2405	0	2326	17	0
2	D	106	0	100	1	0
2	E	110	0	103	1	0
3	A	29	0	0	0	0
3	B	29	0	0	0	0
3	C	13	0	0	0	0
All	All	9598	0	9329	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:789:GLU:HG3	1:B:818:PRO:HG3	1.72	0.72
1:A:789:GLU:HG3	1:A:818:PRO:HG3	1.73	0.71
1:B:899:VAL:HG13	1:B:911:VAL:HG13	1.77	0.65
1:A:790:ASN:HD21	1:A:815:ILE:HG23	1.67	0.60
1:B:511:TYR:HB2	1:B:530:ASP:HB3	1.85	0.58
1:B:818:PRO:HG2	1:B:821:MET:HB3	1.85	0.57
1:A:899:VAL:HG13	1:A:911:VAL:HG13	1.88	0.55
1:A:511:TYR:HB2	1:A:530:ASP:HB3	1.90	0.54
1:C:617:ALA:HB3	1:C:628:VAL:HB	1.90	0.54
1:C:510:GLN:HG2	1:C:531:LEU:HD23	1.90	0.53
1:A:724:SER:HB2	1:A:742:VAL:HG21	1.91	0.52
1:A:846:GLU:OE2	1:A:852:ASN:ND2	2.43	0.51
1:C:722:LEU:HD21	1:C:774:MET:HE2	1.93	0.51
1:A:634:HIS:CE1	1:B:634:HIS:HA	2.46	0.51
1:B:668:MET:HA	1:B:671:ASP:HB2	1.93	0.50
1:A:739:ASP:OD1	1:A:739:ASP:N	2.45	0.50
1:C:745:LEU:HG	1:C:754:CYS:HA	1.95	0.48
1:A:824:GLU:HA	1:A:869:LEU:HD11	1.95	0.48
1:A:490:ASP:HB3	1:A:506:LYS:HB2	1.94	0.48
1:A:569:ILE:HB	1:B:782:VAL:HG21	1.96	0.48
1:C:597:PHE:HB3	1:C:610:GLU:HB3	1.95	0.48
1:A:685:LYS:NZ	1:A:702:ASN:OD1	2.35	0.47
1:C:558:PRO:HG2	1:C:563:HIS:HB2	1.96	0.47
1:B:790:ASN:HD21	1:B:815:ILE:HG23	1.78	0.47
1:B:739:ASP:OD1	1:B:739:ASP:N	2.47	0.47
1:C:724:SER:HB2	1:C:742:VAL:HG21	1.96	0.47
1:A:508:SER:OG	1:A:509:GLU:N	2.46	0.46
1:C:755:ILE:HG23	1:C:769:PRO:HG2	1.98	0.46
1:C:728:ILE:HD11	1:C:742:VAL:HG23	1.97	0.46
1:A:601:ASN:HB3	1:A:607:PHE:HE2	1.81	0.46
1:B:833:LEU:HB3	1:B:858:LEU:HD21	1.98	0.45
1:A:833:LEU:HB3	1:A:858:LEU:HD21	1.98	0.45
1:C:511:TYR:HB2	1:C:530:ASP:HB3	1.99	0.45
1:B:864:LYS:HE3	2:E:215:LEU:HB3	1.98	0.44
1:A:563:HIS:ND1	1:B:880:GLU:OE1	2.34	0.44
1:A:818:PRO:HG2	1:A:821:MET:HB3	2.00	0.44
1:C:496:MET:HG2	1:C:745:LEU:HD22	1.98	0.44
1:A:552:GLY:O	1:A:570:ILE:N	2.36	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:880:GLU:HB3	1:C:603:PHE:CE1	2.53	0.44
1:A:714:GLU:OE1	1:C:649:ARG:NE	2.45	0.44
1:A:791:LYS:HE3	1:A:791:LYS:HB2	1.81	0.44
1:B:648:LYS:HE2	1:B:650:TYR:CZ	2.53	0.44
1:B:496:MET:HG3	1:B:501:TYR:HB3	2.01	0.43
1:A:866:LEU:HD21	1:A:888:GLU:HB2	2.01	0.43
1:A:698:PHE:CG	1:A:744:PRO:HG3	2.54	0.42
1:A:484:THR:O	1:A:492:ARG:HD2	2.19	0.42
1:B:663:ASN:OD1	1:B:663:ASN:N	2.52	0.42
1:C:698:PHE:CG	1:C:744:PRO:HG3	2.54	0.42
1:B:724:SER:HB2	1:B:742:VAL:HG21	2.02	0.42
1:B:866:LEU:HD21	1:B:888:GLU:HB2	2.02	0.42
1:C:751:THR:HB	1:C:773:GLU:HB3	2.01	0.41
1:B:548:GLN:HG3	1:B:551:THR:H	1.84	0.41
1:A:648:LYS:HB3	1:B:717:LYS:HD3	2.02	0.41
1:C:739:ASP:O	1:C:758:LYS:HA	2.20	0.41
1:C:750:ASP:O	1:C:776:ILE:HG12	2.21	0.41
1:B:866:LEU:HD23	1:B:889:LEU:HD23	2.03	0.41
1:B:702:ASN:HB3	1:B:724:SER:HG	1.86	0.41
1:B:704:LEU:HB3	1:B:722:LEU:HB3	2.02	0.41
1:A:864:LYS:HE3	2:D:215:LEU:HB3	2.03	0.40
1:A:652:LYS:NZ	1:A:655:CYS:SG	2.74	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	420/478 (88%)	401 (96%)	18 (4%)	1 (0%)	47 79
1	B	429/478 (90%)	413 (96%)	15 (4%)	1 (0%)	47 79
1	C	293/478 (61%)	275 (94%)	17 (6%)	1 (0%)	41 73

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	11/17 (65%)	11 (100%)	0	0	100	100
2	E	12/17 (71%)	11 (92%)	1 (8%)	0	100	100
All	All	1165/1468 (79%)	1111 (95%)	51 (4%)	3 (0%)	41	73

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	749	TYR
1	C	749	TYR
1	A	749	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	378/422 (90%)	378 (100%)	0	100	100
1	B	385/422 (91%)	384 (100%)	1 (0%)	92	96
1	C	267/422 (63%)	267 (100%)	0	100	100
2	D	13/16 (81%)	13 (100%)	0	100	100
2	E	13/16 (81%)	13 (100%)	0	100	100
All	All	1056/1298 (81%)	1055 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	852	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	634	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	425/478 (88%)	-0.16	1 (0%) 95 90	38, 60, 103, 128	0
1	B	432/478 (90%)	-0.13	4 (0%) 84 69	40, 61, 108, 140	0
1	C	296/478 (61%)	-0.00	17 (5%) 23 11	46, 69, 126, 162	0
2	D	13/17 (76%)	2.22	6 (46%) 0 0	120, 130, 144, 145	0
2	E	14/17 (82%)	2.06	7 (50%) 0 0	121, 135, 145, 148	0
All	All	1180/1468 (80%)	-0.05	35 (2%) 50 27	38, 63, 120, 162	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	207	SER	4.8
1	C	733	GLY	4.4
2	D	208	LEU	4.1
2	E	207	SER	3.8
1	B	666	SER	3.6
2	D	210	ASN	3.5
2	D	209	ARG	3.4
1	C	732	SER	3.3
1	B	927	LYS	3.2
1	C	673	ASN	3.1
1	C	736	GLU	2.9
1	B	665	ASN	2.8
2	E	220	GLY	2.8
2	D	215	LEU	2.7
1	C	729	TRP	2.7
1	C	731	MET	2.6
1	A	510	GLN	2.6
1	B	926	LEU	2.6
1	C	735	LYS	2.5
1	C	724	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	217	ASP	2.4
2	E	219	GLU	2.3
1	C	726	MET	2.3
1	C	737	THR	2.3
1	C	672	ALA	2.3
1	C	740	ILE	2.3
2	D	219	GLU	2.3
1	C	734	GLY	2.2
2	E	213	ASP	2.2
1	C	739	ASP	2.1
2	E	216	ASP	2.1
1	C	730	LYS	2.1
2	E	214	ILE	2.1
1	C	738	THR	2.1
1	C	725	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.