



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 25, 2024 – 04:36 PM EST

PDB ID : 2HPY
Title : Crystallographic model of lumirhodopsin
Authors : Nakamichi, H.; Okada, T.
Deposited on : 2006-07-18
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3657 (2.80-2.80)
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)
RSRZ outliers	164620	3659 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	349	4% (red), 38% (green), 53% (yellow), 9% (orange)
1	B	349	6% (red), 34% (green), 55% (yellow), 10% (orange)
2	C	3	33% (green), 33% (yellow), 33% (orange)
3	D	2	100% (yellow)
3	F	2	50% (green), 50% (orange)
4	E	4	50% (green), 50% (yellow)

2 Entry composition [i](#)


There are 11 unique types of molecules in this entry. The entry contains 5948 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rhodopsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	349	Total 2749	C 1818	N 424	O 481	S 26	0	0	0
1	B	349	Total 2749	C 1818	N 424	O 481	S 26	0	0	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	3	Total 39	C 22	N 2	O 15	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	2	Total 28	C 16	N 2	O 10	0	0	0
3	F	2	Total 28	C 16	N 2	O 10	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
4	E	4	Total	C	N	O	0	0	0
			50	28	2	20			


- Molecule 5 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Hg	0	0
			3	3		
5	B	3	Total	Hg	0	0
			3	3		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	4	Total	Zn	0	0
			4	4		
6	B	3	Total	Zn	0	0
			3	3		

- Molecule 7 is RETINAL (three-letter code: RET) (formula: C₂₀H₂₈O).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C 20 20	0	0
7	B	1	Total C 20 20	0	0

- Molecule 8 is PALMITIC ACID (three-letter code: PLM) (formula: C₁₆H₃₂O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 17 16 1	0	0
8	A	1	Total C O 17 16 1	0	0
8	A	1	Total C 16 16	0	0
8	B	1	Total C O 17 16 1	0	0
8	B	1	Total C O 17 16 1	0	0
8	B	1	Total C 16 16	0	0

- Molecule 9 is heptyl 1-thio-beta-D-glucopyranoside (three-letter code: HTG) (formula: C₁₃H₂₆O₅S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
9	A	1	19	13	5	1	0	0
9	A	1	19	13	5	1	0	0
9	B	1	19	13	5	1	0	0
9	B	1	19	13	5	1	0	0

- Molecule 10 is HEPTANE-1,2,3-TRIOL (three-letter code: HTO) (formula: $C_7H_{16}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	0	0
			10	7	3		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	37	Total	O	0	0
			37	37		
11	B	29	Total	O	0	0
			29	29		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Rhodopsin



- Molecule 1: Rhodopsin





- Molecule 2: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: beta-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics i

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	96.72Å 96.72Å 150.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 50.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.80) 87.6 (50.00-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 2.81Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.218 , 0.238 0.343 , 0.328	Depositor DCC
R_{free} test set	1530 reflections (5.24%)	wwPDB-VP
Wilson B-factor (Å ²)	64.8	Xtrriage
Anisotropy	0.474	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 242.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.138 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	5948	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RET, HTO, BMA, ACE, HG, HTG, ZN, NAG, MAN, PLM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/2831	0.70	0/3859
1	B	0.65	0/2831	0.68	0/3859
All	All	0.65	0/5662	0.69	0/7718

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	306	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2749	0	2709	257	1
1	B	2749	0	2709	327	0
2	C	39	0	34	1	0
3	D	28	0	25	0	0
3	F	28	0	25	1	0
4	E	50	0	43	0	0
5	A	3	0	0	0	0
5	B	3	0	0	0	0
6	A	4	0	0	0	0
6	B	3	0	0	0	0
7	A	20	0	27	2	0
7	B	20	0	27	2	0
8	A	50	0	89	7	0
8	B	50	0	89	4	0
9	A	38	0	52	3	0
9	B	38	0	52	4	1
10	B	10	0	16	1	0
11	A	37	0	0	6	0
11	B	29	0	0	10	0
All	All	5948	0	5897	584	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 50.

The worst 5 of 584 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298:SER:HA	1:A:301:TYR:CE2	1.82	1.15
1:B:298:SER:HA	1:B:301:TYR:CE2	1.81	1.15
1:A:67:LYS:HB2	1:A:337:VAL:HB	1.38	1.02
1:B:64:GLN:HG3	1:B:339:LYS:HE2	1.39	1.02
1:A:345:VAL:HG12	1:A:347:PRO:HD3	1.45	0.98

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:GLN:NE2	9:B:1509:HTG:O6[4_564]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/349 (99%)	264 (76%)	57 (16%)	26 (8%)	1	2
1	B	347/349 (99%)	254 (73%)	69 (20%)	24 (7%)	1	2
All	All	694/698 (99%)	518 (75%)	126 (18%)	50 (7%)	1	2

5 of 50 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	240	SER
1	A	241	ALA
1	A	323	CYS
1	A	328	LEU
1	A	341	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	296/296 (100%)	276 (93%)	20 (7%)	13	38
1	B	296/296 (100%)	273 (92%)	23 (8%)	10	31
All	All	592/592 (100%)	549 (93%)	43 (7%)	11	34

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	161	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	244	GLN
1	B	185	CYS
1	B	229	THR
1	B	249	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	8	ASN
1	B	64	GLN
1	B	302	ASN
1	B	237	GLN
1	B	244	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.63	0	17,19,21	0.85	0
2	NAG	C	2	2	14,14,15	0.77	0	17,19,21	0.93	1 (5%)
2	MAN	C	3	2	11,11,12	0.79	0	15,15,17	0.72	0
3	NAG	D	1	3,1	14,14,15	0.53	0	17,19,21	0.84	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	2	3	14,14,15	0.63	0	17,19,21	1.00	1 (5%)
4	NAG	E	1	4,1	14,14,15	0.60	0	17,19,21	0.67	0
4	NAG	E	2	4	14,14,15	0.65	0	17,19,21	0.96	1 (5%)
4	BMA	E	3	4	11,11,12	0.81	0	15,15,17	0.60	0
4	BMA	E	4	4	11,11,12	0.83	0	15,15,17	0.81	1 (6%)
3	NAG	F	1	3,1	14,14,15	0.64	0	17,19,21	0.84	1 (5%)
3	NAG	F	2	3	14,14,15	0.66	0	17,19,21	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	4/6/23/26	0/1/1/1
2	MAN	C	3	2	-	2/2/19/22	0/1/1/1
3	NAG	D	1	3,1	-	4/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
4	NAG	E	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	E	2	4	-	2/6/23/26	0/1/1/1
4	BMA	E	3	4	-	2/2/19/22	0/1/1/1
4	BMA	E	4	4	-	0/2/19/22	0/1/1/1
3	NAG	F	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2	NAG	C2-N2-C7	-3.20	118.61	122.90
4	E	2	NAG	C2-N2-C7	-2.81	119.13	122.90
3	D	1	NAG	C2-N2-C7	-2.77	119.18	122.90
4	E	4	BMA	C1-O5-C5	2.32	115.30	112.19
2	C	2	NAG	C8-C7-N2	2.24	119.83	116.12

There are no chirality outliers.

5 of 24 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
3	D	1	NAG	C8-C7-N2-C2
3	D	1	NAG	O7-C7-N2-C2
3	F	1	NAG	C8-C7-N2-C2


There are no ring outliers.


3 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	NAG	1	0
3	F	1	NAG	1	0
2	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 13 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
7	RET	A	1296	1	20,20,21	2.56	3 (15%)	27,27,28	2.03	7 (25%)
9	HTG	A	1508	-	19,19,19	2.48	8 (42%)	23,24,24	2.29	2 (8%)
8	PLM	B	1407	-	15,15,17	2.01	3 (20%)	14,14,17	3.41	8 (57%)
8	PLM	B	1322	1	16,16,17	1.04	1 (6%)	15,15,17	0.43	0
8	PLM	B	1323	1	16,16,17	1.01	1 (6%)	15,15,17	0.45	0
10	HTO	B	1401	-	9,9,9	1.81	2 (22%)	10,10,10	1.10	1 (10%)
7	RET	B	1296	1	20,20,21	2.30	4 (20%)	27,27,28	1.90	6 (22%)
8	PLM	A	1322	1	16,16,17	0.99	1 (6%)	15,15,17	0.61	0
9	HTG	A	1507	-	19,19,19	3.25	8 (42%)	23,24,24	1.99	2 (8%)
9	HTG	B	1506	-	19,19,19	3.06	8 (42%)	23,24,24	1.76	2 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	PLM	A	1410	-	15,15,17	2.07	3 (20%)	14,14,17	3.42	7 (50%)
9	HTG	B	1509	-	19,19,19	2.80	8 (42%)	23,24,24	1.98	1 (4%)
8	PLM	A	1323	1	16,16,17	1.00	1 (6%)	15,15,17	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	RET	A	1296	1	-	4/13/30/31	0/1/1/1
9	HTG	A	1508	-	-	4/10/30/30	0/1/1/1
8	PLM	B	1407	-	-	5/13/13/15	-
8	PLM	B	1322	1	-	11/14/14/15	-
8	PLM	B	1323	1	-	9/14/14/15	-
10	HTO	B	1401	-	-	2/10/10/10	-
7	RET	B	1296	1	-	5/13/30/31	0/1/1/1
8	PLM	A	1322	1	-	11/14/14/15	-
9	HTG	A	1507	-	-	5/10/30/30	0/1/1/1
9	HTG	B	1506	-	-	2/10/30/30	0/1/1/1
8	PLM	A	1410	-	-	5/13/13/15	-
9	HTG	B	1509	-	-	5/10/30/30	0/1/1/1
8	PLM	A	1323	1	-	10/14/14/15	-

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	1507	HTG	O5-C1	9.41	1.57	1.42
9	B	1506	HTG	O5-C1	8.98	1.56	1.42
9	B	1509	HTG	O5-C1	7.97	1.54	1.42
7	A	1296	RET	C1-C6	7.82	1.63	1.53
9	A	1508	HTG	O5-C1	7.05	1.53	1.42

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	1508	HTG	C1'-S1-C1	10.12	122.30	100.45
9	B	1509	HTG	C1'-S1-C1	8.41	118.62	100.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	1507	HTG	C1'-S1-C1	8.35	118.47	100.45
8	A	1410	PLM	CD-CC-CB	8.34	156.51	114.37
8	B	1407	PLM	CD-CC-CB	8.21	155.85	114.37

There are no chirality outliers.

5 of 78 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	1296	RET	C1-C6-C7-C8
7	A	1296	RET	C5-C6-C7-C8
7	B	1296	RET	C5-C6-C7-C8
9	B	1509	HTG	O5-C5-C6-O6
8	B	1407	PLM	CA-CB-CC-CD


There are no ring outliers.





12 monomers are involved in 23 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1296	RET	2	0
9	A	1508	HTG	1	0
8	B	1407	PLM	1	0
8	B	1322	PLM	2	0
8	B	1323	PLM	2	0
10	B	1401	HTO	1	0
7	B	1296	RET	2	0
8	A	1322	PLM	6	0
9	A	1507	HTG	2	0
9	B	1506	HTG	1	0
8	A	1410	PLM	1	0
9	B	1509	HTG	3	1


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Warning: The R factor obtained from EDS is 0.2774, which does not match the depositor's R factor of 0.218. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	348/349 (99%)	0.07	14 (4%) 43 35	19, 26, 135, 149	228 (65%)
1	B	348/349 (99%)	0.24	21 (6%) 29 22	19, 27, 154, 163	228 (65%)
All	All	696/698 (99%)	0.15	35 (5%) 35 28	19, 26, 147, 163	456 (65%)

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	146	PHE	6.9
1	B	229	THR	5.4
1	A	328	LEU	5.0
1	B	344	GLN	4.9
1	B	142	PRO	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
3	NAG	F	2	14/15	0.95	0.08	84,89,91,92	0
4	BMA	E	3	11/12	0.95	0.10	104,108,112,116	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	D	1	14/15	0.96	0.07	70,75,80,84	0
3	NAG	F	1	14/15	0.96	0.08	65,69,71,77	0
3	NAG	D	2	14/15	0.97	0.07	91,94,99,101	0
4	NAG	E	1	14/15	0.97	0.07	63,64,69,75	0
4	NAG	E	2	14/15	0.97	0.09	77,83,90,97	0
2	MAN	C	3	11/12	0.97	0.05	94,97,97,99	0
2	NAG	C	1	14/15	0.98	0.06	66,68,74,75	0
2	NAG	C	2	14/15	0.98	0.06	79,82,87,92	0
4	BMA	E	4	11/12	0.98	0.10	120,122,122,124	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	PLM	A	1323	17/18	0.94	0.08	98,105,113,114	0
6	ZN	A	962	1/1	0.95	0.11	80,80,80,80	1
8	PLM	B	1407	16/18	0.95	0.12	75,82,87,88	0
9	HTG	B	1506	19/19	0.95	0.09	88,93,97,99	0
10	HTO	B	1401	10/10	0.95	0.10	56,60,62,63	0
8	PLM	A	1322	17/18	0.96	0.15	94,104,110,110	0
8	PLM	A	1410	16/18	0.96	0.08	74,77,79,79	0
8	PLM	B	1323	17/18	0.96	0.11	117,122,124,124	0
6	ZN	B	963	1/1	0.97	0.14	93,93,93,93	1
7	RET	A	1296	20/21	0.97	0.14	50,52,54,55	20
9	HTG	A	1508	19/19	0.97	0.07	72,78,82,85	0
7	RET	B	1296	20/21	0.97	0.12	51,52,54,54	20
9	HTG	B	1509	19/19	0.97	0.10	93,95,100,100	0
8	PLM	B	1322	17/18	0.97	0.09	105,108,109,110	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	HG	A	905	1/1	0.98	0.12	99,99,99,99	1
9	HTG	A	1507	19/19	0.98	0.10	99,113,115,116	0
6	ZN	A	2011	1/1	0.98	0.04	53,53,53,53	1
6	ZN	B	956	1/1	0.99	0.03	60,60,60,60	1
5	HG	B	902	1/1	0.99	0.04	80,80,80,80	1
5	HG	B	906	1/1	0.99	0.04	121,121,121,121	1
5	HG	A	903	1/1	0.99	0.05	96,96,96,96	1
5	HG	A	901	1/1	0.99	0.05	82,82,82,82	1
6	ZN	A	959	1/1	1.00	0.02	92,92,92,92	1
6	ZN	B	958	1/1	1.00	0.02	43,43,43,43	1
5	HG	B	904	1/1	1.00	0.03	109,109,109,109	1
6	ZN	A	957	1/1	1.00	0.02	53,53,53,53	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





Electron density around PLM B 1407:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around HTG B 1506:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLM A 1322:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PLM A 1410:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLM B 1323:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around RET A 1296:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HTG A 1508:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around RET B 1296:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around HTG B 1509:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around PLM B 1322:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.