



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 04:56 am BST

PDB ID : 6HPS  
Title : Near-infrared dual bioluminescence imaging in vivo using infra-luciferin  
Authors : Parkinson, G.N.; Stowe, C.; Anderson, J.C.  
Deposited on : 2018-09-21  
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.


Mol	Chain	Length	Quality of chain
1	A	543	 4% 89% 8% ..
1	B	543	 3% 88% 10% .




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Luciferin 4-monooxygenase



- Molecule 1: Luciferin 4-monooxygenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.51Å 114.02Å 101.56Å 90.00° 98.18° 90.00°	Depositor
Resolution (Å)	17.86 – 3.10 17.85 – 3.10	Depositor EDS
% Data completeness (in resolution range)	94.1 (17.86-3.10) 94.6 (17.85-3.10)	Depositor EDS
$R_{merge}$	0.23	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.80 (at 3.08Å)	Xtrriage
Refinement program	REFMAC 5.8.0222	Depositor
R, $R_{free}$	0.277 , 0.333 0.280 , 0.336	Depositor DCC
$R_{free}$ test set	924 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.2	Xtrriage
Anisotropy	1.865	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 40.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	8510	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 55.00 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.3700e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GKH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/4297	0.87	2/5822 (0.0%)
1	B	0.56	0/4319	0.88	2/5850 (0.0%)
All	All	0.56	0/8616	0.87	4/11672 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	358	LYS	C-N-CD	-13.69	90.48	120.60
1	B	359	PRO	CA-N-CD	-5.90	103.24	111.50
1	B	395	PRO	CA-N-CD	-5.19	104.23	111.50
1	A	173	PRO	CA-N-CD	-5.12	104.33	111.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	330	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4202	0	4245	27	0
1	B	4224	0	4269	28	0
2	A	42	0	0	1	0
2	B	42	0	0	1	0
All	All	8510	0	8514	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:290:THR:HG21	1:B:524:LYS:HD2	1.56	0.87
1:B:525:GLY:O	1:B:527:THR:N	2.15	0.78
1:B:298:SER:O	1:B:330:ARG:NH1	2.19	0.75
1:A:108:ILE:HD13	1:A:526:LEU:HD13	1.74	0.69
1:B:469:VAL:HG22	1:B:483:ALA:HB2	1.79	0.63
1:B:346:THR:OG1	1:B:396:MET:O	2.17	0.62
1:B:273:PHE:CE2	1:B:277:LEU:HD11	2.36	0.61
1:B:290:THR:CG2	1:B:524:LYS:HD2	2.28	0.60
1:A:523:PRO:O	1:A:530:LEU:HD22	2.01	0.60
1:A:358:LYS:HB3	1:A:361:ALA:HB3	1.84	0.60
1:A:526:LEU:HD23	1:A:526:LEU:O	2.05	0.57
1:B:168:VAL:HG13	1:B:172:LEU:HD12	1.87	0.56
1:B:330:ARG:CZ	1:B:330:ARG:HB3	2.34	0.56
1:B:84:ASN:O	1:B:266:TYR:CD1	2.60	0.55
1:B:198:SER:HB2	1:B:208:VAL:CG2	2.38	0.54
1:A:116:ASN:ND2	1:A:204:LEU:HB2	2.24	0.54
1:A:106:ASN:HB3	1:A:109:TYR:CG	2.45	0.52
1:B:311:GLU:OE1	1:B:337:ARG:NH1	2.43	0.52
1:A:522:VAL:CG1	1:A:530:LEU:HD21	2.41	0.51
1:A:351:ILE:O	1:A:353:PRO:HD3	2.13	0.49
1:A:290:THR:CB	1:A:524:LYS:HD2	2.43	0.48
1:B:290:THR:HG21	1:B:524:LYS:CD	2.35	0.48
1:A:522:VAL:HG12	1:A:530:LEU:HD21	1.95	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:ASN:HB3	1:B:109:TYR:CG	2.49	0.47
1:B:107:ASP:OD1	1:B:107:ASP:N	2.41	0.47
1:A:290:THR:CG2	1:A:528:GLY:O	2.62	0.47
1:B:198:SER:HB2	1:B:208:VAL:HG23	1.97	0.47
1:A:290:THR:HG21	1:A:524:LYS:HG3	1.97	0.47
1:B:192:ILE:HD12	1:B:209:ALA:HB1	1.96	0.46
1:B:273:PHE:CZ	1:B:287:LEU:HD21	2.50	0.46
1:B:206:LYS:HD3	1:B:401:TYR:CE1	2.51	0.46
1:A:339:GLY:O	2:A:601:GKH:N6	2.49	0.46
1:A:74:THR:HA	1:A:99:GLY:O	2.16	0.45
1:B:423:ILE:HB	1:B:435:VAL:HB	1.99	0.45
1:A:290:THR:OG1	1:A:524:LYS:HD2	2.17	0.45
1:A:482:ALA:HB2	1:A:514:GLY:HA3	2.00	0.44
1:B:525:GLY:N	1:B:529:LYS:O	2.51	0.44
1:A:290:THR:HG23	1:A:528:GLY:O	2.19	0.43
1:A:20:GLY:O	1:A:188:ARG:NH1	2.39	0.43
1:B:237:ILE:N	1:B:237:ILE:HD12	2.34	0.42
1:B:194:LEU:HD13	1:B:250:PHE:CZ	2.54	0.42
1:A:273:PHE:CE2	1:A:277:LEU:HD11	2.55	0.42
1:A:290:THR:HG21	1:A:524:LYS:CG	2.49	0.42
1:A:358:LYS:HA	1:A:359:PRO:HD2	1.61	0.42
1:B:339:GLY:O	2:B:601:GKH:N6	2.53	0.41
1:A:423:ILE:HD11	1:A:438:LEU:HD13	2.02	0.41
1:A:289:PRO:HG2	1:A:317:ALA:HB3	2.03	0.41
1:A:79:VAL:HG23	1:A:123:PRO:HB3	2.03	0.41
1:B:290:THR:OG1	1:B:524:LYS:HD2	2.20	0.41
1:A:240:VAL:HG22	1:A:240:VAL:O	2.20	0.41
1:A:109:TYR:CD1	1:A:113:GLU:HB3	2.56	0.41
1:B:506:VAL:HG13	1:B:507:THR:N	2.36	0.41
1:B:290:THR:CB	1:B:524:LYS:HD2	2.50	0.40
1:B:322:GLU:HG2	1:B:460:GLN:HE21	1.86	0.40
1:A:290:THR:HB	1:A:524:LYS:HD2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries



of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	537/543 (99%)	492 (92%)	39 (7%)	6 (1%)	14	46
1	B	541/543 (100%)	481 (89%)	51 (9%)	9 (2%)	9	36
All	All	1078/1086 (99%)	973 (90%)	90 (8%)	15 (1%)	11	40

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	298	SER
1	A	359	PRO
1	A	475	ASP
1	B	359	PRO
1	B	452	ALA
1	B	526	LEU
1	B	296	ALA
1	B	384	VAL
1	A	201	SER
1	A	444	TYR
1	A	309	LEU
1	B	159	GLN
1	B	205	PRO
1	B	445	LYS
1	B	506	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	455/456 (100%)	435 (96%)	20 (4%)	28	61
1	B	456/456 (100%)	429 (94%)	27 (6%)	19	50
All	All	911/912 (100%)	864 (95%)	47 (5%)	23	55

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ASP
1	A	56	TYR
1	A	83	GLU
1	A	109	TYR
1	A	184	GLU
1	A	197	ASN
1	A	204	LEU
1	A	289	PRO
1	A	302	ASP
1	A	340	TYR
1	A	349	ILE
1	A	406	GLU
1	A	427	ASP
1	A	443	LYS
1	A	505	GLN
1	A	523	PRO
1	A	526	LEU
1	A	527	THR
1	A	530	LEU
1	A	533	ARG
1	B	14	PHE
1	B	19	ASP
1	B	56	TYR
1	B	107	ASP
1	B	108	ILE
1	B	109	TYR
1	B	119	ASN
1	B	124	THR
1	B	155	LYS
1	B	163	SER
1	B	169	THR
1	B	201	SER
1	B	205	PRO
1	B	206	LYS
1	B	298	SER
1	B	330	ARG
1	B	357	ASP
1	B	427	ASP
1	B	438	LEU
1	B	453	GLU
1	B	475	ASP
1	B	508	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	520	ASP
1	B	521	GLU
1	B	523	PRO
1	B	524	LYS
1	B	526	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	ASN
1	A	448	GLN
1	B	25	GLN
1	B	116	ASN
1	B	119	ASN
1	B	460	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GKH	A	601	-	42,47,47	4.78	19 (45%)	43,70,70	2.60	16 (37%)
2	GKH	B	601	-	42,47,47	4.51	19 (45%)	43,70,70	2.58	17 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GKH	A	601	-	-	5/11/40/40	0/6/6/6
2	GKH	B	601	-	-	4/11/40/40	0/6/6/6

All (38) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	GKH	CAK-SBN	-19.39	1.46	1.73
2	A	601	GKH	CAK-SBN	-18.05	1.48	1.73
2	A	601	GKH	OBJ-SAP	15.90	1.55	1.42
2	B	601	GKH	OBJ-SAP	10.73	1.51	1.42
2	B	601	GKH	CAH-SAG	-9.60	1.60	1.73
2	A	601	GKH	OBK-SAP	9.17	1.50	1.42
2	A	601	GKH	CAH-SAG	-8.20	1.62	1.73
2	B	601	GKH	OBK-SAP	8.03	1.49	1.42
2	A	601	GKH	CAN-NAO	-6.54	1.31	1.39
2	A	601	GKH	CAM-CAN	-5.63	1.36	1.50
2	B	601	GKH	CAM-CAN	-4.91	1.38	1.50
2	B	601	GKH	O4'-C1'	4.69	1.47	1.41
2	B	601	GKH	CAK-NAL	4.55	1.38	1.31
2	A	601	GKH	CAK-NAL	4.27	1.37	1.31
2	A	601	GKH	O4'-C1'	4.17	1.46	1.41
2	A	601	GKH	C2-N3	3.88	1.38	1.32
2	B	601	GKH	C2-N1	3.85	1.41	1.33
2	A	601	GKH	CAJ-CAI	3.83	1.50	1.31
2	B	601	GKH	CAJ-CAI	3.76	1.50	1.31
2	B	601	GKH	O5'-SAP	-3.69	1.52	1.59
2	B	601	GKH	CAN-NAO	-3.53	1.34	1.39
2	B	601	GKH	C2-N3	3.30	1.37	1.32
2	A	601	GKH	CAH-CAI	-3.30	1.33	1.47
2	B	601	GKH	CAH-CAI	-3.18	1.33	1.47
2	A	601	GKH	SAP-NAO	3.15	1.65	1.59
2	B	601	GKH	CAK-CAJ	-3.15	1.33	1.47

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	GKH	CAK-CAJ	-3.09	1.34	1.47
2	B	601	GKH	CAD-CAE	-2.99	1.36	1.41
2	A	601	GKH	CAD-CAE	-2.91	1.36	1.41
2	B	601	GKH	C6-C5	-2.57	1.33	1.43
2	B	601	GKH	CAD-CAC	2.54	1.41	1.36
2	A	601	GKH	C2-N1	2.52	1.38	1.33
2	A	601	GKH	CAD-CAC	2.52	1.41	1.36
2	B	601	GKH	C5-C4	-2.44	1.34	1.40
2	A	601	GKH	O5'-SAP	-2.40	1.54	1.59
2	A	601	GKH	C5-C4	-2.38	1.34	1.40
2	A	601	GKH	C6-C5	-2.34	1.34	1.43
2	B	601	GKH	CAH-NBO	2.25	1.34	1.31

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	GKH	OBK-SAP-OBJ	-10.80	103.94	120.76
2	B	601	GKH	OBK-SAP-OBJ	-9.84	105.43	120.76
2	B	601	GKH	N3-C2-N1	-6.57	118.41	128.68
2	A	601	GKH	N3-C2-N1	-6.17	119.04	128.68
2	B	601	GKH	CAM-CBM-SBN	-4.22	106.61	111.79
2	A	601	GKH	CAM-CAN-NAO	3.92	118.62	114.03
2	B	601	GKH	C5'-O5'-SAP	3.60	124.80	117.37
2	B	601	GKH	CAM-CAN-NAO	3.45	118.07	114.03
2	A	601	GKH	CAM-CBM-SBN	-3.43	107.58	111.79
2	A	601	GKH	CAK-CAJ-CAI	-3.26	116.72	123.69
2	B	601	GKH	CAK-CAJ-CAI	-3.12	117.02	123.69
2	B	601	GKH	C5-C6-N6	-2.87	115.99	120.35
2	A	601	GKH	C1'-N9-C4	-2.79	121.74	126.64
2	B	601	GKH	N6-C6-N1	2.79	124.36	118.57
2	A	601	GKH	C2'-C3'-C4'	-2.77	97.26	102.64
2	B	601	GKH	CAH-CAI-CAJ	-2.76	117.78	123.69
2	A	601	GKH	OBL-CAN-NAO	-2.65	117.84	121.08
2	A	601	GKH	O4'-C1'-C2'	-2.58	103.15	106.93
2	B	601	GKH	CAD-CAE-CAF	-2.54	116.07	120.53
2	A	601	GKH	CAF-CAE-NBO	2.49	113.83	108.04
2	A	601	GKH	CAC-CAB-CBP	-2.46	117.25	120.39
2	B	601	GKH	CAF-CAE-NBO	2.45	113.73	108.04
2	B	601	GKH	O4'-C1'-C2'	-2.36	103.48	106.93
2	A	601	GKH	C5'-O5'-SAP	2.36	122.24	117.37
2	B	601	GKH	C3'-C2'-C1'	2.31	104.45	100.98
2	A	601	GKH	C5-C6-N6	-2.27	116.90	120.35

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	GKH	O4'-C4'-C3'	-2.24	100.69	105.11
2	B	601	GKH	C2'-C3'-C4'	-2.20	98.37	102.64
2	B	601	GKH	CAC-CAB-CBP	-2.19	117.59	120.39
2	A	601	GKH	O5'-SAP-OBJ	2.17	112.23	105.59
2	B	601	GKH	CAE-CAF-SAG	-2.16	108.99	111.85
2	A	601	GKH	CAD-CAE-CAF	-2.11	116.81	120.53
2	B	601	GKH	C1'-N9-C4	-2.05	123.05	126.64

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	GKH	CAM-CAN-NAO-SAP
2	A	601	GKH	OBL-CAN-NAO-SAP
2	A	601	GKH	CAN-NAO-SAP-OBJ
2	A	601	GKH	C3'-C4'-C5'-O5'
2	B	601	GKH	CAN-NAO-SAP-OBJ
2	B	601	GKH	CAM-CAN-NAO-SAP
2	B	601	GKH	OBL-CAN-NAO-SAP
2	A	601	GKH	O4'-C4'-C5'-O5'
2	B	601	GKH	C3'-C4'-C5'-O5'


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	GKH	1	0
2	B	601	GKH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	539/543 (99%)	0.41	24 (4%) 33 16	16, 35, 61, 99	0
1	B	543/543 (100%)	0.40	17 (3%) 49 26	20, 35, 63, 93	0
All	All	1082/1086 (99%)	0.41	41 (3%) 40 20	16, 35, 63, 99	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	203	GLY	6.0
1	B	204	LEU	5.2
1	B	490	GLY	4.9
1	A	299	THR	3.8
1	B	283	GLN	3.8
1	A	204	LEU	3.7
1	B	160	GLY	3.6
1	A	283	GLN	3.5
1	A	302	ASP	3.5
1	A	298	SER	3.3
1	A	202	THR	3.2
1	A	355	GLY	3.1
1	B	201	SER	2.8
1	A	490	GLY	2.7
1	A	476	ASP	2.7
1	A	15	TYR	2.7
1	B	202	THR	2.6
1	A	266	TYR	2.6
1	A	489	HIS	2.6
1	B	359	PRO	2.6
1	B	71	GLY	2.5
1	A	508	THR	2.5
1	B	508	THR	2.4
1	B	489	HIS	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	174	PRO	2.3
1	B	12	ALA	2.3
1	A	402	VAL	2.3
1	B	355	GLY	2.2
1	A	106	ASN	2.2
1	A	175	GLY	2.2
1	B	234	ASP	2.2
1	A	290	THR	2.2
1	B	528	GLY	2.2
1	A	354	GLU	2.2
1	B	269	GLU	2.1
1	A	360	GLY	2.1
1	B	460	GLN	2.1
1	B	203	GLY	2.1
1	A	334	PRO	2.1
1	A	494	THR	2.0
1	A	199	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GKH	B	601	42/42	0.86	0.27	17,24,29,31	0
2	GKH	A	601	42/42	0.90	0.23	15,21,23,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around GKH B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GKH A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.