



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:40 PM UTC

PDB ID : 9HTX / pdb\_00009htx  
Title : Glycosyltransferase C from the *Limosilactobacillus reuteri* accessory secretion system. Apo form.  
Authors : Pfalzgraf, H.E.; Griffiths, R.; Juge, N.; Hemmings, A.M.  
Deposited on : 2024-12-20  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	<div> <div>6%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>• 5%</div> </div> </div>
1	B	354	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>•• 5%</div> </div> </div>
1	C	354	<div> <div>3%</div> <div> <div></div> <div>81%</div> <div>11%</div> <div>•• 5%</div> </div> </div>
1	D	354	<div> <div>9%</div> <div> <div></div> <div>80%</div> <div>13%</div> <div>•• 5%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 21219 atoms, of which 10218 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucosyltransferase 3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	335	Total	C	H	N	O	S	0	0	0
			5255	1744	2552	449	503	7			
1	B	338	Total	C	H	N	O	S	0	0	0
			5303	1757	2576	455	508	7			
1	C	335	Total	C	H	N	O	S	0	0	0
			5244	1736	2548	452	502	6			
1	D	336	Total	C	H	N	O	S	0	0	0
			5206	1725	2522	449	503	7			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP B3XPQ7
A	-18	GLY	-	expression tag	UNP B3XPQ7
A	-17	SER	-	expression tag	UNP B3XPQ7
A	-16	SER	-	expression tag	UNP B3XPQ7
A	-15	HIS	-	expression tag	UNP B3XPQ7
A	-14	HIS	-	expression tag	UNP B3XPQ7
A	-13	HIS	-	expression tag	UNP B3XPQ7
A	-12	HIS	-	expression tag	UNP B3XPQ7
A	-11	HIS	-	expression tag	UNP B3XPQ7
A	-10	HIS	-	expression tag	UNP B3XPQ7
A	-9	SER	-	expression tag	UNP B3XPQ7
A	-8	SER	-	expression tag	UNP B3XPQ7
A	-7	GLY	-	expression tag	UNP B3XPQ7
A	-6	LEU	-	expression tag	UNP B3XPQ7
A	-5	VAL	-	expression tag	UNP B3XPQ7
A	-4	PRO	-	expression tag	UNP B3XPQ7
A	-3	ARG	-	expression tag	UNP B3XPQ7
A	-2	GLY	-	expression tag	UNP B3XPQ7
A	-1	SER	-	expression tag	UNP B3XPQ7
A	0	HIS	-	expression tag	UNP B3XPQ7
A	1	LEU	-	expression tag	UNP B3XPQ7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	initiating methionine	UNP B3XPQ7
B	-18	GLY	-	expression tag	UNP B3XPQ7
B	-17	SER	-	expression tag	UNP B3XPQ7
B	-16	SER	-	expression tag	UNP B3XPQ7
B	-15	HIS	-	expression tag	UNP B3XPQ7
B	-14	HIS	-	expression tag	UNP B3XPQ7
B	-13	HIS	-	expression tag	UNP B3XPQ7
B	-12	HIS	-	expression tag	UNP B3XPQ7
B	-11	HIS	-	expression tag	UNP B3XPQ7
B	-10	HIS	-	expression tag	UNP B3XPQ7
B	-9	SER	-	expression tag	UNP B3XPQ7
B	-8	SER	-	expression tag	UNP B3XPQ7
B	-7	GLY	-	expression tag	UNP B3XPQ7
B	-6	LEU	-	expression tag	UNP B3XPQ7
B	-5	VAL	-	expression tag	UNP B3XPQ7
B	-4	PRO	-	expression tag	UNP B3XPQ7
B	-3	ARG	-	expression tag	UNP B3XPQ7
B	-2	GLY	-	expression tag	UNP B3XPQ7
B	-1	SER	-	expression tag	UNP B3XPQ7
B	0	HIS	-	expression tag	UNP B3XPQ7
B	1	LEU	-	expression tag	UNP B3XPQ7
C	-19	MET	-	initiating methionine	UNP B3XPQ7
C	-18	GLY	-	expression tag	UNP B3XPQ7
C	-17	SER	-	expression tag	UNP B3XPQ7
C	-16	SER	-	expression tag	UNP B3XPQ7
C	-15	HIS	-	expression tag	UNP B3XPQ7
C	-14	HIS	-	expression tag	UNP B3XPQ7
C	-13	HIS	-	expression tag	UNP B3XPQ7
C	-12	HIS	-	expression tag	UNP B3XPQ7
C	-11	HIS	-	expression tag	UNP B3XPQ7
C	-10	HIS	-	expression tag	UNP B3XPQ7
C	-9	SER	-	expression tag	UNP B3XPQ7
C	-8	SER	-	expression tag	UNP B3XPQ7
C	-7	GLY	-	expression tag	UNP B3XPQ7
C	-6	LEU	-	expression tag	UNP B3XPQ7
C	-5	VAL	-	expression tag	UNP B3XPQ7
C	-4	PRO	-	expression tag	UNP B3XPQ7
C	-3	ARG	-	expression tag	UNP B3XPQ7
C	-2	GLY	-	expression tag	UNP B3XPQ7
C	-1	SER	-	expression tag	UNP B3XPQ7
C	0	HIS	-	expression tag	UNP B3XPQ7
C	1	LEU	-	expression tag	UNP B3XPQ7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-19	MET	-	initiating methionine	UNP B3XPQ7
D	-18	GLY	-	expression tag	UNP B3XPQ7
D	-17	SER	-	expression tag	UNP B3XPQ7
D	-16	SER	-	expression tag	UNP B3XPQ7
D	-15	HIS	-	expression tag	UNP B3XPQ7
D	-14	HIS	-	expression tag	UNP B3XPQ7
D	-13	HIS	-	expression tag	UNP B3XPQ7
D	-12	HIS	-	expression tag	UNP B3XPQ7
D	-11	HIS	-	expression tag	UNP B3XPQ7
D	-10	HIS	-	expression tag	UNP B3XPQ7
D	-9	SER	-	expression tag	UNP B3XPQ7
D	-8	SER	-	expression tag	UNP B3XPQ7
D	-7	GLY	-	expression tag	UNP B3XPQ7
D	-6	LEU	-	expression tag	UNP B3XPQ7
D	-5	VAL	-	expression tag	UNP B3XPQ7
D	-4	PRO	-	expression tag	UNP B3XPQ7
D	-3	ARG	-	expression tag	UNP B3XPQ7
D	-2	GLY	-	expression tag	UNP B3XPQ7
D	-1	SER	-	expression tag	UNP B3XPQ7
D	0	HIS	-	expression tag	UNP B3XPQ7
D	1	LEU	-	expression tag	UNP B3XPQ7

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	0	0
			11	3	5	3		
3	B	1	Total	C	H	O	0	0
			11	3	5	3		
3	B	1	Total	C	H	O	0	0
			11	3	5	3		
3	C	1	Total	C	H	O	0	0
			11	3	5	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	25	Total	O	0	0
			25	25		
4	B	70	Total	O	0	0
			70	70		
4	C	59	Total	O	0	0
			59	59		
4	D	11	Total	O	0	0
			11	11		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucosyltransferase 3





- Molecule 1: Glucosyltransferase 3



- Molecule 1: Glucosyltransferase 3



- Molecule 1: Glucosyltransferase 3





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.44Å 140.49Å 145.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.77 – 2.00 72.77 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (72.77-2.00) 88.8 (72.77-2.00)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.12 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.8.0430 (refmacat 0.4.105)	Depositor
R, $R_{free}$	0.207 , 0.253 0.202 , 0.248	Depositor DCC
$R_{free}$ test set	5089 reflections (3.80%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.2	Xtriage
Anisotropy	0.459	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 48.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	21219	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.84	0/2776	1.37	21/3785 (0.6%)
1	B	1.10	5/2800 (0.2%)	1.49	23/3814 (0.6%)
1	C	1.08	2/2768 (0.1%)	1.49	26/3770 (0.7%)
1	D	0.85	1/2754 (0.0%)	1.36	20/3758 (0.5%)
All	All	0.98	8/11098 (0.1%)	1.43	90/15127 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
1	C	0	5
All	All	0	10

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	100	HIS	CG-CD2	-8.19	1.26	1.35
1	D	45	ASP	CG-OD1	7.47	1.39	1.25
1	B	253	SER	CA-CB	7.17	1.64	1.53
1	B	313	ARG	C-O	-5.74	1.17	1.24
1	B	230	PHE	C-N	5.68	1.36	1.33

The worst 5 of 90 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	225	ARG	CD-NE-CZ	15.88	146.63	124.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	227	ASN	CB-CA-C	12.38	133.48	109.72
1	D	334	TYR	CA-C-O	-9.39	104.83	120.80
1	C	313	ARG	NE-CZ-NH2	9.07	127.37	119.20
1	B	1	LEU	CD1-CG-CD2	-8.84	91.34	110.80

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	158	ALA	Peptide
1	A	313	ARG	Sidechain
1	B	1	LEU	Peptide
1	B	225	ARG	Sidechain
1	B	313	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2703	2552	2614	14	1
1	B	2727	2576	2639	21	0
1	C	2696	2548	2608	21	0
1	D	2684	2522	2585	20	1
2	A	1	0	0	0	0
2	C	1	0	0	0	0
3	B	18	15	24	2	0
3	C	6	5	8	3	0
4	A	25	0	0	0	0
4	B	70	0	0	1	0
4	C	59	0	0	0	0
4	D	11	0	0	0	0
All	All	11001	10218	10478	70	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 70 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:53:THR:HG21	1:D:50:ALA:O	1.60	1.02
1:C:53:THR:CG2	1:D:50:ALA:O	2.23	0.87
1:C:250:HIS:HB3	3:C:402:GOL:O1	1.90	0.71
1:C:102:ILE:HG22	1:C:104:PRO:HD2	1.72	0.71
1:B:52:ASP:HB3	3:B:401:GOL:H12	1.73	0.70

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:GLU:OE1	1:D:239:TYR:H[3_545]	1.60	0.00

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/354 (94%)	326 (98%)	6 (2%)	1 (0%)	36	35
1	B	336/354 (95%)	325 (97%)	10 (3%)	1 (0%)	36	35
1	C	331/354 (94%)	323 (98%)	7 (2%)	1 (0%)	36	35
1	D	334/354 (94%)	324 (97%)	9 (3%)	1 (0%)	36	35
All	All	1334/1416 (94%)	1298 (97%)	32 (2%)	4 (0%)	36	35

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	77	SER
1	B	77	SER
1	D	77	SER
1	C	77	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/309 (94%)	282 (97%)	9 (3%)	35	37
1	B	293/309 (95%)	288 (98%)	5 (2%)	53	60
1	C	290/309 (94%)	279 (96%)	11 (4%)	29	29
1	D	287/309 (93%)	267 (93%)	20 (7%)	14	10
All	All	1161/1236 (94%)	1116 (96%)	45 (4%)	28	28

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	93	VAL
1	D	160	VAL
1	D	105	LEU
1	D	143	LEU
1	D	179	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	132	GLN
1	C	306	GLN
1	C	245	HIS
1	D	34	ASN
1	B	132	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	C	402	-	5,5,5	0.31	0	5,5,5	0.58	0
3	GOL	B	401	-	5,5,5	0.48	0	5,5,5	1.21	1 (20%)
3	GOL	B	402	-	5,5,5	0.40	0	5,5,5	0.91	0
3	GOL	B	403	-	5,5,5	0.31	0	5,5,5	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	402	-	-	4/4/4/4	-
3	GOL	B	401	-	-	2/4/4/4	-
3	GOL	B	402	-	-	2/4/4/4	-
3	GOL	B	403	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	GOL	O1-C1-C2	2.27	120.60	110.38

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	402	GOL	C1-C2-C3-O3
3	C	402	GOL	O2-C2-C3-O3
3	B	401	GOL	C1-C2-C3-O3
3	B	402	GOL	C1-C2-C3-O3
3	B	403	GOL	O1-C1-C2-C3

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	402	GOL	3	0
3	B	401	GOL	1	0
3	B	402	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	335/354 (94%)	0.55	23 (6%)	23 21	37, 76, 138, 174	0
1	B	338/354 (95%)	0.14	10 (2%)	52 51	33, 53, 103, 157	0
1	C	335/354 (94%)	0.07	9 (2%)	56 55	34, 52, 89, 150	0
1	D	336/354 (94%)	0.76	31 (9%)	14 13	43, 85, 140, 199	0
All	All	1344/1416 (94%)	0.38	73 (5%)	31 30	33, 64, 127, 199	0

The worst 5 of 73 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	183	LEU	4.7
1	D	183	LEU	4.6
1	A	184	VAL	4.5
1	D	187	TRP	4.4
1	D	178	LEU	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	B	403	6/6	0.83	0.15	60,69,90,96	0
3	GOL	B	401	6/6	0.86	0.12	60,66,75,75	0
3	GOL	C	402	6/6	0.90	0.13	69,80,82,88	0
3	GOL	B	402	6/6	0.91	0.14	50,74,85,98	0
2	CA	C	401	1/1	0.97	0.06	50,50,50,50	0
2	CA	A	401	1/1	0.99	0.04	45,45,45,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.