



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 04:59 AM EDT

PDB ID : 5HUE  
Title : DAHP synthase from *Corynebacterium glutamicum* in complex with tryptophan  
Authors : Burschowsky, D.; Heim, J.B.; Thorbjørnsrud, H.V.; Kregel, U.  
Deposited on : 2016-01-27  
Resolution : 2.65 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	472	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MPD	A	505	-	-	-	X

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 3571 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-Deoxy-D-arabino-heptulosonate 7-phosphate (DAHP) synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	447	3456	2151	614	674	17	0	0	0


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q8NNL5
A	2	HIS	-	expression tag	UNP Q8NNL5
A	3	HIS	-	expression tag	UNP Q8NNL5
A	4	HIS	-	expression tag	UNP Q8NNL5
A	5	HIS	-	expression tag	UNP Q8NNL5
A	6	HIS	-	expression tag	UNP Q8NNL5
A	7	HIS	-	expression tag	UNP Q8NNL5
A	8	SER	-	expression tag	UNP Q8NNL5
A	9	SER	-	expression tag	UNP Q8NNL5
A	10	GLY	-	expression tag	UNP Q8NNL5

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mn	0	0
			1	1		

- Molecule 3 is PHOSPHOENOLPYRUVATE (three-letter code: PEP) (formula: C<sub>3</sub>H<sub>5</sub>O<sub>6</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 4 is TRYPTOPHAN (three-letter code: TRP) (formula:  $C_{11}H_{12}N_2O_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			15	11	2	2		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula:  $C_6H_{14}O_2$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 8 6 2	0	0
5	A	1	Total C O 8 6 2	0	0
5	A	1	Total C O 8 6 2	0	0
5	A	1	Total C O 8 6 2	0	0
5	A	1	Total C O 8 6 2	0	0

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			5	4	1		
7	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 8 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
8	A	1	8	4	1	3	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
9	A	21	21	21	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-Deoxy-D-arabino-heptulosonate 7-phosphate (DAHP) synthase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	109.19Å 109.19Å 279.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.01 – 2.65 36.83 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.01-2.65) 99.9 (36.83-2.65)	Depositor EDS
$R_{merge}$	0.28	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.15 (at 2.65Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.247 , 0.285 0.251 , 0.282	Depositor DCC
$R_{free}$ test set	1436 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	72.2	Xtrriage
Anisotropy	0.473	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 66.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3571	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	109.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, PEP, TRS, PO4, MN, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3525	0.64	1/4780 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	28	LEU	CA-CB-CG	5.75	128.53	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3456	0	3361	7	0
2	A	1	0	0	0	0
3	A	10	0	2	0	0
4	A	15	0	9	0	0
5	A	40	0	70	5	0
6	A	10	0	14	0	0
7	A	10	0	0	0	0
8	A	8	0	12	0	0
9	A	21	0	0	0	0
All	All	3571	0	3468	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:THR:HG21	5:A:506:MPD:H51	1.83	0.60
1:A:143:LYS:HG2	1:A:450:CYS:SG	2.44	0.57
1:A:313:ILE:HD12	1:A:341:ARG:HE	1.71	0.56
1:A:42:ARG:O	5:A:504:MPD:HM1	2.12	0.49
5:A:504:MPD:H52	5:A:508:MPD:H4	1.98	0.46
1:A:309:ILE:O	1:A:341:ARG:NH1	2.52	0.43
5:A:504:MPD:H52	5:A:504:MPD:H11	2.01	0.43
1:A:113:VAL:HG13	1:A:253:ILE:CD1	2.49	0.42
1:A:127:ALA:HB2	1:A:470:LEU:HD13	2.02	0.42
5:A:504:MPD:H11	5:A:508:MPD:H52	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	445/472 (94%)	425 (96%)	17 (4%)	3 (1%)	22 33

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	30	GLU
1	A	29	PRO
1	A	144	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	365/389 (94%)	346 (95%)	19 (5%)	23 36

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LEU
1	A	50	TRP
1	A	92	LEU
1	A	102	GLU
1	A	133	LYS
1	A	134	MET
1	A	145	ARG
1	A	154	LEU
1	A	249	ARG
1	A	260	LEU
1	A	261	LEU
1	A	275	GLU
1	A	290	TRP
1	A	317	ILE
1	A	331	ASP
1	A	341	ARG
1	A	347	ARG
1	A	395	ASP
1	A	453	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	MPD	A	506	-	7,7,7	0.30	0	9,10,10	0.45	0
7	PO4	A	510	-	4,4,4	0.96	0	6,6,6	1.13	1 (16%)
7	PO4	A	511	-	4,4,4	1.01	0	6,6,6	0.54	0
5	MPD	A	508	-	7,7,7	0.31	0	9,10,10	0.52	0
8	TRS	A	512	-	7,7,7	0.44	0	9,9,9	0.51	0
5	MPD	A	505	-	7,7,7	0.28	0	9,10,10	0.44	0
5	MPD	A	507	-	7,7,7	0.27	0	9,10,10	0.50	0
4	TRP	A	503	-	14,16,16	0.88	1 (7%)	16,22,22	1.07	2 (12%)
5	MPD	A	504	-	7,7,7	0.30	0	9,10,10	0.56	0
6	PGE	A	509	-	9,9,9	0.48	0	8,8,8	0.27	0
3	PEP	A	502	-	9,9,9	1.51	2 (22%)	11,13,13	1.77	3 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MPD	A	506	-	-	1/5/5/5	-
5	MPD	A	508	-	-	1/5/5/5	-
8	TRS	A	512	-	-	4/9/9/9	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MPD	A	505	-	-	1/5/5/5	-
5	MPD	A	507	-	-	1/5/5/5	-
4	TRP	A	503	-	-	0/7/8/8	0/2/2/2
5	MPD	A	504	-	-	1/5/5/5	-
6	PGE	A	509	-	-	4/7/7/7	-
3	PEP	A	502	-	-	4/9/9/9	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	PEP	O2-C2	2.53	1.46	1.39
3	A	502	PEP	C2-C1	-2.33	1.47	1.49
4	A	503	TRP	OXT-C	-2.05	1.23	1.30

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	PEP	O2'-C1-C2	3.44	119.77	113.91
3	A	502	PEP	O2-C2-C3	-2.51	119.95	124.79
4	A	503	TRP	OXT-C-O	-2.35	118.76	124.09
3	A	502	PEP	O1-C1-C2	-2.24	118.41	121.79
4	A	503	TRP	OXT-C-CA	2.15	120.69	113.38
7	A	510	PO4	O4-P-O2	2.02	114.45	107.97

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	PEP	O1-C1-C2-C3
3	A	502	PEP	O1-C1-C2-O2
3	A	502	PEP	O2'-C1-C2-C3
3	A	502	PEP	O2'-C1-C2-O2
5	A	504	MPD	C2-C3-C4-O4
5	A	506	MPD	C2-C3-C4-O4
8	A	512	TRS	N-C-C1-O1
8	A	512	TRS	N-C-C3-O3
6	A	509	PGE	O3-C5-C6-O4
8	A	512	TRS	C1-C-C3-O3
6	A	509	PGE	C1-C2-O2-C3
5	A	505	MPD	C2-C3-C4-C5
5	A	507	MPD	C2-C3-C4-C5

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
5	A	508	MPD	C2-C3-C4-C5
8	A	512	TRS	C3-C-C1-O1
6	A	509	PGE	O1-C1-C2-O2
6	A	509	PGE	C6-C5-O3-C4

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	506	MPD	1	0
5	A	508	MPD	2	0
5	A	504	MPD	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	447/472 (94%)	0.40	40 (8%) <b>9</b> <b>8</b>	72, 102, 158, 198	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	276	GLU	6.5
1	A	28	LEU	6.1
1	A	388	GLY	4.9
1	A	27	PRO	4.6
1	A	389	TYR	4.1
1	A	167	ALA	4.0
1	A	444	GLY	3.9
1	A	440	VAL	3.7
1	A	443	PRO	3.7
1	A	386	SER	3.5
1	A	93	GLN	3.3
1	A	26	PRO	3.2
1	A	255	CYS	3.1
1	A	94	GLY	3.0
1	A	33	GLN	3.0
1	A	387	ASN	2.9
1	A	420	ILE	2.9
1	A	134	MET	2.8
1	A	385	ALA	2.8
1	A	256	SER	2.7
1	A	390	LYS	2.7
1	A	170	GLU	2.7
1	A	428	THR	2.7
1	A	275	GLU	2.6
1	A	290	TRP	2.6
1	A	419	HIS	2.5
1	A	439	ASP	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	95	GLY	2.5
1	A	438	THR	2.5
1	A	135	ALA	2.4
1	A	153	GLY	2.4
1	A	278	ASN	2.4
1	A	166	GLU	2.3
1	A	92	LEU	2.3
1	A	152	ASN	2.2
1	A	277	GLY	2.2
1	A	435	GLU	2.2
1	A	383	PHE	2.1
1	A	417	GLY	2.1
1	A	274	ASP	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	MPD	A	505	8/8	0.76	0.47	148,155,157,159	0
5	MPD	A	506	8/8	0.83	0.35	143,146,147,148	0
7	PO4	A	511	5/5	0.84	0.24	68,70,74,76	1
5	MPD	A	508	8/8	0.85	0.36	137,143,144,146	0
7	PO4	A	510	5/5	0.86	0.24	62,69,69,78	1
5	MPD	A	507	8/8	0.87	0.52	118,128,137,139	0
5	MPD	A	504	8/8	0.89	0.27	132,140,145,147	0
8	TRS	A	512	8/8	0.91	0.22	90,103,110,112	0
6	PGE	A	509	10/10	0.94	0.28	98,103,106,107	0
3	PEP	A	502	10/10	0.94	0.25	103,116,124,125	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	TRP	A	503	15/15	0.95	0.17	90,93,97,101	0
2	MN	A	501	1/1	0.96	0.13	106,106,106,106	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.