



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 3, 2024 – 07:35 PM EST

PDB ID : 2HVG
Title : Crystal Structure of Adenylosuccinate Lyase from Plasmodium Vivax
Authors : Wernimont, A.K.; Dong, A.; Lew, J.; Wasney, G.A.; Vedadi, M.; Ren, H.; Alam, Z.; Qiu, W.; Kozieradzki, I.; Weigelt, J.; Sundstrom, M.; Edwards, A.M.; Arrowsmith, C.H.; Bochkarev, A.; Hui, R.; Hills, T.; Structural Genomics Consortium (SGC)
Deposited on : 2006-07-28
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	 2% 80% 11% • 7%
1	B	482	 5% 83% 9% • 7%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7783 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylosuccinate lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	447	3570	2288	607	658	12	5	0	0	0
1	B	447	3584	2293	614	660	12	5	0	0	0

There are 50 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	cloning artifact	UNP Q8WSJ9
A	-17	GLY	-	cloning artifact	UNP Q8WSJ9
A	-16	SER	-	cloning artifact	UNP Q8WSJ9
A	-15	SER	-	cloning artifact	UNP Q8WSJ9
A	-14	HIS	-	cloning artifact	UNP Q8WSJ9
A	-13	HIS	-	cloning artifact	UNP Q8WSJ9
A	-12	HIS	-	cloning artifact	UNP Q8WSJ9
A	-11	HIS	-	cloning artifact	UNP Q8WSJ9
A	-10	HIS	-	cloning artifact	UNP Q8WSJ9
A	-9	HIS	-	cloning artifact	UNP Q8WSJ9
A	-8	SER	-	cloning artifact	UNP Q8WSJ9
A	-7	SER	-	cloning artifact	UNP Q8WSJ9
A	-6	GLY	-	cloning artifact	UNP Q8WSJ9
A	-5	LEU	-	cloning artifact	UNP Q8WSJ9
A	-4	VAL	-	cloning artifact	UNP Q8WSJ9
A	-3	PRO	-	cloning artifact	UNP Q8WSJ9
A	-2	ARG	-	cloning artifact	UNP Q8WSJ9
A	-1	GLY	-	cloning artifact	UNP Q8WSJ9
A	0	SER	-	cloning artifact	UNP Q8WSJ9
A	148	MSE	MET	modified residue	UNP Q8WSJ9
A	181	MSE	MET	modified residue	UNP Q8WSJ9
A	296	MSE	MET	modified residue	UNP Q8WSJ9
A	364	LYS	ILE	SEE REMARK 999	UNP Q8WSJ9
A	391	MSE	MET	modified residue	UNP Q8WSJ9
A	417	MSE	MET	modified residue	UNP Q8WSJ9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	MSE	-	cloning artifact	UNP Q8WSJ9
B	-17	GLY	-	cloning artifact	UNP Q8WSJ9
B	-16	SER	-	cloning artifact	UNP Q8WSJ9
B	-15	SER	-	cloning artifact	UNP Q8WSJ9
B	-14	HIS	-	cloning artifact	UNP Q8WSJ9
B	-13	HIS	-	cloning artifact	UNP Q8WSJ9
B	-12	HIS	-	cloning artifact	UNP Q8WSJ9
B	-11	HIS	-	cloning artifact	UNP Q8WSJ9
B	-10	HIS	-	cloning artifact	UNP Q8WSJ9
B	-9	HIS	-	cloning artifact	UNP Q8WSJ9
B	-8	SER	-	cloning artifact	UNP Q8WSJ9
B	-7	SER	-	cloning artifact	UNP Q8WSJ9
B	-6	GLY	-	cloning artifact	UNP Q8WSJ9
B	-5	LEU	-	cloning artifact	UNP Q8WSJ9
B	-4	VAL	-	cloning artifact	UNP Q8WSJ9
B	-3	PRO	-	cloning artifact	UNP Q8WSJ9
B	-2	ARG	-	cloning artifact	UNP Q8WSJ9
B	-1	GLY	-	cloning artifact	UNP Q8WSJ9
B	0	SER	-	cloning artifact	UNP Q8WSJ9
B	148	MSE	MET	modified residue	UNP Q8WSJ9
B	181	MSE	MET	modified residue	UNP Q8WSJ9
B	296	MSE	MET	modified residue	UNP Q8WSJ9
B	364	LYS	ILE	SEE REMARK 999	UNP Q8WSJ9
B	391	MSE	MET	modified residue	UNP Q8WSJ9
B	417	MSE	MET	modified residue	UNP Q8WSJ9

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	290	Total O 290 290	0	0
3	B	274	Total O 274 274	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenylosuccinate lyase



- Molecule 1: Adenylosuccinate lyase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	120.42Å 177.60Å 68.61Å 90.00° 99.16° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 50.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.00-2.30) 99.3 (50.00-2.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.60 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.207 , 0.251 0.205 , 0.248	Depositor DCC
R_{free} test set	3170 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	31.4	Xtrriage
Anisotropy	0.089	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7783	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.60	1/3634 (0.0%)	0.65	0/4909
1	B	0.59	0/3648	0.64	1/4927 (0.0%)
All	All	0.59	1/7282 (0.0%)	0.65	1/9836 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	118	CYS	CB-SG	-5.37	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	337	ASP	CB-CG-OD1	5.14	122.93	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3570	0	3575	36	0
1	B	3584	0	3594	25	0
2	A	30	0	0	0	0
2	B	35	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	290	0	0	8	0
3	B	274	0	0	6	0
All	All	7783	0	7169	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:HIS:HD2	3:A:658:HOH:O	1.65	0.78
1:A:55:THR:HG22	1:A:106:ASP:OD2	1.90	0.72
1:A:92:GLU:HG3	1:A:114:VAL:HG12	1.71	0.71
1:B:57:HIS:ND1	1:B:104:ARG:NH2	2.40	0.69
1:B:410:LYS:HE2	3:B:641:HOH:O	1.93	0.67
1:A:320:LYS:HD3	3:B:794:HOH:O	1.96	0.64
1:A:333:ARG:O	1:A:333:ARG:HG3	1.98	0.63
1:B:85:ASN:N	3:B:686:HOH:O	2.31	0.62
1:B:459:SER:O	1:B:459:SER:OG	2.16	0.61
1:B:89:LYS:HE2	1:B:92:GLU:OE2	2.02	0.59
1:A:152:LYS:HD2	1:A:456:GLU:HB2	1.85	0.59
1:A:332:GLN:O	1:A:333:ARG:HB3	2.02	0.58
1:B:249:HIS:HD2	3:B:673:HOH:O	1.86	0.58
1:A:337:ASP:OD2	3:A:644:HOH:O	2.17	0.58
1:B:393:ARG:HH21	1:B:394:HIS:HE1	1.50	0.58
1:A:108:LEU:HA	1:A:111:LYS:HD3	1.85	0.57
1:A:100:LYS:HB2	1:A:111:LYS:HE3	1.88	0.56
1:A:217:ALA:HB2	1:B:389:ILE:HD13	1.87	0.55
1:B:85:ASN:O	1:B:89:LYS:HB2	2.07	0.55
1:A:369:ARG:O	1:A:373:GLU:HG2	2.07	0.54
1:B:46:GLU:HA	1:B:53:LYS:HE3	1.91	0.53
1:A:32:HIS:O	1:A:36:VAL:HG23	2.08	0.53
1:B:214:HIS:HB3	1:B:223:TRP:CE2	2.43	0.52
1:A:19:GLU:OE2	3:A:800:HOH:O	2.19	0.52
1:B:108:LEU:HA	1:B:111:LYS:HD3	1.92	0.52
1:A:12:ARG:HD3	1:A:13:TYR:CE1	2.45	0.50
1:B:112:GLU:HB2	3:B:644:HOH:O	2.11	0.50
1:B:12:ARG:HD3	1:B:13:TYR:CE1	2.47	0.50
1:B:419:GLU:OE1	1:B:423:THR:CG2	2.60	0.49
1:A:283:LYS:HE2	1:A:367:ILE:O	2.12	0.49
1:B:245:GLN:HB3	1:B:333:ARG:HG2	1.93	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:393:ARG:HH21	1:A:394:HIS:HE1	1.61	0.49
1:B:419:GLU:O	1:B:423:THR:HG23	2.12	0.48
1:A:214:HIS:HB3	1:A:223:TRP:CE2	2.49	0.47
1:A:55:THR:HG23	1:A:58:SER:H	1.79	0.47
1:A:279:ASN:O	1:A:280:ASN:HB2	2.15	0.46
1:B:150:LYS:HE3	1:B:363:ASN:OD1	2.16	0.45
1:A:422:LYS:HD3	1:A:437:LEU:HD21	1.99	0.45
1:A:422:LYS:HE3	3:A:636:HOH:O	2.16	0.45
1:A:413:ASP:HB2	3:A:730:HOH:O	2.17	0.45
1:A:316:ASN:O	1:A:320:LYS:HG3	2.17	0.44
1:B:148:MSE:HG3	1:B:188:ILE:HG22	1.99	0.44
1:A:196:ARG:HE	1:A:196:ARG:HB3	1.40	0.44
1:A:200:VAL:HG13	1:A:251:TYR:CE1	2.52	0.44
1:A:89:LYS:HE3	1:A:93:TYR:CE1	2.53	0.44
1:B:394:HIS:O	1:B:395:ASN:HB2	2.18	0.44
1:B:418:GLN:O	1:B:422:LYS:HD3	2.18	0.44
1:A:43:PHE:HA	1:A:46:GLU:HG2	2.00	0.43
1:A:261:ARG:HG2	3:B:758:HOH:O	2.17	0.42
1:B:10:ASP:HA	1:B:14:LYS:HE2	2.01	0.42
1:B:211:PHE:HB3	1:B:214:HIS:HB2	2.02	0.42
1:A:33:ARG:NH1	3:A:733:HOH:O	2.52	0.42
1:A:257:ASP:OD2	1:B:261:ARG:NH1	2.46	0.42
1:A:373:GLU:HG3	3:A:787:HOH:O	2.20	0.42
1:A:164:LEU:HD22	1:A:375:GLU:HG2	2.02	0.41
1:A:192:VAL:O	1:A:196:ARG:HG2	2.21	0.41
1:B:296:MSE:HA	1:B:297:PRO:HD3	1.90	0.40
1:A:393:ARG:NH2	1:A:394:HIS:HE1	2.19	0.40
1:A:300:VAL:HG13	3:A:729:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/482 (92%)	425 (96%)	15 (3%)	1 (0%)	44	55
1	B	441/482 (92%)	426 (97%)	14 (3%)	1 (0%)	44	55
All	All	882/964 (92%)	851 (96%)	29 (3%)	2 (0%)	44	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	333	ARG
1	B	333	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	392/425 (92%)	377 (96%)	15 (4%)	28	42
1	B	395/425 (93%)	378 (96%)	17 (4%)	25	36
All	All	787/850 (93%)	755 (96%)	32 (4%)	26	39

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	ARG
1	A	58	SER
1	A	59	VAL
1	A	98	LYS
1	A	106	ASP
1	A	125	ASN
1	A	196	ARG
1	A	203	LYS
1	A	208	VAL
1	A	229	LEU
1	A	324	SER
1	A	341	LEU
1	A	422	LYS
1	A	443	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	458	LEU
1	B	12	ARG
1	B	59	VAL
1	B	78	LYS
1	B	87	ASP
1	B	89	LYS
1	B	106	ASP
1	B	111	LYS
1	B	125	ASN
1	B	203	LYS
1	B	208	VAL
1	B	229	LEU
1	B	333	ARG
1	B	339	THR
1	B	422	LYS
1	B	423	THR
1	B	458	LEU
1	B	459	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	307	ASN
1	A	394	HIS
1	B	27	HIS
1	B	249	HIS
1	B	394	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	602	-	4,4,4	0.25	0	6,6,6	0.24	0
2	SO4	A	601	-	4,4,4	0.31	0	6,6,6	0.45	0
2	SO4	B	610	-	4,4,4	0.24	0	6,6,6	0.29	0
2	SO4	A	606	-	4,4,4	0.23	0	6,6,6	0.08	0
2	SO4	B	613	-	4,4,4	0.34	0	6,6,6	0.30	0
2	SO4	A	603	-	4,4,4	0.23	0	6,6,6	0.29	0
2	SO4	B	608	-	4,4,4	0.24	0	6,6,6	0.33	0
2	SO4	B	609	-	4,4,4	0.23	0	6,6,6	0.23	0
2	SO4	A	604	-	4,4,4	0.24	0	6,6,6	0.20	0
2	SO4	B	611	-	4,4,4	0.24	0	6,6,6	0.26	0
2	SO4	A	605	-	4,4,4	0.23	0	6,6,6	0.14	0
2	SO4	B	612	-	4,4,4	0.21	0	6,6,6	0.38	0
2	SO4	B	607	-	4,4,4	0.25	0	6,6,6	0.32	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/482 (91%)	0.33	12 (2%) 56 57	17, 29, 48, 64	0
1	B	442/482 (91%)	0.42	23 (5%) 34 35	17, 29, 47, 65	0
All	All	884/964 (91%)	0.38	35 (3%) 43 44	17, 29, 47, 65	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	85	ASN	3.7
1	A	86	HIS	3.4
1	B	106	ASP	3.2
1	B	403	LEU	3.1
1	B	419	GLU	3.1
1	A	405	GLN	3.0
1	B	86	HIS	3.0
1	B	68	ASN	3.0
1	A	101	ASN	2.9
1	B	103	LYS	2.9
1	B	101	ASN	2.8
1	B	80	ILE	2.7
1	A	93	TYR	2.6
1	B	16	ALA	2.5
1	A	220	ASP	2.5
1	A	55	THR	2.5
1	A	403	LEU	2.4
1	B	434	ASP	2.4
1	B	459	SER	2.3
1	B	109	LYS	2.3
1	B	73	ASP	2.3
1	B	69	ILE	2.3
1	B	405	GLN	2.3
1	A	68	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	431	ASP	2.2
1	B	438	GLU	2.1
1	B	93	TYR	2.1
1	B	220	ASP	2.1
1	A	81	GLU	2.1
1	B	374	GLU	2.1
1	A	438	GLU	2.0
1	B	54	VAL	2.0
1	B	387	ILE	2.0
1	A	105	GLU	2.0
1	B	61	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	613	5/5	0.68	0.16	76,76,77,78	0
2	SO4	B	609	5/5	0.79	0.16	80,80,81,81	0
2	SO4	B	608	5/5	0.83	0.18	67,67,68,68	0
2	SO4	A	602	5/5	0.83	0.17	62,63,64,64	0
2	SO4	A	603	5/5	0.83	0.18	74,74,74,74	0
2	SO4	B	607	5/5	0.85	0.22	69,70,70,70	0
2	SO4	B	610	5/5	0.85	0.19	69,69,70,70	0
2	SO4	A	604	5/5	0.85	0.12	74,74,74,75	0
2	SO4	B	611	5/5	0.87	0.13	65,65,65,66	0
2	SO4	B	612	5/5	0.89	0.21	61,61,63,63	0
2	SO4	A	605	5/5	0.90	0.14	67,67,67,68	0
2	SO4	A	601	5/5	0.91	0.15	50,51,54,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SO4	A	606	5/5	0.92	0.14	55,56,56,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.