



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 07:37 AM BST

PDB ID : 4HYX  
Title : Crystal Structure Analysis of the Bacteriorhodopsin in Facial Amphiphile-4 DMPC Bicelle  
Authors : Lee, S.; Stout, C.D.; Zhang, Q.  
Deposited on : 2012-11-14  
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	 3% 74% 10% • 14%
1	B	262	 3% 75% 11% • 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
2	RET	A	301	-	-	X	-
2	RET	B	301	-	-	X	-
7	GOL	B	309	-	-	X	-

## 2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 3754 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacteriorhodopsin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	226	Total 1749	C 1176	N 267	O 297	S 9	0	0	0
1	B	231	Total 1788	C 1200	N 272	O 307	S 9	0	0	0

- Molecule 2 is RETINAL (three-letter code: RET) (formula: C<sub>20</sub>H<sub>28</sub>O).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C		
2	A	1	Total 20	C 20	0	0
2	B	1	Total 20	C 20	0	0

- Molecule 3 is N-OCTANE (three-letter code: OCT) (formula: C<sub>8</sub>H<sub>18</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 8 8	0	0
3	A	1	Total C 8 8	0	0
3	B	1	Total C 8 8	0	0
3	B	1	Total C 8 8	0	0

- Molecule 4 is DECANE (three-letter code: D10) (formula:  $C_{10}H_{22}$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C 10 10	0	0
4	A	1	Total C 10 10	0	0
4	A	1	Total C 10 10	0	0
4	A	1	Total C 10 10	0	0
4	A	1	Total C 10 10	0	0
4	B	1	Total C 10 10	0	0
4	B	1	Total C 10 10	0	0
4	B	1	Total C 10 10	0	0

- Molecule 5 is HEXANE (three-letter code: HEX) (formula: C<sub>6</sub>H<sub>14</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C 6 6	0	0

- Molecule 6 is TETRADECANE (three-letter code: C14) (formula: C<sub>14</sub>H<sub>30</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C 14 14	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C O 6 3 3	0	0

- Molecule 8 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
8	A	22	Total 22	O 22	0	0
8	B	17	Total 17	O 17	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bacteriorhodopsin



- Molecule 1: Bacteriorhodopsin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.89Å 112.92Å 56.29Å 90.00° 113.83° 90.00°	Depositor
Resolution (Å)	46.85 – 1.99 46.85 – 1.99	Depositor EDS
% Data completeness (in resolution range)	95.7 (46.85-1.99) 95.5 (46.85-1.99)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	14.97 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.179 , 0.214 0.185 , 0.215	Depositor DCC
$R_{free}$ test set	1759 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.8	Xtrriage
Anisotropy	0.404	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 73.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.098 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3754	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, D10, HEX, RET, OCT, C14

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.14	2/1797 (0.1%)	0.98	4/2454 (0.2%)
1	B	1.08	0/1837	0.98	3/2509 (0.1%)
All	All	1.11	2/3634 (0.1%)	0.98	7/4963 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	43	TYR	CE1-CZ	-5.73	1.31	1.38
1	A	26	TYR	CB-CG	-5.58	1.43	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	209	MET	CG-SD-CE	-8.04	87.33	100.20
1	A	209	MET	CG-SD-CE	-8.01	87.38	100.20
1	B	60	MET	CG-SD-CE	6.18	110.09	100.20
1	B	85	ASP	CB-CG-OD1	5.79	123.51	118.30
1	A	85	ASP	CB-CG-OD1	5.76	123.48	118.30
1	A	95	LEU	CB-CG-CD2	5.54	120.42	111.00
1	A	28	LEU	CA-CB-CG	5.25	127.38	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1749	0	1808	26	0
1	B	1788	0	1840	38	0
2	A	20	0	27	12	0
2	B	20	0	27	10	0
3	A	16	0	36	2	0
3	B	16	0	36	0	0
4	A	50	0	110	1	0
4	B	30	0	66	1	0
5	B	6	0	14	1	0
6	B	14	0	30	1	0
7	B	6	0	8	8	0
8	A	22	0	0	1	0
8	B	17	0	0	5	0
All	All	3754	0	4002	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:216:LYS:HZ2	2:B:301:RET:C15	1.39	1.30
1:A:216:LYS:HZ2	2:A:301:RET:C15	1.58	1.10
1:A:216:LYS:HZ1	2:A:301:RET:C15	1.75	0.92
1:B:216:LYS:HZ1	2:B:301:RET:C15	1.69	0.92
1:B:117:ILE:CG2	8:B:414:HOH:O	2.21	0.88
1:B:216:LYS:HZ1	2:B:301:RET:C14	1.90	0.83
2:A:301:RET:H171	2:A:301:RET:H8	1.58	0.83
1:B:216:LYS:NZ	2:B:301:RET:C14	2.43	0.81
1:B:201:LEU:HD13	7:B:309:GOL:H11	1.63	0.80
1:A:216:LYS:HZ1	2:A:301:RET:C14	1.96	0.77
1:A:15:LEU:HD22	1:A:209:MET:CE	2.14	0.77
1:A:216:LYS:NZ	2:A:301:RET:C14	2.50	0.75
1:B:128:THR:HG23	4:B:302:D10:H81	1.67	0.75
1:A:163:MET:HG3	8:A:403:HOH:O	1.89	0.73
1:B:216:LYS:CE	2:B:301:RET:C15	2.67	0.72
1:B:15:LEU:HD22	1:B:209:MET:CE	2.19	0.72
1:B:79:TYR:HD2	7:B:309:GOL:H2	1.55	0.71
1:B:121:THR:HG21	8:B:414:HOH:O	1.91	0.69
1:B:117:ILE:HG23	8:B:414:HOH:O	1.89	0.67

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:LEU:HD13	1:A:58:LEU:C	2.15	0.67
1:A:15:LEU:HD22	1:A:209:MET:HE1	1.79	0.64
1:A:216:LYS:CE	2:A:301:RET:C15	2.75	0.64
1:B:201:LEU:CD1	7:B:309:GOL:H11	2.26	0.63
1:A:51:ALA:O	1:A:55:THR:HG23	1.99	0.62
1:B:51:ALA:O	1:B:55:THR:HG23	2.00	0.62
1:B:144:ALA:HB2	6:B:305:C14:H143	1.82	0.61
1:B:15:LEU:HD22	1:B:209:MET:HE1	1.84	0.60
1:B:202:ASN:HB2	8:B:417:HOH:O	2.02	0.60
1:A:15:LEU:HD21	1:A:19:LEU:HD11	1.84	0.59
1:B:58:LEU:C	1:B:58:LEU:HD13	2.22	0.59
1:B:15:LEU:HD21	1:B:19:LEU:HD11	1.85	0.57
1:B:163:MET:HG2	1:B:167:VAL:HG11	1.87	0.57
2:B:301:RET:H171	2:B:301:RET:H8	1.90	0.53
1:A:163:MET:HG2	1:A:167:VAL:HG11	1.91	0.52
1:A:15:LEU:HD22	1:A:209:MET:SD	2.50	0.52
1:B:140:ILE:HG22	8:B:414:HOH:O	2.11	0.51
1:A:86:TRP:CD1	2:A:301:RET:H14	2.47	0.49
1:B:201:LEU:HD13	7:B:309:GOL:C1	2.38	0.49
1:B:15:LEU:HD22	1:B:209:MET:SD	2.54	0.48
1:B:65:GLY:HA3	1:B:81:ALA:HB2	1.95	0.48
1:A:202:ASN:H	1:A:202:ASN:HD22	1.61	0.48
2:B:301:RET:H181	2:B:301:RET:H7	1.64	0.48
1:B:65:GLY:HA2	1:B:80:TRP:CE2	2.49	0.48
1:B:201:LEU:CD1	7:B:309:GOL:C1	2.91	0.47
1:B:193:SER:OG	7:B:309:GOL:H11	2.15	0.46
1:A:142:THR:HA	2:A:301:RET:H182	1.97	0.46
1:B:202:ASN:H	1:B:202:ASN:HD22	1.61	0.46
1:B:86:TRP:CD1	2:B:301:RET:H14	2.51	0.46
1:A:176:ASN:HD22	4:A:304:D10:H103	1.81	0.46
1:A:56:MET:HG3	1:A:85:ASP:HB2	1.98	0.46
1:B:216:LYS:HZ1	2:B:301:RET:H14	1.77	0.45
1:B:77:PRO:O	7:B:309:GOL:O1	2.28	0.45
1:A:65:GLY:HA3	1:A:81:ALA:HB2	1.98	0.45
1:B:15:LEU:CD2	1:B:19:LEU:CD1	2.95	0.45
1:A:15:LEU:CD2	1:A:19:LEU:HD11	2.48	0.44
1:A:15:LEU:CD2	1:A:19:LEU:CD1	2.96	0.44
1:A:46:THR:O	1:A:50:PRO:HD2	2.17	0.43
2:B:301:RET:H161	2:B:301:RET:H8	2.00	0.43
2:A:301:RET:H171	2:A:301:RET:C8	2.35	0.43
1:A:186:PRO:HB3	2:A:301:RET:H183	2.00	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:110:ALA:HB2	5:B:304:HEX:H22	2.01	0.42
1:A:216:LYS:HZ1	2:A:301:RET:H14	1.81	0.42
1:B:28:LEU:O	1:B:32:MET:HG2	2.20	0.42
1:B:202:ASN:N	1:B:202:ASN:HD22	2.18	0.42
3:A:308:OCT:H62	3:A:308:OCT:H32	1.85	0.41
1:B:15:LEU:HD23	1:B:15:LEU:C	2.40	0.41
1:B:82:ARG:HE	7:B:309:GOL:H31	1.86	0.41
1:A:90:THR:HG22	2:A:301:RET:H202	2.04	0.40
1:B:15:LEU:HD23	1:B:15:LEU:O	2.22	0.40
1:A:183:SER:HA	3:A:308:OCT:H81	2.03	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/262 (86%)	222 (99%)	2 (1%)	0	100	100
1	B	227/262 (87%)	225 (99%)	2 (1%)	0	100	100
All	All	451/524 (86%)	447 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/206 (88%)	172 (95%)	9 (5%)	24	20
1	B	186/206 (90%)	178 (96%)	8 (4%)	29	26
All	All	367/412 (89%)	350 (95%)	17 (5%)	27	23

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LEU
1	A	25	LEU
1	A	28	LEU
1	A	95	LEU
1	A	131	TYR
1	A	161	GLU
1	A	169	SER
1	A	186	PRO
1	A	202	ASN
1	B	7	ARG
1	B	22	LEU
1	B	25	LEU
1	B	28	LEU
1	B	95	LEU
1	B	131	TYR
1	B	163	MET
1	B	202	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	GLN
1	A	202	ASN
1	B	75	GLN
1	B	202	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	D10	B	308	-	9,9,9	0.18	0	8,8,8	0.44	0
4	D10	B	303	-	9,9,9	0.33	0	8,8,8	0.26	0
4	D10	B	302	-	9,9,9	0.26	0	8,8,8	0.53	0
3	OCT	B	307	-	7,7,7	0.25	0	6,6,6	0.42	0
3	OCT	A	302	-	7,7,7	0.30	0	6,6,6	0.25	0
3	OCT	B	306	-	7,7,7	0.24	0	6,6,6	0.31	0
4	D10	A	304	-	9,9,9	0.31	0	8,8,8	0.44	0
6	C14	B	305	-	13,13,13	0.25	0	12,12,12	0.57	0
3	OCT	A	308	-	7,7,7	0.21	0	6,6,6	0.66	0
7	GOL	B	309	-	5,5,5	0.55	0	5,5,5	0.52	0
4	D10	A	307	-	9,9,9	0.36	0	8,8,8	0.26	0
4	D10	A	303	-	9,9,9	0.22	0	8,8,8	0.53	0
2	RET	B	301	1	20,20,21	1.50	6 (30%)	27,27,28	2.18	11 (40%)
2	RET	A	301	1	20,20,21	1.47	4 (20%)	27,27,28	2.30	10 (37%)
4	D10	A	306	-	9,9,9	0.28	0	8,8,8	0.41	0
5	HEX	B	304	-	5,5,5	0.23	0	4,4,4	0.33	0
4	D10	A	305	-	9,9,9	0.25	0	8,8,8	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	D10	B	308	-	-	1/7/7/7	-
4	D10	B	303	-	-	4/7/7/7	-

*Continued on next page...*



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	D10	B	302	-	-	3/7/7/7	-
3	OCT	B	307	-	-	3/5/5/5	-
3	OCT	A	302	-	-	4/5/5/5	-
3	OCT	B	306	-	-	2/5/5/5	-
4	D10	A	304	-	-	2/7/7/7	-
6	C14	B	305	-	-	6/11/11/11	-
3	OCT	A	308	-	-	2/5/5/5	-
7	GOL	B	309	-	-	4/4/4/4	-
4	D10	A	307	-	-	4/7/7/7	-
4	D10	A	303	-	-	4/7/7/7	-
2	RET	B	301	1	-	0/13/30/31	0/1/1/1
2	RET	A	301	1	-	0/13/30/31	0/1/1/1
4	D10	A	306	-	-	4/7/7/7	-
5	HEX	B	304	-	-	2/3/3/3	-
4	D10	A	305	-	-	2/7/7/7	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	RET	C14-C13	-3.23	1.31	1.33
2	A	301	RET	C5-C6	2.74	1.39	1.34
2	B	301	RET	C20-C13	-2.73	1.45	1.50
2	A	301	RET	C15-C14	-2.51	1.40	1.49
2	B	301	RET	C18-C5	-2.46	1.46	1.50
2	B	301	RET	C2-C1	-2.45	1.48	1.54
2	B	301	RET	C19-C9	-2.33	1.46	1.50
2	B	301	RET	C4-C5	-2.10	1.46	1.51
2	A	301	RET	C1-C6	2.09	1.56	1.53
2	B	301	RET	C8-C9	-2.02	1.41	1.45

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	RET	C20-C13-C12	7.48	129.87	118.08
2	B	301	RET	C20-C13-C12	5.97	127.48	118.08
2	B	301	RET	C8-C9-C10	-4.03	112.76	118.94
2	B	301	RET	C2-C1-C6	3.73	116.23	110.48
2	A	301	RET	C8-C9-C10	-3.25	113.95	118.94
2	B	301	RET	C10-C11-C12	-2.93	114.08	123.22

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	RET	C19-C9-C8	2.89	122.63	118.08
2	B	301	RET	C19-C9-C8	2.89	122.62	118.08
2	A	301	RET	C15-C14-C13	-2.73	111.90	127.65
2	A	301	RET	C18-C5-C4	-2.59	108.63	113.62
2	B	301	RET	C20-C13-C14	-2.51	116.30	123.71
2	B	301	RET	C1-C6-C7	2.49	122.83	115.78
2	A	301	RET	C10-C11-C12	-2.41	115.71	123.22
2	A	301	RET	C3-C2-C1	2.37	123.08	114.60
2	B	301	RET	C7-C6-C5	-2.35	115.78	121.46
2	A	301	RET	C20-C13-C14	-2.34	116.78	123.71
2	B	301	RET	C3-C4-C5	2.28	118.15	114.08
2	B	301	RET	C15-C14-C13	-2.23	114.77	127.65
2	B	301	RET	C2-C3-C4	2.08	116.03	111.38
2	A	301	RET	C1-C6-C7	2.04	121.56	115.78
2	A	301	RET	C7-C6-C5	-2.03	116.53	121.46

There are no chirality outliers.

All (47) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	309	GOL	O1-C1-C2-C3
4	B	302	D10	C4-C5-C6-C7
3	A	308	OCT	C3-C4-C5-C6
4	A	307	D10	C5-C6-C7-C8
3	B	306	OCT	C4-C5-C6-C7
4	A	303	D10	C6-C7-C8-C9
3	A	302	OCT	C3-C4-C5-C6
4	A	307	D10	C6-C7-C8-C9
7	B	309	GOL	C1-C2-C3-O3
4	A	304	D10	C4-C5-C6-C7
4	B	302	D10	C2-C3-C4-C5
4	A	305	D10	C6-C7-C8-C9
3	B	306	OCT	C2-C3-C4-C5
6	B	305	C14	C07-C08-C09-C10
6	B	305	C14	C09-C10-C11-C12
7	B	309	GOL	O1-C1-C2-O2
4	B	303	D10	C3-C4-C5-C6
4	A	307	D10	C4-C5-C6-C7
4	A	304	D10	C6-C7-C8-C9
4	A	303	D10	C5-C6-C7-C8
6	B	305	C14	C04-C05-C06-C07
3	B	307	OCT	C1-C2-C3-C4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
6	B	305	C14	C01-C02-C03-C04
4	A	306	D10	C1-C2-C3-C4
3	B	307	OCT	C4-C5-C6-C7
4	B	302	D10	C7-C8-C9-C10
4	A	306	D10	C2-C3-C4-C5
4	B	308	D10	C5-C6-C7-C8
3	A	302	OCT	C4-C5-C6-C7
4	B	303	D10	C6-C7-C8-C9
4	B	303	D10	C4-C5-C6-C7
4	A	306	D10	C6-C7-C8-C9
4	A	307	D10	C3-C4-C5-C6
5	B	304	HEX	C1-C2-C3-C4
3	A	302	OCT	C2-C3-C4-C5
6	B	305	C14	C02-C03-C04-C05
3	A	302	OCT	C1-C2-C3-C4
3	B	307	OCT	C2-C3-C4-C5
4	A	305	D10	C2-C3-C4-C5
6	B	305	C14	C03-C04-C05-C06
7	B	309	GOL	O2-C2-C3-O3
4	A	303	D10	C1-C2-C3-C4
3	A	308	OCT	C4-C5-C6-C7
5	B	304	HEX	C2-C3-C4-C5
4	B	303	D10	C2-C3-C4-C5
4	A	303	D10	C4-C5-C6-C7
4	A	306	D10	C5-C6-C7-C8


There are no ring outliers.

8 monomers are involved in 36 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	302	D10	1	0
4	A	304	D10	1	0
6	B	305	C14	1	0
3	A	308	OCT	2	0
7	B	309	GOL	8	0
2	B	301	RET	10	0
2	A	301	RET	12	0
5	B	304	HEX	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	226/262 (86%)	0.24	9 (3%) 38 37	20, 28, 52, 70	0
1	B	231/262 (88%)	0.20	8 (3%) 44 43	21, 29, 56, 80	0
All	All	457/524 (87%)	0.22	17 (3%) 41 41	20, 28, 53, 80	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	231	GLY	7.0
1	B	239	SER	5.7
1	A	6	GLY	5.0
1	A	73	GLY	4.7
1	B	6	GLY	4.6
1	A	230	PHE	4.1
1	B	161	GLU	3.8
1	A	38	ASP	3.7
1	A	227	ARG	3.5
1	B	35	SER	2.6
1	B	102	ASP	2.5
1	B	38	ASP	2.4
1	A	68	MET	2.4
1	B	227	ARG	2.3
1	A	225	ARG	2.3
1	A	70	PRO	2.1
1	B	37	PRO	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	D10	B	303	10/10	0.47	0.26	56,69,74,75	0
3	OCT	B	307	8/8	0.57	0.32	57,60,66,67	0
3	OCT	B	306	8/8	0.60	0.25	60,66,68,68	0
4	D10	A	307	10/10	0.61	0.24	54,61,64,65	0
6	C14	B	305	14/14	0.63	0.20	68,76,78,80	0
4	D10	A	305	10/10	0.66	0.17	65,72,77,78	0
4	D10	A	306	10/10	0.70	0.30	61,67,73,74	0
4	D10	B	302	10/10	0.75	0.24	54,66,69,72	0
4	D10	B	308	10/10	0.79	0.25	39,46,61,63	0
4	D10	A	304	10/10	0.79	0.20	55,62,64,68	0
3	OCT	A	302	8/8	0.83	0.20	66,69,70,72	0
7	GOL	B	309	6/6	0.85	0.59	59,64,64,70	0
5	HEX	B	304	6/6	0.88	0.26	54,60,65,66	0
3	OCT	A	308	8/8	0.89	0.22	41,48,54,57	0
4	D10	A	303	10/10	0.89	0.20	42,60,65,68	0
2	RET	A	301	20/21	0.92	0.13	22,29,37,39	0
2	RET	B	301	20/21	0.96	0.14	20,26,31,34	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around RET A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around RET B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.