



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 11, 2021 – 03:48 PM EDT

PDB ID : 2HZM
Title : Structure of the Mediator head subcomplex Med18/20
Authors : Lariviere, L.; Geiger, S.; Hoepfner, S.; Rother, S.; Straesser, K.; Cramer, P.
Deposited on : 2006-08-09
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	 2% 74% 19% 8%
1	C	212	 8% 65% 23% 8%
1	E	212	 23% 55% 34% 7%
1	G	212	 28% 56% 34% 8%
2	B	317	 4% 50% 18% 28%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	317	
2	F	317	
2	H	317	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA polymerase II mediator complex subunit 20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	205	1574	994	265	310	5	0	0	0
1	C	195	1504	952	253	294	5	0	0	0
1	E	197	1520	961	257	297	5	0	0	0
1	G	196	1511	957	254	295	5	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ALA	-	cloning artifact	UNP P34162
A	0	SER	-	cloning artifact	UNP P34162
C	-1	ALA	-	cloning artifact	UNP P34162
C	0	SER	-	cloning artifact	UNP P34162
E	-1	ALA	-	cloning artifact	UNP P34162
E	0	SER	-	cloning artifact	UNP P34162
G	-1	ALA	-	cloning artifact	UNP P34162
G	0	SER	-	cloning artifact	UNP P34162

- Molecule 2 is a protein called RNA polymerase II mediator complex subunit 18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	227	1785	1145	291	342	7	0	0	0
2	D	233	1820	1165	299	347	9	0	0	0
2	F	238	1869	1199	302	358	10	0	0	0
2	H	226	1770	1137	287	337	9	0	0	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	274	VAL	ALA	engineered mutation	UNP P32585
B	308	ALA	-	cloning artifact	UNP P32585
B	309	ALA	-	cloning artifact	UNP P32585
B	310	ALA	-	cloning artifact	UNP P32585
B	311	LEU	-	cloning artifact	UNP P32585
B	312	GLU	-	cloning artifact	UNP P32585
B	313	HIS	-	expression tag	UNP P32585
B	314	HIS	-	expression tag	UNP P32585
B	315	HIS	-	expression tag	UNP P32585
B	316	HIS	-	expression tag	UNP P32585
B	317	HIS	-	expression tag	UNP P32585
B	318	HIS	-	expression tag	UNP P32585
D	274	VAL	ALA	engineered mutation	UNP P32585
D	308	ALA	-	cloning artifact	UNP P32585
D	309	ALA	-	cloning artifact	UNP P32585
D	310	ALA	-	cloning artifact	UNP P32585
D	311	LEU	-	cloning artifact	UNP P32585
D	312	GLU	-	cloning artifact	UNP P32585
D	313	HIS	-	expression tag	UNP P32585
D	314	HIS	-	expression tag	UNP P32585
D	315	HIS	-	expression tag	UNP P32585
D	316	HIS	-	expression tag	UNP P32585
D	317	HIS	-	expression tag	UNP P32585
D	318	HIS	-	expression tag	UNP P32585
F	274	VAL	ALA	engineered mutation	UNP P32585
F	308	ALA	-	cloning artifact	UNP P32585
F	309	ALA	-	cloning artifact	UNP P32585
F	310	ALA	-	cloning artifact	UNP P32585
F	311	LEU	-	cloning artifact	UNP P32585
F	312	GLU	-	cloning artifact	UNP P32585
F	313	HIS	-	expression tag	UNP P32585
F	314	HIS	-	expression tag	UNP P32585
F	315	HIS	-	expression tag	UNP P32585
F	316	HIS	-	expression tag	UNP P32585
F	317	HIS	-	expression tag	UNP P32585
F	318	HIS	-	expression tag	UNP P32585
H	274	VAL	ALA	engineered mutation	UNP P32585
H	308	ALA	-	cloning artifact	UNP P32585
H	309	ALA	-	cloning artifact	UNP P32585
H	310	ALA	-	cloning artifact	UNP P32585
H	311	LEU	-	cloning artifact	UNP P32585
H	312	GLU	-	cloning artifact	UNP P32585

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	313	HIS	-	expression tag	UNP P32585
H	314	HIS	-	expression tag	UNP P32585
H	315	HIS	-	expression tag	UNP P32585
H	316	HIS	-	expression tag	UNP P32585
H	317	HIS	-	expression tag	UNP P32585
H	318	HIS	-	expression tag	UNP P32585

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	76	Total O 76 76	0	0
4	B	40	Total O 40 40	0	0
4	C	16	Total O 16 16	0	0
4	D	41	Total O 41 41	0	0
4	E	6	Total O 6 6	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	F	40	Total 40	O 40	0	0
4	G	2	Total 2	O 2	0	0
4	H	8	Total 8	O 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA polymerase II mediator complex subunit 20




- Molecule 1: RNA polymerase II mediator complex subunit 20




- Molecule 1: RNA polymerase II mediator complex subunit 20





• Molecule 2: RNA polymerase II mediator complex subunit 18



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.90Å 129.35Å 241.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40 19.98 – 2.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.40) 98.7 (19.98-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.24 (at 2.41Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.228 , 0.264 0.228 , 0.263	Depositor DCC
R_{free} test set	4276 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	44.8	Xtrriage
Anisotropy	0.469	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13587	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1598	0.72	0/2162
1	C	0.36	0/1526	0.60	0/2064
1	E	0.31	0/1542	0.54	0/2083
1	G	0.34	0/1533	0.58	0/2072
2	B	0.48	0/1816	0.75	3/2453 (0.1%)
2	D	0.43	0/1854	0.68	1/2508 (0.0%)
2	F	0.44	0/1903	0.67	0/2573
2	H	0.36	0/1801	0.63	2/2433 (0.1%)
All	All	0.42	0/13573	0.65	6/18348 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	84	PRO	N-CA-C	7.68	132.08	112.10
2	H	82	ASP	N-CA-C	6.66	128.98	111.00
2	H	241	ALA	N-CA-C	-6.21	94.22	111.00
2	B	85	LEU	CA-CB-CG	5.80	128.65	115.30
2	D	79	THR	N-CA-C	-5.28	96.73	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1574	0	1588	36	0
1	C	1504	0	1517	45	0
1	E	1520	0	1534	77	0
1	G	1511	0	1526	81	0
2	B	1785	0	1799	68	0
2	D	1820	0	1837	68	0
2	F	1869	0	1885	42	0
2	H	1770	0	1791	74	0
3	A	5	0	0	0	0
4	A	76	0	0	0	0
4	B	40	0	0	1	0
4	C	16	0	0	0	0
4	D	41	0	0	0	0
4	E	6	0	0	0	0
4	F	40	0	0	1	0
4	G	2	0	0	0	0
4	H	8	0	0	0	0
All	All	13587	0	13477	465	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 465 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:83:LYS:HB3	2:B:84:PRO:HD2	1.20	1.07
1:A:114:ASN:HD22	2:B:49:VAL:HG11	1.24	1.02
1:E:40:ARG:HH11	1:E:118:GLN:HE22	1.10	0.95
2:B:83:LYS:CB	2:B:84:PRO:HD2	1.96	0.95
2:B:4:GLN:HE21	2:B:258:ASN:HD21	1.16	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	201/212 (95%)	193 (96%)	5 (2%)	3 (2%)	10	14
1	C	189/212 (89%)	172 (91%)	14 (7%)	3 (2%)	9	13
1	E	191/212 (90%)	163 (85%)	23 (12%)	5 (3%)	5	5
1	G	190/212 (90%)	161 (85%)	23 (12%)	6 (3%)	4	3
2	B	217/317 (68%)	209 (96%)	5 (2%)	3 (1%)	11	15
2	D	227/317 (72%)	209 (92%)	14 (6%)	4 (2%)	8	10
2	F	230/317 (73%)	221 (96%)	8 (4%)	1 (0%)	34	48
2	H	216/317 (68%)	195 (90%)	15 (7%)	6 (3%)	5	4
All	All	1661/2116 (78%)	1523 (92%)	107 (6%)	31 (2%)	8	10

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
1	A	193	THR
1	C	161	GLY
1	C	187	ASP
2	D	79	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/179 (97%)	163 (94%)	11 (6%)	18	28
1	C	166/179 (93%)	158 (95%)	8 (5%)	25	41
1	E	167/179 (93%)	160 (96%)	7 (4%)	30	47
1	G	166/179 (93%)	159 (96%)	7 (4%)	30	47
2	B	204/288 (71%)	192 (94%)	12 (6%)	19	32
2	D	207/288 (72%)	194 (94%)	13 (6%)	18	28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	F	214/288 (74%)	207 (97%)	7 (3%)	38 57
2	H	203/288 (70%)	196 (97%)	7 (3%)	37 56
All	All	1501/1868 (80%)	1429 (95%)	72 (5%)	25 41

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	261	ARG
2	H	261	ARG
1	G	11	ARG
2	H	66	LEU
1	C	71	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	118	GLN
1	G	68	GLN
1	E	120	GLN
2	F	237	GLN
1	G	120	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	211	-	4,4,4	1.62	0	6,6,6	0.45	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	205/212 (96%)	-0.14	5 (2%) 59 57	20, 31, 57, 91	0
1	C	195/212 (91%)	0.43	18 (9%) 9 8	29, 58, 84, 91	0
1	E	197/212 (92%)	1.33	49 (24%) 0 0	30, 88, 100, 100	0
1	G	196/212 (92%)	1.47	59 (30%) 0 0	36, 91, 100, 100	0
2	B	227/317 (71%)	0.05	13 (5%) 23 22	20, 39, 71, 94	0
2	D	233/317 (73%)	0.28	18 (7%) 13 12	24, 45, 82, 91	0
2	F	238/317 (75%)	0.05	14 (5%) 22 21	21, 43, 77, 95	0
2	H	226/317 (71%)	0.90	43 (19%) 1 1	35, 74, 99, 100	0
All	All	1717/2116 (81%)	0.52	219 (12%) 3 3	20, 55, 99, 100	0

The worst 5 of 219 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	160	ALA	12.2
1	G	2	GLY	9.7
1	G	160	ALA	9.7
1	G	65	HIS	7.2
2	H	262	GLY	7.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	A	211	5/5	0.99	0.10	43,44,47,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.