



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2024 – 08:27 AM EDT

PDB ID : 4HZ8
Title : Crystal structure of BglB with natural substrate
Authors : Hwang, K.Y.; Nam, K.H.
Deposited on : 2012-11-14
Resolution : 1.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1492 (1.18-1.10)
Clashscore	141614	1537 (1.18-1.10)
Ramachandran outliers	138981	1483 (1.18-1.10)
Sidechain outliers	138945	1480 (1.18-1.10)
RSRZ outliers	127900	1464 (1.18-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4011 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-glucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	439	3513	2243	610	648	12	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASN	GLU	engineered mutation	UNP Q0GMU3
A	40	VAL	LEU	engineered mutation	UNP Q0GMU3
A	41	LYS	GLN	engineered mutation	UNP Q0GMU3
A	42	LYS	PRO	engineered mutation	UNP Q0GMU3
A	45	GLU	LYS	engineered mutation	UNP Q0GMU3
A	477	ASN	ARG	engineered mutation	UNP Q0GMU3
A	481	GLU	ALA	engineered mutation	UNP Q0GMU3
A	482	ASP	ALA	engineered mutation	UNP Q0GMU3

- Molecule 2 is beta-D-glucopyranose (three-letter code: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	486	Total	O	0	0
			486	486		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-glucosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.38Å 70.56Å 86.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.80 – 1.14 19.83 – 1.14	Depositor EDS
% Data completeness (in resolution range)	90.5 (19.80-1.14) 90.4 (19.83-1.14)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.14Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.135 , 0.161 0.134 , 0.160	Depositor DCC
R_{free} test set	7152 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	9.3	Xtrriage
Anisotropy	0.127	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.010 for k,h,-l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4011	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.18	13/3616 (0.4%)	1.24	34/4920 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	315	GLU	CB-CG	-9.07	1.34	1.52
1	A	393	ASP	CB-CG	8.98	1.70	1.51
1	A	175	GLU	CD-OE2	8.12	1.34	1.25
1	A	429	ARG	CZ-NH2	7.52	1.42	1.33
1	A	212	TYR	CD1-CE1	7.29	1.50	1.39
1	A	393	ASP	CG-OD2	6.68	1.40	1.25
1	A	69	GLU	CD-OE1	5.78	1.32	1.25
1	A	203	GLU	CD-OE1	5.61	1.31	1.25
1	A	429	ARG	NE-CZ	5.51	1.40	1.33
1	A	337	VAL	CA-CB	5.43	1.66	1.54
1	A	282	SER	CB-OG	-5.30	1.35	1.42
1	A	175	GLU	CG-CD	5.24	1.59	1.51
1	A	45	GLU	CB-CG	-5.02	1.42	1.52

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	429	ARG	NE-CZ-NH2	15.78	128.19	120.30
1	A	429	ARG	NE-CZ-NH1	-13.49	113.56	120.30
1	A	381	LEU	CB-CG-CD2	11.07	129.82	111.00
1	A	393	ASP	CB-CG-OD2	10.21	127.49	118.30
1	A	381	LEU	CB-CG-CD1	-9.46	94.92	111.00
1	A	246	ARG	NE-CZ-NH1	9.40	125.00	120.30
1	A	336	ARG	NE-CZ-NH1	9.13	124.86	120.30
1	A	136	ASP	CB-CG-OD2	-8.39	110.75	118.30
1	A	381	LEU	CD1-CG-CD2	-8.33	85.50	110.50
1	A	139	ARG	NE-CZ-NH2	8.22	124.41	120.30
1	A	77	ARG	NE-CZ-NH2	8.18	124.39	120.30
1	A	74	ARG	NE-CZ-NH1	-7.91	116.34	120.30
1	A	429	ARG	CD-NE-CZ	7.82	134.55	123.60
1	A	407	ARG	NE-CZ-NH1	7.56	124.08	120.30
1	A	420	ARG	NE-CZ-NH1	7.33	123.97	120.30
1	A	336	ARG	NE-CZ-NH2	-7.25	116.68	120.30
1	A	74	ARG	NE-CZ-NH2	7.14	123.87	120.30
1	A	246	ARG	NE-CZ-NH2	-6.92	116.84	120.30
1	A	397	ASP	CB-CG-OD1	6.78	124.41	118.30
1	A	419	ARG	NE-CZ-NH1	6.51	123.56	120.30
1	A	422	LEU	CB-CG-CD2	-6.49	99.97	111.00
1	A	69	GLU	OE1-CD-OE2	6.07	130.58	123.30
1	A	128	ARG	NE-CZ-NH1	-5.90	117.35	120.30
1	A	133	ARG	NE-CZ-NH2	5.76	123.18	120.30
1	A	74	ARG	CG-CD-NE	-5.64	99.95	111.80
1	A	77	ARG	NE-CZ-NH1	-5.60	117.50	120.30
1	A	140	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	278	ARG	NE-CZ-NH1	5.26	122.93	120.30
1	A	420	ARG	NE-CZ-NH2	-5.26	117.67	120.30
1	A	462	ARG	NE-CZ-NH1	5.26	122.93	120.30
1	A	279	ARG	CB-CG-CD	5.21	125.14	111.60
1	A	83	LYS	CD-CE-NZ	-5.08	100.03	111.70
1	A	212	TYR	CG-CD1-CE1	-5.04	117.27	121.30
1	A	139	ARG	NH1-CZ-NH2	-5.02	113.88	119.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	393	ASP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3513	0	3361	38	0
2	A	12	0	11	1	0
3	A	486	0	0	10	1
All	All	4011	0	3372	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:ARG:NH2	3:A:830:HOH:O	1.98	0.94
1:A:131:ASN:HD22	1:A:134:GLY:H	1.35	0.73
1:A:278:ARG:HH11	1:A:278:ARG:CG	2.03	0.71
1:A:278:ARG:HH11	1:A:278:ARG:HG2	1.58	0.68
1:A:439:ASN:ND2	3:A:1030:HOH:O	2.25	0.68
1:A:262:ASN:HD22	1:A:284:GLN:HE21	1.43	0.66
1:A:99:GLU:OE1	3:A:1085:HOH:O	2.14	0.65
1:A:387:GLU:OE2	2:A:501:BGC:H1	1.97	0.64
1:A:315:GLU:HG2	3:A:910:HOH:O	1.99	0.63
1:A:280:MET:HE2	1:A:335:MET:HE3	1.80	0.62
1:A:129:GLN:NE2	3:A:1058:HOH:O	2.33	0.61
1:A:275:GLU:HG2	1:A:279:ARG:HD3	1.90	0.54
1:A:337:VAL:HG13	1:A:346:ILE:HG23	1.89	0.53
1:A:280:MET:HE2	1:A:335:MET:CE	2.41	0.50
1:A:219:PRO:HG2	1:A:221:LEU:HD13	1.94	0.50
1:A:262:ASN:ND2	1:A:284:GLN:HE21	2.09	0.49
1:A:74:ARG:HD3	1:A:77:ARG:NH1	2.28	0.49
1:A:201:HIS:HE1	1:A:237:SER:OG	1.97	0.48
1:A:131:ASN:ND2	1:A:134:GLY:H	2.06	0.47
1:A:74:ARG:CD	1:A:77:ARG:NH1	2.77	0.46
1:A:128:ARG:NH2	3:A:985:HOH:O	1.91	0.45
1:A:100:GLN:NE2	3:A:736:HOH:O	2.47	0.45
1:A:280:MET:CE	1:A:335:MET:CE	2.96	0.44
1:A:44:PRO:HD3	1:A:422:LEU:HD21	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:LEU:HG	1:A:196:PRO:HG2	2.00	0.43
1:A:128:ARG:NE	3:A:849:HOH:O	2.25	0.43
1:A:74:ARG:NH2	1:A:74:ARG:CG	2.81	0.42
1:A:86:ASP:OD1	3:A:1082:HOH:O	2.22	0.42
1:A:113:TYR:CE2	1:A:115:PHE:HB3	2.55	0.42
1:A:343:PRO:HA	1:A:344:PRO:HA	1.81	0.41
1:A:401:GLN:OE1	3:A:912:HOH:O	2.22	0.41
1:A:335:MET:HB3	1:A:335:MET:HE2	1.65	0.41
1:A:381:LEU:CD2	1:A:381:LEU:H	2.33	0.41
1:A:278:ARG:CG	1:A:278:ARG:NH1	2.71	0.41
1:A:74:ARG:NH2	1:A:74:ARG:HG3	2.33	0.41
1:A:280:MET:CE	1:A:335:MET:HE3	2.49	0.41
1:A:340:SER:O	1:A:345:GLY:HA2	2.21	0.41
1:A:74:ARG:HH21	1:A:74:ARG:HB3	1.85	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:891:HOH:O	3:A:900:HOH:O[4_545]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	437/444 (98%)	430 (98%)	7 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	361/366 (99%)	355 (98%)	6 (2%)	60 22

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	GLU
1	A	74	ARG
1	A	278	ARG
1	A	279	ARG
1	A	307	ASN
1	A	381	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	129	GLN
1	A	131	ASN
1	A	132	GLN
1	A	163	GLN
1	A	201	HIS
1	A	262	ASN
1	A	296	GLN
1	A	307	ASN
1	A	342	GLN
1	A	388	ASN
1	A	461	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	A	501	-	12,12,12	1.51	2 (16%)	17,17,17	2.48	8 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	501	-	-	0/2/22/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	BGC	O1-C1	3.28	1.49	1.39
2	A	501	BGC	O6-C6	-2.79	1.30	1.42

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	BGC	O5-C1-C2	5.72	120.36	110.30
2	A	501	BGC	O1-C1-O5	-5.69	93.51	110.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	BGC	O5-C5-C4	-2.96	104.37	109.70
2	A	501	BGC	O2-C2-C1	2.38	114.73	109.25
2	A	501	BGC	C6-C5-C4	2.29	118.63	113.02
2	A	501	BGC	O2-C2-C3	2.25	115.67	110.38
2	A	501	BGC	O3-C3-C2	-2.19	105.21	110.38
2	A	501	BGC	C1-O5-C5	-2.02	109.74	113.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	BGC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/444 (98%)	-0.11	16 (3%) 42 42	5, 10, 22, 38	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	398	GLN	6.0
1	A	399	SER	4.5
1	A	343	PRO	4.4
1	A	350	GLN	4.0
1	A	352	GLU	3.4
1	A	209	TRP	3.4
1	A	303	MET	2.9
1	A	344	PRO	2.8
1	A	400	GLY	2.6
1	A	393	ASP	2.6
1	A	342	GLN	2.4
1	A	299	GLN	2.3
1	A	353	SER	2.3
1	A	45	GLU	2.3
1	A	224	PRO	2.2
1	A	314	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	A	501	12/12	0.94	0.16	9,13,21,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.