



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 12, 2024 – 08:06 AM EST

PDB ID : 2I2O  
Title : Crystal Structure of an eIF4G-like Protein from *Danio rerio*  
Authors : Bitto, E.; Wesenberg, G.E.; Phillips Jr., G.N.; Mccoy, J.G.; Bingman, C.A.;  
Center for Eukaryotic Structural Genomics (CESG)  
Deposited on : 2006-08-16  
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1028 (1.92-1.92)
Clashscore	180529	1100 (1.92-1.92)
Ramachandran outliers	177936	1087 (1.92-1.92)
Sidechain outliers	177891	1087 (1.92-1.92)
RSRZ outliers	164620	1028 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	224	
1	B	224	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called eIF4G-like protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	211	1739	1106	295	327	5	6	0	2	0
1	B	210	1737	1106	295	325	5	6	0	3	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ALA	-	cloning artifact	UNP Q5EAQ1
A	0	ILE	-	cloning artifact	UNP Q5EAQ1
A	1	ALA	-	cloning artifact	UNP Q5EAQ1
B	-1	ALA	-	cloning artifact	UNP Q5EAQ1
B	0	ILE	-	cloning artifact	UNP Q5EAQ1
B	1	ALA	-	cloning artifact	UNP Q5EAQ1

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ni	0	0
			1	1		
2	B	1	Total	Ni	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	202	Total	O	0	0
			202	202		
3	B	216	Total	O	0	0
			216	216		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: eIF4G-like protein



- Molecule 1: eIF4G-like protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	317.80Å 40.95Å 40.92Å 90.00° 90.26° 90.00°	Depositor
Resolution (Å)	40.61 – 1.92 40.61 – 1.92	Depositor EDS
% Data completeness (in resolution range)	97.4 (40.61-1.92) 97.4 (40.61-1.92)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.52 (at 1.92Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.190 , 0.235 0.247 , 0.284	Depositor DCC
$R_{free}$ test set	1997 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.0	Xtrriage
Anisotropy	0.044	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 31.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.032 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3896	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	8.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	0/1763	0.73	3/2364 (0.1%)
1	B	0.69	0/1761	0.68	0/2362
All	All	0.67	0/3524	0.71	3/4726 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	192	ARG	NE-CZ-NH2	-9.16	115.72	120.30
1	A	192	ARG	NE-CZ-NH1	8.22	124.41	120.30
1	A	192	ARG	CD-NE-CZ	5.06	130.68	123.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	118	ASN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1739	0	1753	32	0
1	B	1737	0	1755	29	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	202	0	0	4	0
3	B	216	0	0	3	0
All	All	3896	0	3508	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181[B]:PHE:CZ	1:B:181[B]:PHE:HE2	1.70	1.08
1:B:117:VAL:O	1:B:120:MSE:HB3	1.63	0.98
1:B:26:LEU:HD21	1:B:58:ILE:HG13	1.47	0.96
1:A:181[B]:PHE:CZ	1:B:181[B]:PHE:CE2	2.61	0.88
1:A:208:LEU:HD22	3:A:650:HOH:O	1.80	0.81
1:A:117:VAL:O	1:A:120:MSE:HB3	1.82	0.79
1:B:99:GLN:NE2	1:B:146:GLU:OE2	2.19	0.75
1:A:181[B]:PHE:CE1	1:B:181[B]:PHE:HE2	2.09	0.71
1:B:119:ASN:O	1:B:121:PRO:HD3	1.95	0.67
1:B:43:ASP:OD1	1:B:82:ARG:NH2	2.22	0.66
1:A:181[B]:PHE:CE1	1:B:181[B]:PHE:CE2	2.82	0.66
1:A:167:VAL:HB	3:A:531:HOH:O	1.96	0.66
1:A:91:GLU:HG2	3:A:540:HOH:O	1.96	0.65
1:A:99:GLN:NE2	1:A:146:GLU:OE2	2.31	0.64
1:A:119:ASN:O	1:A:121:PRO:HD3	2.00	0.62
1:B:111:ILE:HG23	1:B:115:LEU:HD22	1.82	0.60
1:A:118:ASN:CG	1:A:119:ASN:H	2.04	0.60
1:B:140:ASP:HA	1:B:143:LYS:HD3	1.84	0.59
1:A:123:VAL:HG13	3:A:534:HOH:O	2.02	0.59
1:A:117:VAL:HB	1:A:120:MSE:HE2	1.86	0.57
1:A:42:VAL:HG11	1:A:82:ARG:HG3	1.89	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:181[A]:PHE:HE1	1:B:196:LEU:CA	2.22	0.53
1:A:126:VAL:HG21	1:A:165[B]:MSE:SE	2.60	0.51
1:A:181[B]:PHE:HD2	1:A:199:LEU:HD12	1.75	0.51
1:B:214:LYS:HG3	3:B:658:HOH:O	2.11	0.51
1:B:126:VAL:HG21	1:B:165[A]:MSE:HG2	1.93	0.50
1:A:26:LEU:HD21	1:A:58:ILE:HG13	1.93	0.49
1:A:116:LYS:HE2	1:A:121:PRO:HG3	1.95	0.49
1:A:117:VAL:O	1:A:118:ASN:HB3	2.12	0.49
1:A:181[A]:PHE:HE1	1:B:181[A]:PHE:CZ	2.31	0.48
1:A:23:LYS:HA	1:A:26:LEU:HD12	1.94	0.48
1:A:117:VAL:HB	1:A:120:MSE:CE	2.44	0.48
1:B:118:ASN:HB3	1:B:120:MSE:H	1.77	0.48
1:A:126:VAL:CG2	1:A:165[B]:MSE:HG2	2.44	0.47
1:B:126:VAL:CG1	1:B:165[A]:MSE:HG2	2.44	0.47
1:B:126:VAL:HG11	1:B:165[A]:MSE:HG2	1.96	0.47
1:B:126:VAL:CG2	1:B:165[A]:MSE:HG2	2.44	0.47
1:B:42:VAL:HG11	1:B:82:ARG:HG3	1.98	0.46
1:A:126:VAL:HG21	1:A:165[B]:MSE:HG2	1.98	0.46
1:B:185:GLU:CD	3:B:614:HOH:O	2.54	0.46
1:A:181[B]:PHE:CE1	1:A:192:ARG:HG3	2.51	0.45
1:A:73:SER:OG	1:A:77:ARG:NH1	2.49	0.45
1:B:126:VAL:HG21	1:B:165[A]:MSE:SE	2.67	0.44
1:A:118:ASN:CG	1:A:119:ASN:N	2.71	0.43
1:A:181[B]:PHE:HE2	1:A:196:LEU:CA	2.33	0.42
1:B:117:VAL:CG2	1:B:120:MSE:HE3	2.50	0.41
1:B:161:GLN:O	1:B:165[B]:MSE:HG2	2.20	0.41
1:A:60:TYR:O	1:A:64:GLN:HG2	2.20	0.41
1:B:165[B]:MSE:HE3	3:B:521:HOH:O	2.19	0.41
1:B:181[A]:PHE:HD1	1:B:199:LEU:HD12	1.85	0.41
1:A:181[B]:PHE:CE2	1:A:196:LEU:HB2	2.55	0.41
1:A:69:GLN:HG2	1:A:70:THR:HG23	2.04	0.40
1:B:116:LYS:HE2	1:B:121:PRO:HG3	2.04	0.40
1:B:187:LEU:HB2	1:B:192:ARG:NH1	2.36	0.40
1:B:47:LYS:HA	1:B:47:LYS:HD3	1.93	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/224 (94%)	204 (97%)	6 (3%)	1 (0%)	25	13
1	B	211/224 (94%)	206 (98%)	4 (2%)	1 (0%)	25	13
All	All	422/448 (94%)	410 (97%)	10 (2%)	2 (0%)	25	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	118	ASN
1	B	118	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/199 (98%)	192 (98%)	3 (2%)	60	49
1	B	195/199 (98%)	191 (98%)	4 (2%)	48	34
All	All	390/398 (98%)	383 (98%)	7 (2%)	58	42

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	ASP
1	A	27	LYS
1	A	192	ARG
1	B	117	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	143	LYS
1	B	181[A]	PHE
1	B	181[B]	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

**Warning:** The R factor obtained from EDS is 0.2486, which does not match the depositor's R factor of 0.19. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	206/224 (91%)	0.45	10 (4%) 36 42	3, 7, 13, 19	1 (0%)
1	B	205/224 (91%)	0.68	27 (13%) 8 10	2, 7, 13, 19	3 (1%)
All	All	411/448 (91%)	0.57	37 (9%) 17 21	2, 7, 13, 19	4 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	118	ASN	5.9
1	A	119	ASN	5.4
1	B	119	ASN	5.1
1	B	118	ASN	4.4
1	B	16	LEU	3.9
1	B	8	ASP	3.7
1	B	28	ASP	3.4
1	B	31	SER	3.3
1	A	8	ASP	3.2
1	B	69	GLN	3.1
1	A	117	VAL	3.0
1	B	67	ALA	2.9
1	B	183	LEU	2.9
1	B	68	LYS	2.9
1	B	17	GLU	2.8
1	A	183	LEU	2.8
1	A	17	GLU	2.7
1	B	12	GLN	2.7
1	A	15	ASP	2.6
1	B	92	GLU	2.6
1	B	61	THR	2.6
1	B	24	THR	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	71	ASN	2.5
1	B	71	ASN	2.5
1	B	182	LEU	2.4
1	B	14	PHE	2.4
1	B	73	SER	2.4
1	B	15	ASP	2.3
1	B	60	TYR	2.3
1	B	57	ARG	2.2
1	A	31	SER	2.1
1	B	64	GLN	2.1
1	B	20	LYS	2.1
1	B	27	LYS	2.1
1	B	26	LEU	2.0
1	A	7	GLU	2.0
1	B	65	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q < 0.9
2	NI	A	502	1/1	0.93	0.08	20,20,20,20	1
2	NI	B	501	1/1	0.95	0.06	17,17,17,17	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.