



# wwPDB EM Validation Summary Report ⓘ

Apr 28, 2024 – 02:54 AM JST

PDB ID : 8I4M  
EMDB ID : EMD-35175  
Title : Portal-tail complex structure of the Cyanophage P-SCSP1u  
Authors : Liu, H.; Dang, S.  
Deposited on : 2023-01-19  
Resolution : 3.81 Å (reported)  
Based on initial model : ?

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	S	565	
1	T	565	
1	U	565	
1	V	565	
1	W	565	
1	X	565	
1	Y	565	
1	Z	565	
1	a	565	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	b	565	14% 96%
1	c	565	14% 96%
1	d	565	14% 97%
2	A	806	98%
2	B	806	98%
2	C	806	98%
2	D	806	98%
2	E	806	98%
2	F	806	98%
3	G	200	96%
3	H	200	95%
3	I	200	96%
3	J	200	96%
3	K	200	97%
3	L	200	98%
3	q	200	97%
3	r	200	96%
3	s	200	98%
3	t	200	96%
3	u	200	97%
3	v	200	96%
4	M	1079	17% 82%
4	N	1079	18% 82%
4	O	1079	18% 82%
4	P	1079	17% 82%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	Q	1079	 18% 82%
4	R	1079	 18% 82%
4	e	1079	 17% 82%
4	f	1079	 18% 82%
4	g	1079	 17% 82%
4	h	1079	 17% 82%
4	i	1079	 17% 82%
4	j	1079	 18% 82%
4	k	1079	 18% 82%
4	l	1079	 17% 82%
4	m	1079	 18% 82%
4	n	1079	 17% 82%
4	o	1079	 18% 82%
4	p	1079	 18% 82%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 134400 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Portal protein(gp 16) of the cyanophage P-SCSP1u.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	S	564	4368	2714	769	864	21	0	0
1	T	564	4368	2714	769	864	21	0	0
1	U	564	4368	2714	769	864	21	0	0
1	V	564	4368	2714	769	864	21	0	0
1	W	564	4368	2714	769	864	21	0	0
1	X	564	4368	2714	769	864	21	0	0
1	Y	564	4368	2714	769	864	21	0	0
1	Z	564	4368	2714	769	864	21	0	0
1	a	564	4368	2714	769	864	21	0	0
1	b	564	4368	2714	769	864	21	0	0
1	c	564	4368	2714	769	864	21	0	0
1	d	564	4368	2714	769	864	21	0	0

- Molecule 2 is a protein called Nozzle protein(gp 23) of the cyanophage P-SCSP1u.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	A	806	6159	3899	1022	1220	18	0	0
2	B	806	6159	3899	1022	1220	18	0	0
2	C	806	6159	3899	1022	1220	18	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
2	D	806	Total	C	N	O	S	0	0
			6159	3899	1022	1220	18		
2	E	806	Total	C	N	O	S	0	0
			6159	3899	1022	1220	18		
2	F	806	Total	C	N	O	S	0	0
			6159	3899	1022	1220	18		

- Molecule 3 is a protein called Adaptor protein(gp22) of the cyanophage P-SCSP1u.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	H	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	I	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	J	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	K	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	L	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	q	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	r	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	s	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	t	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	u	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		
3	v	198	Total	C	N	O	S	0	0
			1582	980	279	320	3		

- Molecule 4 is a protein called Fiber protein(gp 28) of the cyanophage P-SCSP1u.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	e	192	Total	C	N	O	S	0	0
			1447	898	244	304	1		
4	f	192	Total	C	N	O	S	0	0
			1447	898	244	304	1		

*Continued on next page...*




*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
4	g	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	h	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	j	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	l	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	i	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	k	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	m	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	n	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	o	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	p	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	M	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	N	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	O	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	P	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	Q	192	Total 1447	C 898	N 244	O 304	S 1	0	0
4	R	192	Total 1447	C 898	N 244	O 304	S 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Portal protein(gp 16) of the cyanophage P-SCSP1u









- Molecule 1: Portal protein(gp 16) of the cyanophage P-SCSP1u

Chain V: 14% 96%




- Molecule 1: Portal protein(gp 16) of the cyanophage P-SCSP1u


Chain W: 14% 96%




- Molecule 1: Portal protein(gp 16) of the cyanophage P-SCSP1u

Chain X: 14% 97%







• Molecule 1: Portal protein(gp 16) of the cyanophage P-SCSP1u




• Molecule 1: Portal protein(gp 16) of the cyanophage P-SCSP1u




• Molecule 1: Portal protein(gp 16) of the cyanophage P-SCSP1u




• Molecule 1: Portal protein(gp 16) of the cyanophage P-SCSP1u




- Molecule 1: Portal protein(gp 16) of the cyanophage P-SCSP1u




- Molecule 1: Portal protein(gp 16) of the cyanophage P-SCSP1u




- Molecule 2: Nozzle protein(gp 23) of the cyanophage P-SCSP1u




- Molecule 2: Nozzle protein(gp 23) of the cyanophage P-SCSP1u




- Molecule 2: Nozzle protein(gp 23) of the cyanophage P-SCSP1u




- Molecule 2: Nozzle protein(gp 23) of the cyanophage P-SCSP1u




- Molecule 2: Nozzle protein(gp 23) of the cyanophage P-SCSP1u



- Molecule 2: Nozzle protein(gp 23) of the cyanophage P-SCSP1u




- Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u



- Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u





- Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u

Chain I: 96%



- Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u

Chain J: 96%



- Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u

Chain K: 97%



- Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u

Chain L: 98%



- Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u

Chain q: 97%



- Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u

Chain r: 96%




- Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u

Chain s:  98% ..



• Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u

Chain t:  96% ..




• Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u

Chain u:  97% ..




• Molecule 3: Adaptor protein(gp22) of the cyanophage P-SCSP1u

Chain v:  96% ..



• Molecule 4: Fiber protein(gp 28) of the cyanophage P-SCSP1u

Chain e:  17%  82%



ALA ASP ALA SER ILE TYR THR LYS ILE GLN ASN VAL SER ASP ASP THR ARG ILE LEU

VAL GLU SER ALA SER ILE ASP ASN VAL VAL HIS THR LYS ARG ILE LEU VAL THR ASP VAL THR ASP THR THR PHE THR SER

ASN LEU ASP THR VAL PRO THR GLN ASN THR LYS THR ASN THR VAL VAL VAL THR THR THR THR THR THR THR THR THR THR THR

ALA LYS GLU THR VAL ILE THR ASN THR ASN THR LYS PHE PRO THR THR VAL ALA ALA THR VAL VAL ASN ASN PHE VAL THR VAL THR

LEU VAL ILE ASN VAL ASN GLY LEU SER ASP ASN THR VAL GLY THR VAL THR THR THR THR THR THR THR THR THR THR THR THR

LYS GLU ASP VAL ASN GLN LEU SER ASP ASN THR PHE ASN THR VAL THR THR THR THR THR THR THR THR THR THR THR THR THR THR












PHE  
LEU  
ILE  
ILE  
ASN  
LYS  
LYS  
TYR  
PHE  
ASN  
LEU  
THR  
GLU  
VAL  
SER

● Molecule 4: Fiber protein(gp 28) of the cyanophage P-SCSP1u

Chain 1:

17% 82%



ILE  
LEU  
GLY  
ARG  
ASP  
SER  
GLY  
ALA  
VAL  
ILE  
GLY  
GLU  
ILE

ASN  
ALA  
ILE  
GLU  
GLU  
SER  
SER  
ALA  
ASP  
VAL  
THR  
VAL  
VAL  
SER  
ASN  
VAL  
GLY  
THR  
ASN

THR  
SER  
GLN  
PRO  
LEU  
ASP  
SER  
GLU  
SER  
LEU  
SER  
GLY  
ALA  
MET  
THR  
SER

VAL  
VAL  
ASN  
PHE  
VAL  
VAL  
ALA  
ASP  
ASN  
GLN  
ILE  
PRO  
ALA  
PRO  
VAL  
GLY  
ILE

SER  
ASP  
ASN  
VAL  
THR  
ILE  
LYS  
ASN  
PHE  
PRO  
ALA  
SER  
LEU  
ASN  
ARG  
VAL  
THR

TYR  
ARG  
VAL  
GLU  
ASN  
THR  
LEU  
PRO  
ALA  
ASN  
ASN  
SER  
SER  
ASN  
SER  
HIS  
TYR

ALA  
PHE  
GLY  
SER  
LEU  
GLN  
ASP  
PHE  
ILE  
ASN  
THR  
PRO  
ALA  
SER  
ILE  
ASN  
ASN

ASP  
TYR  
PHE  
ALA  
ILE  
VAL  
SER  
ILE  
GLY  
THR  
ASN  
PRO  
SER  
ASN  
PRO  
THR  
SER  
LYS

GLY  
ASP  
GLY  
SER  
ASP  
GLY  
TYR  
LEU  
THR  
GLN  
LEU  
VAL  
ASN  
CYS  
SER  
SER  
GLN  
ILE

VAL  
THR  
ASP  
VAL  
ASN  
THR  
SER  
PRO  
GLN  
LEU  
GLY  
GLY  
ASN  
CYS  
ASN  
ASP  
LYS  
VAL

ASP  
LEU  
HIS  
LEU  
ARG  
LEU  
ASN  
GLY  
SER  
GLU  
GLY  
ILE  
VAL  
ASN  
ASN  
ALA  
VAL  
SER

MET  
GLY  
SER  
GLY  
ASN  
ALA  
GLN  
LEU  
HIS  
GLY  
SER  
ASN  
HIS  
GLN  
TYR  
PHE  
GLY  
ASN

VAL  
THR  
VAL  
SER  
GLY  
SER  
VAL  
THR  
HIS  
HIS  
PHE  
SER  
THR  
SER  
SER  
PRO  
ARG  
GLY


GLY  
HIS  
LEU  
ARG  
PRO  
TYR  
ALA  
SER  
ASN  
ASN  
THR  
HIS  
TRP  
ARG  
ARG  
ASN  
VAL  
GLY  
GLY  
SER

ARG  
THR  
SER  
LYS  
TYR  
LYS  
PHE  
ASN  
LEU  
THR  
GLU  
VAL  
SER


● Molecule 4: Fiber protein(gp 28) of the cyanophage P-SCSP1u

Chain i:

17% 82%



● Molecule 4: Fiber protein(gp 28) of the cyanophage P-SCSP1u

























## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	86469	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50.0	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.099	Depositor
Minimum map value	-0.051	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.01	Depositor
Map size (Å)	435.2, 435.2, 435.2	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.7, 1.7, 1.7	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	S	0.26	0/4428	0.56	2/5983 (0.0%)
1	T	0.27	0/4428	0.57	3/5983 (0.1%)
1	U	0.26	0/4428	0.57	4/5983 (0.1%)
1	V	0.27	0/4428	0.57	2/5983 (0.0%)
1	W	0.27	0/4428	0.57	3/5983 (0.1%)
1	X	0.26	0/4428	0.55	2/5983 (0.0%)
1	Y	0.26	0/4428	0.54	1/5983 (0.0%)
1	Z	0.26	0/4428	0.56	4/5983 (0.1%)
1	a	0.27	0/4428	0.57	3/5983 (0.1%)
1	b	0.27	0/4428	0.57	2/5983 (0.0%)
1	c	0.27	0/4428	0.58	3/5983 (0.1%)
1	d	0.26	0/4428	0.55	2/5983 (0.0%)
2	A	0.27	0/6293	0.52	1/8571 (0.0%)
2	B	0.27	0/6293	0.52	1/8571 (0.0%)
2	C	0.27	0/6293	0.52	0/8571
2	D	0.27	0/6293	0.53	1/8571 (0.0%)
2	E	0.27	0/6293	0.53	1/8571 (0.0%)
2	F	0.27	0/6293	0.53	1/8571 (0.0%)
3	G	0.27	0/1603	0.51	0/2171
3	H	0.29	0/1603	0.51	0/2171
3	I	0.27	0/1603	0.54	1/2171 (0.0%)
3	J	0.27	0/1603	0.51	0/2171
3	K	0.27	0/1603	0.52	0/2171
3	L	0.27	0/1603	0.51	0/2171
3	q	0.27	0/1603	0.51	0/2171
3	r	0.28	0/1603	0.51	0/2171
3	s	0.27	0/1603	0.53	1/2171 (0.0%)
3	t	0.28	0/1603	0.52	0/2171
3	u	0.27	0/1603	0.53	0/2171
3	v	0.27	0/1603	0.51	0/2171
4	M	0.26	0/1462	0.50	0/1985
4	N	0.26	0/1462	0.48	0/1985
4	O	0.26	0/1462	0.49	0/1985
4	P	0.27	0/1462	0.52	0/1985



Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
4	Q	0.26	0/1462	0.48	0/1985
4	R	0.27	0/1462	0.51	0/1985
4	e	0.27	0/1462	0.50	0/1985
4	f	0.26	0/1462	0.48	0/1985
4	g	0.28	0/1462	0.50	0/1985
4	h	0.26	0/1462	0.49	0/1985
4	i	0.27	0/1462	0.51	0/1985
4	j	0.26	0/1462	0.50	0/1985
4	k	0.26	0/1462	0.49	0/1985
4	l	0.26	0/1462	0.49	0/1985
4	m	0.27	0/1462	0.51	0/1985
4	n	0.26	0/1462	0.50	0/1985
4	o	0.26	0/1462	0.48	0/1985
4	p	0.27	0/1462	0.50	0/1985
All	All	0.27	0/136446	0.53	38/185004 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	U	0	1
1	V	0	1
1	Z	0	1
1	b	0	1
All	All	0	4

There are no bond length outliers.

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	W	464	PRO	CA-N-CD	-10.90	96.24	111.50
1	c	464	PRO	CA-N-CD	-10.90	96.24	111.50
1	a	464	PRO	CA-N-CD	-7.46	101.05	111.50
1	T	464	PRO	CA-N-CD	-7.43	101.10	111.50
1	Y	360	MET	CA-CB-CG	6.34	124.07	113.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	U	474	GLU	Peptide
1	V	474	GLU	Peptide
1	Z	474	GLU	Peptide
1	b	474	GLU	Peptide

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	562/565 (100%)	525 (93%)	35 (6%)	2 (0%)	34	70
1	T	562/565 (100%)	527 (94%)	33 (6%)	2 (0%)	34	70
1	U	562/565 (100%)	528 (94%)	30 (5%)	4 (1%)	22	59
1	V	562/565 (100%)	530 (94%)	30 (5%)	2 (0%)	34	70
1	W	562/565 (100%)	529 (94%)	30 (5%)	3 (0%)	29	66
1	X	562/565 (100%)	525 (93%)	34 (6%)	3 (0%)	29	66
1	Y	562/565 (100%)	526 (94%)	33 (6%)	3 (0%)	29	66
1	Z	562/565 (100%)	528 (94%)	30 (5%)	4 (1%)	22	59
1	a	562/565 (100%)	527 (94%)	33 (6%)	2 (0%)	34	70
1	b	562/565 (100%)	529 (94%)	31 (6%)	2 (0%)	34	70
1	c	562/565 (100%)	526 (94%)	33 (6%)	3 (0%)	29	66
1	d	562/565 (100%)	520 (92%)	40 (7%)	2 (0%)	34	70
2	A	804/806 (100%)	764 (95%)	39 (5%)	1 (0%)	51	83
2	B	804/806 (100%)	764 (95%)	39 (5%)	1 (0%)	51	83
2	C	804/806 (100%)	771 (96%)	33 (4%)	0	100	100
2	D	804/806 (100%)	763 (95%)	40 (5%)	1 (0%)	51	83

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	804/806 (100%)	765 (95%)	37 (5%)	2 (0%)	47	78
2	F	804/806 (100%)	768 (96%)	35 (4%)	1 (0%)	51	83
3	G	196/200 (98%)	189 (96%)	7 (4%)	0	100	100
3	H	196/200 (98%)	187 (95%)	9 (5%)	0	100	100
3	I	196/200 (98%)	188 (96%)	8 (4%)	0	100	100
3	J	196/200 (98%)	186 (95%)	10 (5%)	0	100	100
3	K	196/200 (98%)	192 (98%)	4 (2%)	0	100	100
3	L	196/200 (98%)	188 (96%)	8 (4%)	0	100	100
3	q	196/200 (98%)	192 (98%)	4 (2%)	0	100	100
3	r	196/200 (98%)	187 (95%)	9 (5%)	0	100	100
3	s	196/200 (98%)	186 (95%)	10 (5%)	0	100	100
3	t	196/200 (98%)	185 (94%)	11 (6%)	0	100	100
3	u	196/200 (98%)	189 (96%)	7 (4%)	0	100	100
3	v	196/200 (98%)	190 (97%)	6 (3%)	0	100	100
4	M	190/1079 (18%)	181 (95%)	9 (5%)	0	100	100
4	N	190/1079 (18%)	181 (95%)	9 (5%)	0	100	100
4	O	190/1079 (18%)	182 (96%)	8 (4%)	0	100	100
4	P	190/1079 (18%)	180 (95%)	10 (5%)	0	100	100
4	Q	190/1079 (18%)	183 (96%)	7 (4%)	0	100	100
4	R	190/1079 (18%)	181 (95%)	9 (5%)	0	100	100
4	e	190/1079 (18%)	177 (93%)	13 (7%)	0	100	100
4	f	190/1079 (18%)	181 (95%)	9 (5%)	0	100	100
4	g	190/1079 (18%)	180 (95%)	10 (5%)	0	100	100
4	h	190/1079 (18%)	178 (94%)	12 (6%)	0	100	100
4	i	190/1079 (18%)	179 (94%)	11 (6%)	0	100	100
4	j	190/1079 (18%)	179 (94%)	11 (6%)	0	100	100
4	k	190/1079 (18%)	182 (96%)	8 (4%)	0	100	100
4	l	190/1079 (18%)	184 (97%)	6 (3%)	0	100	100
4	m	190/1079 (18%)	184 (97%)	6 (3%)	0	100	100
4	n	190/1079 (18%)	181 (95%)	9 (5%)	0	100	100
4	o	190/1079 (18%)	182 (96%)	8 (4%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	p	190/1079 (18%)	181 (95%)	9 (5%)	0	100	100
All	All	17340/33438 (52%)	16430 (95%)	872 (5%)	38 (0%)	50	78

5 of 38 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	T	486	ILE
1	U	486	ILE
1	V	486	ILE
1	W	486	ILE
1	Y	486	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	S	466/467 (100%)	451 (97%)	15 (3%)	39	65
1	T	466/467 (100%)	452 (97%)	14 (3%)	41	66
1	U	466/467 (100%)	450 (97%)	16 (3%)	37	64
1	V	466/467 (100%)	447 (96%)	19 (4%)	30	59
1	W	466/467 (100%)	450 (97%)	16 (3%)	37	64
1	X	466/467 (100%)	453 (97%)	13 (3%)	43	67
1	Y	466/467 (100%)	452 (97%)	14 (3%)	41	66
1	Z	466/467 (100%)	448 (96%)	18 (4%)	32	60
1	a	466/467 (100%)	451 (97%)	15 (3%)	39	65
1	b	466/467 (100%)	449 (96%)	17 (4%)	35	63
1	c	466/467 (100%)	452 (97%)	14 (3%)	41	66
1	d	466/467 (100%)	451 (97%)	15 (3%)	39	65
2	A	686/686 (100%)	671 (98%)	15 (2%)	52	72
2	B	686/686 (100%)	672 (98%)	14 (2%)	55	75
2	C	686/686 (100%)	668 (97%)	18 (3%)	46	69

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	686/686 (100%)	670 (98%)	16 (2%)	50	71
2	E	686/686 (100%)	674 (98%)	12 (2%)	60	78
2	F	686/686 (100%)	672 (98%)	14 (2%)	55	75
3	G	177/179 (99%)	172 (97%)	5 (3%)	43	67
3	H	177/179 (99%)	169 (96%)	8 (4%)	27	57
3	I	177/179 (99%)	172 (97%)	5 (3%)	43	67
3	J	177/179 (99%)	171 (97%)	6 (3%)	37	64
3	K	177/179 (99%)	173 (98%)	4 (2%)	50	71
3	L	177/179 (99%)	175 (99%)	2 (1%)	73	85
3	q	177/179 (99%)	173 (98%)	4 (2%)	50	71
3	r	177/179 (99%)	172 (97%)	5 (3%)	43	67
3	s	177/179 (99%)	175 (99%)	2 (1%)	73	85
3	t	177/179 (99%)	170 (96%)	7 (4%)	31	59
3	u	177/179 (99%)	173 (98%)	4 (2%)	50	71
3	v	177/179 (99%)	172 (97%)	5 (3%)	43	67
4	M	163/886 (18%)	157 (96%)	6 (4%)	34	61
4	N	163/886 (18%)	160 (98%)	3 (2%)	59	77
4	O	163/886 (18%)	160 (98%)	3 (2%)	59	77
4	P	163/886 (18%)	157 (96%)	6 (4%)	34	61
4	Q	163/886 (18%)	160 (98%)	3 (2%)	59	77
4	R	163/886 (18%)	160 (98%)	3 (2%)	59	77
4	e	163/886 (18%)	159 (98%)	4 (2%)	47	70
4	f	163/886 (18%)	161 (99%)	2 (1%)	71	84
4	g	163/886 (18%)	158 (97%)	5 (3%)	40	65
4	h	163/886 (18%)	156 (96%)	7 (4%)	29	58
4	i	163/886 (18%)	158 (97%)	5 (3%)	40	65
4	j	163/886 (18%)	161 (99%)	2 (1%)	71	84
4	k	163/886 (18%)	162 (99%)	1 (1%)	86	92
4	l	163/886 (18%)	159 (98%)	4 (2%)	47	70
4	m	163/886 (18%)	160 (98%)	3 (2%)	59	77
4	n	163/886 (18%)	158 (97%)	5 (3%)	40	65

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	o	163/886 (18%)	162 (99%)	1 (1%)	86	92
4	p	163/886 (18%)	162 (99%)	1 (1%)	86	92
All	All	14766/27816 (53%)	14370 (97%)	396 (3%)	48	68

5 of 396 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	788	CYS
3	H	180	ARG
2	D	254	HIS
2	E	792	SER
3	K	96	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
4	l	167	ASN
4	Q	146	GLN
4	i	35	ASN
4	M	167	ASN
2	C	791	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-35175. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.




## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


### 6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 143




Y Index: 143




Z Index: 95


### 6.3.2 Raw map



X Index: 143



Y Index: 113




Z Index: 141


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_35175\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 3611 nm<sup>3</sup>; this corresponds to an approximate mass of 3262 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum i




\*Reported resolution corresponds to spatial frequency of 0.262 Å<sup>-1</sup>



## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.262 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.81	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.09	4.67	4.16

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-35175 and PDB model 8I4M. Per-residue inclusion information can be found in section 3 on page 8.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).







































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 90% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary





























The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8740	 0.3940
A	 0.9720	 0.4630
B	 0.9710	 0.4640
C	 0.9700	 0.4640
D	 0.9720	 0.4640
E	 0.9710	 0.4640
F	 0.9700	 0.4630
G	 0.9560	 0.4740
H	 0.9540	 0.4650
I	 0.9580	 0.4770
J	 0.9480	 0.4610
K	 0.9520	 0.4730
L	 0.9540	 0.4650
M	 0.8180	 0.3470
N	 0.8810	 0.3710
O	 0.8900	 0.3970
P	 0.8230	 0.3470
Q	 0.8830	 0.3660
R	 0.8860	 0.3940
S	 0.7810	 0.3280
T	 0.7820	 0.3320
U	 0.7800	 0.3260
V	 0.7800	 0.3220
W	 0.7830	 0.3300
X	 0.7790	 0.3280
Y	 0.7810	 0.3330
Z	 0.7800	 0.3260
a	 0.7810	 0.3330
b	 0.7820	 0.3250
c	 0.7840	 0.3320
d	 0.7810	 0.3340
e	 0.8160	 0.3430
f	 0.8910	 0.3720
g	 0.8990	 0.3960
h	 0.8190	 0.3460



*Continued on next page...*

*Continued from previous page...*

Chain	Atom inclusion	Q-score
i	 0.8210	 0.3460
j	 0.8810	 0.3660
k	 0.8810	 0.3640
l	 0.8900	 0.4000
m	 0.8870	 0.3940
n	 0.8210	 0.3460
o	 0.8930	 0.3740
p	 0.8970	 0.3990
q	 0.9550	 0.4770
r	 0.9550	 0.4650
s	 0.9560	 0.4770
t	 0.9480	 0.4600
u	 0.9540	 0.4730
v	 0.9550	 0.4650